

## Water Requirements of Avocado in Israel. I Tree and Soil Parameters†

D. Kalmar<sup>A</sup> and E. Lahav<sup>B</sup>

<sup>A</sup> Division of Environmental Physiology and Irrigation, Agricultural Research Organization,  
Volcani Center, Bet Dagan, Israel.

<sup>B</sup> Division of Subtropical Horticulture, Agricultural Research Organization,  
Volcani Center.

† Contribution No. 116-E, 1976 Series, Agric. Res. Organ., Volcani Center.

### Abstract

The effect of different irrigation intervals on the avocado tree was tested during the years 1968-1974 in the northern coastal plain of Israel. The objectives of this experiment were: (a) to make measurements of soil and plant parameters to help understand mechanisms of response to irrigation; and (b) to establish the most suitable irrigation schedule for highest yields of export quality fruit.

The trial consisted of four irrigation treatments, at intervals of 7, 14, 21 and 28 days; the respective average annual water applications were 8890, 7450, 6680, and 5940 m<sup>3</sup>/ha. The cultivars Ettinger, Fuerte and Hass were tested in five replications in randomized blocks. Soil moisture consumption was estimated by using a neutron probe and tensiometers.

Most of the water consumption was from the upper 60-cm soil layer. With the 7-day interval, the calculated daily loss from the soil profile was greater throughout the whole irrigation season than with the other intervals.

The least salt accumulation occurred under the long interval treatments since the large amounts of water supplied at each irrigation leached the salts continuously.

Reduced intervals between irrigations resulted in increased trunk growth and increased tree size. It is concluded that with a 21-day interval the vegetative growth of producing trees was restricted and therefore relatively smaller trees with easier fruit picking and delayed orchard thinning were obtained.

### Introduction

Most avocado plantations in the northern coastal plain of Israel are planted jointly with, or are incorporated into, existing banana plantations. For the first few years the avocado trees are irrigated according to the requirements of the banana plants. After the removal of the bananas, irrigation schedules are changed to those which are optimal for avocados.

Very little research has been carried out in Israel or elsewhere on the water requirements of avocado trees, and the little done was mostly on the effect of intervals between irrigations, as determined with the aid of tensiometers (Richards 1950; Richards *et al.* 1962). In a single irrigation experiment with cv. Hass in California, 31-46 irrigations per season resulted in increases in yields and fruit size as compared with 17-26 or 8-11 irrigations per season (Richards *et al.* 1958, 1962). The present paper reports on the first comprehensive irrigation trial, carried out with the aims of (a) measuring soil and plant parameters to help understand mechanisms of response to irrigation, and (b) determining the best irrigation schedule for achieving highest yields of export quality fruits (Lahav and Kalmar 1977).

## Methods

In 1963 an avocado orchard was planted together with bananas at the Akko Experiment Station in the northern coastal plain of Israel. The plantation was situated on a slope of 4‰. In the summer of 1966 the banana trees were removed and soon afterwards ridges were made for surface drainage during the rainy season.

The trial was initiated in 1968 and terminated in 1974. The average rainfall and evaporation for this period are summarized in Fig. 1. Sprinklers were placed equidistant between four avocado trees planted 6 by 6 m apart. Water quality was very good (50 ppm chloride) and constant throughout the irrigation season. The soil was a grumusol with above 60% clay in all layers down to 150 cm.




Fig. 1. Six years' average rainfall and class A pan evaporation (Akko Exp. Stn., 1968-1973).

The Ettinger, Fuerte and Hass cultivars were tested in five replications in randomized blocks. Each plot comprised nine trees and double guard rows (900 m<sup>2</sup>).

The trial consisted of four irrigation treatments, at intervals of 7, 14, 21 and 28 days. The respective average annual amounts of water applied were 8890, 7450, 6680, and 5940 m<sup>3</sup>/ha. In the first two treatments (7- and 14-day intervals) water was given to wet the soil to a depth of 30 cm; additional quantities were given to wet the soil down to 90 cm whenever this was found necessary according to soil moisture tests. For irrigation treatments at longer intervals, water quantities were fixed according to the requirements of the 0-60 cm soil layer; additional amounts were given whenever needed to keep the soil moist to a depth of 120 cm. In all treatments no excess water was given to the trees and therefore no underground drainage occurred.

The differential irrigation treatments were started in early June after the fruit had set, to avoid excessive fruit drop. Irrigation treatments were applied until the end of October.

### Soil and Water Tests

Moisture tests were carried out with a neutron probe before and after each irrigation. For each treatment measurements were made in one plot with 36 access tubes, one for each sq metre between four trees. In the other four replications, four access tubes were placed according to measurements of water distribution by the sprinklers (Christiansen 1942). The neutron probe was calibrated separately to

0–30 cm and 30–180 cm soil layers by means of parallel gravimetric measurements (Cohen 1964).

Six tensiometers were placed in a single plot of each treatment with the object of studying their use to determine irrigation needs. The depths of tensiometer placements were 30, 45, 90 and 120 cm in 7, 14, 21 and 28-day irrigation intervals respectively.

### *Soil Salinity*

The soil was analysed before the start of the experiment and then again every spring and autumn. Each plot was sampled and analysed separately. Soil layers (of 30 cm each) were sampled to a depth of 150 cm. Estimations of pH, exchangeable cations, sodium, chloride, potassium, and calcium+magnesium were made in the soil water extract. The sodium adsorption ratio was also calculated.

### *Tree Size*

Measurements were made each spring and autumn. The tree radius—from the trunk to the edge of the canopy—and the tree height were recorded. The tree volume was calculated according to Turrell (1946). The formulae used were: for cvv. Fuerte and Hass,  $V = \frac{2}{3}ab^2$ , and for cv. Ettinger  $V = \frac{4}{3}ab^2$ , where  $V$  is the tree volume,  $a$  the height and  $b$  the radius. Regression analysis for tree volume on the amount of water applied was made, and the deviation from linearity calculated.

The trunk circumference was measured 20 cm above the graft union. Trunk growth in winter and summer was calculated.

Daily trunk expansion and contraction were registered continuously with dendrographs sensitive to changes of 10  $\mu\text{m}$ . As the preliminary tests did not produce differences among cultivars but significant differences according to alternate bearing (Lahav *et al.* 1973–74, 1975), Hass trees only, in their 'on-year', were measured. Trees irrigated at 7-day intervals were compared with those irrigated every 28 days. During the 6 years of experimentation many trunks were measured. The results presented here can be regarded as typical, as the findings recorded were observed repeatedly in other trees and seasons.

## **Results**

### *Soil parameters*

Water extraction from the different layers is shown in Fig. 2. More water was consumed in 1972 than in 1968 because of tree growth, but the relative differences between treatments were unchanged.

The daily loss from the soil profile was affected by meteorological conditions and irrigation treatments. As temperature and evaporation increased and irrigation intervals decreased, the daily loss from the profile increased.

The relationship between the water deficit and the evaporation from a class A pan ( $E_t/E_o$ ) varied with the season, with a peak in August–September (Fig. 3). The irrigation interval was found to be negatively correlated with  $E_t/E_o$ , being high in the 7- and 14-day interval treatments compared with the other two treatments.

Most of the changes in relative water consumption occurred to a depth of 60 cm. Water extraction in 1972 was from deeper soil layers than in 1968. The contribution of the upper layers to the total water consumption decreased with increased irrigation







Fig. 2. Effect of irrigation interval on relative water consumption from the soil in two years.

Fig. 3. Effect of irrigation interval on the daily water deficit ( $E_t$ ) and the relationship between the water deficit and the evaporation from a class A pan ( $E_t/E_o$ ). Irrigation intervals:  $\circ$  7 days;  $\square$  14 days;  $\triangle$  21 days;  $\blacktriangle$  28 days.

Fig. 4. Effect of a 21-day irrigation interval on soil moisture tension at the 90-cm depth throughout the irrigation season.

intervals. Average seasonal irrigation efficiency (the ratio between the additional moisture in the soil and the amount of water supplied) was 94–96%, without differences among treatments.

The tensiometers showed the following effects of irrigation intervals on soil water tension:

- 7-day—the tensiometers never showed the maximal water tension (70 centibars) even when placed only 30 cm deep.
- 14-day—tensiometers located to a depth of 45 cm showed maximal tension after 12–14 days.
- 21-day—tensiometers 90 cm deep showed increasing tension before each irrigation, and during the irrigation season (Fig. 4).
- 28-day—deeper-located tensiometers (120 cm) did not respond to irrigation, but a continuous increase in tension was registered during the irrigation season.




Fig. 5. Effect of irrigation interval on salt accumulation in the soil profile (0–150 cm) from spring 1968 (dashed lines) until autumn 1974 (solid lines, left-hand side) and during the six experimental years (0–90 cm, right-hand side). Symbols for irrigation intervals as in Fig. 3.

The effect of irrigation interval and water quantity on soil salinity is shown in Fig. 5. During the 6 years of experimentation, pronounced salt accumulation occurred in all soil layers and in all treatments, up to 2.0 mmhos/cm (mostly below 90 cm). The treatment effect was found from the third year onward. In the upper layers, salt accumulation was higher with the 7-day interval than with the other treatments, while with the 28-day interval it was lowest. Generally, as irrigation intervals shortened and the amount of water per year consequently increased (though in each irrigation it decreased), more salts accumulated in the soil profile.

### Tree parameters

#### Tree Size

The dimensions of trees irrigated at 7-day intervals were largest, and those of trees irrigated at 28-day intervals were smallest (Table 1).

#### Trunk Circumference

As the trees became older and thicker, the annual trunk growth decreased. Among the three cultivars, the increase in circumference of cv. Ettinger was highest, while that of Fuerte was smallest (Fig. 6). As irrigation intervals were shortened, the rate of increase in circumference rose.

Table 1. Effect of irrigation interval on avocado tree volume  
Each figure is a mean for five plots

Cultivar	Irrigation interval (days)	Tree volume (m <sup>3</sup> )			Volume increase (%)
		1969	1974	Difference	
Ettinger	7	37.0	88.9	51.9	140.2
	14	35.0	87.4	52.4	149.7
	21	34.4	70.8	36.4	105.0
	28	32.2	64.5	32.3	100.0
	Average	34.7	77.9	43.3	124.8
Fuerte	7	21.1	43.5	22.4	106.2
	14	18.7	35.7	17.0	90.9
	21	19.7	35.0	15.3	77.7
	28	19.0	33.2	14.2	74.7
	Average	19.6	36.9	17.2	87.8
Hass	7	21.5	45.3	23.8	110.7
	14	18.0	39.9	21.9	121.7
	21	22.0	42.1	20.1	91.4
	28	17.5	34.6	17.1	97.7
	Average	19.8	40.5	20.7	104.5
Standard error of means		1.5	1.9	2.2	
Significance of linear component		NS	0.01	0.05	
Deviation from linearity		NS	NS	NS	

Differences were found between the effect of treatments on the winter increase in circumference compared with that of summer. Increase in trunk circumference in the summer was more rapid at shorter irrigation intervals, as mentioned above. In the winter the tendency was reversed, i.e. the 'dry' treatments showed higher growth rates. However, since summer growth was four times that of winter, the overall effect was not much changed.

Daily expansion and contraction of radius as measured by dendrometers showed remarkable differences between trees irrigated at 7-day and 28-day intervals (Fig. 7). The trunk contraction in particular was affected, being twice as high in trees irrigated at long intervals. Irrigation reduced trunk contraction in trees irrigated at long

intervals for 9–15 days, but for 3–4 days only in trees irrigated every week. In the 'dry' treatment, trunk contraction disappeared almost completely for the first day after irrigation. Yet the total increase in radius achieved in one 28-day cycle, as compared with four intervals of 7 days each, did not differ greatly.




Fig. 6. Effect of irrigation interval on the summer (*upper curves*) and winter (*lower curves*) growth of the trunk circumference.

Fig. 7. Differences in effect of irrigation on changes in the trunk radius between trees irrigated at 7-day intervals (*top*) and those irrigated at 28-day intervals (*bottom*).

## Discussion

On the basis of yield records presented in paper II (Lahav and Kalmar 1977), it was shown that the best treatment was that of the 21-day interval. In this treatment 600 m<sup>3</sup> water/ha was supplied at each irrigation during June and July. Since water losses increased later in the season, irrigation rates were then increased up to 800 m<sup>3</sup>/ha. It should be pointed out that in this treatment the average daily amount of water applied was only 32 m<sup>3</sup>/ha ( $E_d/E_o = 0.50$ ), and the seasonal amount was 6680 m<sup>3</sup>/ha. These values are only 75% of the corresponding 7-day interval values, which are normal for avocado plantations in Israel. The neutron probe measurements enabled water consumption in the whole tree area to be calculated down to a considerable depth. At the 7-day interval, the daily loss from the soil profile was higher



during the whole irrigation season than at other intervals (Fig. 3). This can be explained by the larger tree volume caused by frequent irrigations and the higher annual total amount of water applied.

It seems that with a 21-day interval a better wetting and drying regime is obtained. It has been established that an irrigation schedule which permits considerable drying of the heavy soil results in a better soil structure, in contrast to the destruction caused by frequent irrigations, which is in fact an irrigation regime without a drying period.\*

On the basis of the present findings, the relation between water deficit and the class A evaporation ( $E_i/E_o$ ) in avocado plantations in heavy soil can be reduced from 0.6 to 0.43 in June–August.

Under prevailing experimental conditions, tensiometers recorded the soil water tension for the 7- and 14-day irrigation intervals. It would appear that the range of tensiometers is not large enough to cover the changes in tension down to 60 cm in the longer irrigation intervals.

It is well known that under uniform soil and water conditions, the irrigation rate is the main factor affecting salt accumulation. In the present experiment the chloride content in the irrigation water was less than 50 mg/l, which is far below the concentration found to be damaging to avocado trees even when grafted on Mexican rootstocks (Kadman 1963). However, the annual amount of chlorides introduced with the irrigation water (400–600 kg/ha) was not negligible. Heavy winter rainfall during 1968–1971 leached out all the salts which had accumulated since the commencement of the experiment. Later, with reduced precipitation, a gradual increase in soil salinity was registered.

With the 7-day interval treatment, large amounts of water were supplied (6-year av., 8890 m<sup>3</sup>/ha). In consequence, this treatment caused larger amounts of salt to be introduced into the soil. This treatment, of light but frequent irrigations, reduced the leaching process, which led to an additional relative increase in salinity in the upper soil layer. Increased water consumption by the larger trees under this treatment may add to the already low leaching capacity of this irrigation regime. In an irrigation experiment conducted with Hass trees in California, frequent irrigations also resulted in increased chloride concentrations (Bingham and Richards 1958).

The lowest salt accumulations occurred with applications at 28-day intervals, since the large amounts of water (up to 1000 m<sup>3</sup>/ha per irrigation) leached the salts continuously. These findings should be of considerable importance in areas where avocado is grown under conditions of water shortage and high water salinity.

The average increase in tree volume during the 1968–1974 period was over 30 m<sup>3</sup> for trees irrigated at 7- and 14-day intervals, and 22 m<sup>3</sup> for trees irrigated at greater intervals. The response to the irrigation regime was highest with cv. Ettinger and lowest with cv. Fuerte (Fig. 8). Restricted growth as a result of a reduced water supply was also observed with Hass trees in California (Richards *et al.* 1962).

The trees were relatively young and not yet producing when the experiment was initiated (fifth year). Naturally, the effect of water stress on tree volume was more pronounced at this stage of growth. The main effect was on the number of flushes per season. There were three or four flushes on the trees irrigated at 7-day and 14-day intervals compared with only one or two flushes with the other two treatments.

\* Kalmar, D., Dynamics of grumusol structure. Ph.D Thesis, Hebrew Univ. Jerusalem (in Hebrew, in preparation).



The water effect on summer growth was reversed in autumn with the onset of the rains, but temperatures were still high. The autumn-accelerated growth of trees irrigated at 21-day and 28-day intervals was best observed in trunk circumference measurements, and in the daily trunk growth and contraction curves (Lahav *et al.* 1975).

Irrigation always caused an increase in circumference. The annual average difference in trunk radius between trees irrigated at 7-day and 28-day intervals was 1.4 mm (= 1.7% of trunk radius). A qualitative difference in trunk growth was observed 18 days after irrigation (28-day interval). This corresponds with the cessation of fruit growth (Lahav *et al.* 1975).




Fig. 8. Effect of amount of irrigation water on the relative efficiency of canopy growth during 6 years (1968-1974). Correlation coefficients: Ettinger, 0.499; Fuerte, 0.987\*\*; Hass, 0.587.

In conclusion, it can be stated that a well-planned irrigation program can be used to control avocado tree size. The growth of young trees can be accelerated and bearing thereby advanced, or the balance between the reproductive and vegetative phases may be changed by growth restriction. To ease fruit picking and delay orchard thinning, relatively small trees were obtained by irrigating every 21 days, with a total of 6680 m<sup>3</sup>/ha water per year. This treatment did not result in lower yields, and compared favourably with the other treatments tested (Lahav and Kalmar 1977).

## References

- Bingham, F. T., and Richards, S. J. (1958). Effects of irrigation treatments and rates of nitrogen fertilization on young Hass avocado trees. III. Changes in soil chemical properties. *Proc. Am. Soc. Hortic. Sci.* **71**, 304-9.
- Christiansen, J. E. (1942). Irrigation by sprinkling. Bull. Calif. Agric. Exp. Stn. No. 670.
- Cohen, O. P. (1964). A procedure for calibrating neutron moisture probes in the field. *Israel J. Agric. Res.* **14**, 169-78.
- Kadman, A. (1963). The uptake and accumulation of chloride in avocado leaves and the tolerance of avocado seedlings under saline conditions. *Proc. Am. Soc. Hortic. Sci.* **83**, 280-6.
- Lahav, E., and Kalmar, D. (1977). Water requirements of avocado in Israel. II. Influence on yield, fruit growth and oil content. *Aust. J. Agric. Res.* **28**, 869-77.
- Lahav, E., Kalmar, D., and Zamet, D. (1973-74). The value of physiological indicators in determining the irrigation requirements of the avocado tree. *Yearb. Calif. Avocado Soc.* **56**, 137-46.

- Lahav, E., Kalmar, D., and Zamet, D. (1975). Daily fluctuations in the growth of avocado trunks and fruits and the effects of irrigation and meteorological conditions on them. Pamph. Agric. Res. Organ., Volcani Center, No. 149. (Hebrew, with English summary).
- Richards, S. J. (1950). Soil moisture conditions in avocado groves. *Yearb. Calif. Avocado Soc.* 35, 70-2.
- Richards, S. J., Moore, P. W., Bingham, F. T., Embleton, T. W., and Labanauskas, C. K. (1958). Avocado irrigation and nitrogen fertilization plots at the Citrus Experiment Station. *Yearb. Calif. Avocado Soc.* 42, 25-9.
- Richards, S. J., Warneke, J. E., and Bingham, F. T. (1962). Avocado tree growth response to irrigation. *Yearb. Calif. Avocado Soc.* 46, 83-7.
- Turrell, F. M. (1946). 'Tables of Surfaces and of Prolate and Oblate Spheroids and Spheroidal Coefficients.' (Univ. of California Press: Berkeley.)

Manuscript received 21 January 1977

## **Water Requirements of Avocado in Israel. II\* Influence on Yield, Fruit Growth and Oil Content†**

*E. Lahav<sup>A</sup> and D. Kalmar<sup>B</sup>*

<sup>A</sup> Division of Subtropical Horticulture, Agricultural Research Organization,  
Volcani Center, Bet Dagan, Israel.

<sup>B</sup> Division of Environmental Physiology and Irrigation, Agricultural Research Organization,  
Volcani Center.

### *Abstract*

The effect of four different irrigation intervals, 7, 14, 21 and 28 days, on the avocado tree was tested during the years 1968-1974 in the northern coastal plain of Israel. The respective average annual water applications were 8890, 7450, 6680 and 5940 m<sup>3</sup>/ha. The cultivars Ettinger, Fuerte and Hass were tested in five replications in randomized blocks.

After 6 years no significant reduction in yield was observed with Ettinger or Fuerte trees irrigated once in 21 or 28 days. With the Hass cultivar the 28-day interval gave lower yields than with the three other intervals tested.

Shortening of irrigation intervals increased the growth rate and size of individual fruits, which may be of economic importance for cv. Hass, where overproduction leads to small fruits unsuitable for export. Shortening of irrigation intervals tended to increase the oil percentage of the fruit, which may advance the harvest date. In view of the equal yields obtained and the predominance of cv. Hass in avocado plantations, it was concluded that the 21-day interval was the optimum irrigation frequency under the experimental conditions.

### **Introduction**

An irrigation experiment on the avocado tree was conducted during 6 years in the northern coastal plain of Israel. The objectives were (a) to find soil or plant physiological indicators for ascertaining the water status and (b) to determine the best irrigation schedule for achieving highest yields of export quality fruit.

Soil moisture tests showed that most of the water consumed came from the upper 60 cm. When irrigation intervals were increased, the contribution of the uppermost soil layer to the overall water consumption lessened. It was likewise found that decreased irrigation intervals and increased water application rates resulted in increased accumulation of salts and increased growth rate of the tree (Kalmar and Lahav 1977).

### **Methods**

Four irrigation treatments were given, with intervals of 7, 14, 21 and 28 days between irrigations and respective average annual water applications of 8890, 7450, 6680 and 5940 m<sup>3</sup>/ha. The detailed layout of the experiment was given previously.

### *Evaluation of Yields and Fruit Size*

The fruits of each tree were counted and weighed.

\* Part I, *Aust. J. Agric. Res.*, 1977, **28**, 859.

† Contribution No. 117-E, 1976 Series, Agric. Res. Organ., Volcani Center.



### *Fruit Growth*

Experimentation was aimed at establishing parameters for the determination of irrigation timing for avocado plantations. Growth of avocado fruits in 1969–1970 was determined by measuring the distance between two metal discs glued to opposite sides of a fruit. The measuring gauge used was sensitive to changes of  $10\text{ }\mu\text{m}$ . Subsequently the data were recorded by means of fruitographs (Fig. 1) which recorded all stages of the daily and seasonal growth and shrinkage of fruits (Lahav and Kalmar 1972; Lahav *et al.* 1975).

Measurements were carried out in the Hass cultivar only (distinguished by small fruits). Fruits of trees irrigated at 7-day intervals were measured and compared with those irrigated at 28-day intervals. During the 6 years of experimentation, the growth pattern of many fruits was studied. The data presented here can be regarded as typical, as the findings recorded were observed repeatedly in other trees, fruits and seasons.




Fig. 1. Measuring growth of avocado fruit diameter by means of a fruitograph.

Significant differences were observed in the growth of fruits between trees in an 'on year' and those in an 'off year'. As problems of fruit size arise principally in years of abundant yields, the results reported here are only for trees of high productivity.

### *Oil Content*

In view of the importance of the oil percentage on harvest date, representative uniform-sized fruits of the Fuerte and Ettinger cultivars were analysed for oil content. Samples comprised five fruits of several sizes from each of the 20 plots, taken at harvest time. Analyses were made by the Halowax method, in which the refractive index is measured (Gazit and Spodheim 1969).

For all parameters regression analyses on the amount of water applied were made, and the deviation from linearity calculated.

## Results

### *Yields and grading*

Yields by weight and number of fruits are given in Tables 1 and 2 respectively. The Ettinger and Fuerte cultivars gave satisfactory yields. Yields decreased with increase in irrigation intervals, but only in one case (Ettinger in 1973-74) were significant differences found between treatments. The drop in yield was in most cases expressed only in the weight of the crop and not in the number of fruits, which remained nearly constant over all treatments.

Table 1. Effect of irrigation interval on the average yield of avocado

Each figure is the mean for five plots

Cultivar	Year	Yield (kg/tree)				SE of means	Signif. of linear component	Deviation from linearity
		Irrigation interval (days):						
		7	14	21	28			
<i>Ettinger</i>	1968-69	10.2	17.6	20.7	18.8	3.5	0.05	NS
	1969-70	19.5	24.4	18.3	16.4	3.8	NS	NS
	1970-71	73.6	73.0	66.5	63.1	5.0	NS	NS
	1971-72	49.8	45.0	45.0	54.0	3.2	NS	NS
	1972-73	53.0	42.9	49.9	47.9	3.4	NS	NS
	1973-74	92.5	79.4	76.7	72.4	4.9	0.01	NS
Av., 1970-71 to 1973-74		67.2	60.1	59.5	59.3	2.6	NS	NS
Av., 6 years		49.8	47.0	46.2	45.4	1.5	NS	NS
<i>Fuerte</i>	1968-69	7.3	4.7	11.3	6.0	2.7	NS	NS
	1969-70	41.9	32.2	30.1	28.6	4.4	NS	NS
	1970-71	52.9	56.9	47.8	46.9	4.9	NS	NS
	1971-72	45.2	37.4	40.4	43.4	2.9	NS	NS
	1972-73	52.7	66.1	61.2	50.4	4.4	NS	NS
	1973-74	50.9	43.6	46.5	50.6	3.2	NS	NS
Av., 1969-70 to 1973-74		48.7	47.2	45.2	44.0	1.6	NS	NS
Av., 6 years		41.8	40.1	39.5	37.6	1.6	NS	NS
<i>Hass</i>	1968-69	41.7	57.2	64.6	58.0	2.7	NS	NS
	1969-70	86.3	77.3	41.1	33.7	4.2	0.01	NS
	1970-71	32.5	64.1	95.0	97.0	5.2	0.01	NS
	1971-72	104.3	63.3	44.1	47.0	5.3	0.01	NS
	1972-73	18.1	39.5	72.4	40.2	4.4	0.05	NS
	1973-74	93.8	56.6	13.5	40.3	4.9	0.01	NS
Av., 1969-70 to 1973-74		67.0	60.2	53.2	51.6	2.2	0.05	NS
Av., 6 years		62.8	59.7	55.1	52.7	1.8	0.05	NS

The Hass cultivar gave the highest overall yield during the 6 years of the experiment. The treatments had a significant effect on the yield and on the number of fruits, starting in the second year of the experiment.

At the conclusion of the experiment, differences in yields between trees irrigated at 7-day and 28-day intervals were found to be about 10 kg/tree, which was significant.



### Effect of Irrigation Regime on Alternate Bearing

In the second year of the experiment, irrigation treatments applied at 21-day and 28-day intervals resulted in a 50% reduction of yields of cv. Hass from those irrigated at 7-day and 14-day intervals. These yield differences resulted in alternate bearing.

Table 2. Effect of irrigation interval on the number of avocado fruits per tree (1968-69 to 1973-74 average)

Each figure is the mean for five plots

Cultivar	No. of fruits per tree				SE of means	Signif. of linear component	Deviation from linearity
	Irrigation interval (days):						
	7	14	21	28			
Ettinger	100	106	99	104	6	NS	NS
Fuerte	125	116	124	118	5	NS	NS
Hass	276	286	255	254	10	0.05	NS

In Fig. 2, yields for cv. Hass are expressed as percentages of 'on year' trees (more than 50 kg/tree) during each of the six years, and in the year following (1974-75). Alternate bearing of trees irrigated at 7- and 14-day intervals was the opposite of that of trees irrigated at 21- and 28-day intervals. The 7-day interval trees tended more to alternate bearing than did trees irrigated at the other intervals.




Fig. 2. Effect of irrigation regime on the percentage of 'on year' Hass trees (> 50 kg/tree) during the six experimental years and the following year (1974). Irrigation intervals: ○ 7 days; □ 14 days; △ 21 days; ◆ 28 days.

### Fruit Size

In most cases, the largest-sized fruits occurred under the 7-day interval treatment and the smallest under the 28-day treatment (Table 3). Fruits of cvv. Ettinger and Hass were more affected than cv. Fuerte fruits. Whereas the average fruit size difference between trees irrigated at 7-day and 28-day intervals in cv. Fuerte was 9 g (about 3% of fruit weight), similar treatments with Hass trees produced an average difference of 18 g (9%), and with cv. Ettinger of 35 g (11%).

Contrary to expectations, fruits of Ettinger trees irrigated at 21-day intervals were about the same, and those of Hass relatively larger, than those irrigated at 14-day intervals. With cv. Hass this observation was recorded in five out of the six years of experimentation. In some cases (Fuerte in 1972-73, 1973-74), the fruit size was affected more by the yield than by the irrigation interval.



Table 3. Effect of irrigation interval on mean avocado fruit weight

Each figure is the mean for five plots

Cultivar	Year	Mean fruit weight (g)				SE of means	Signif. of linear component	Deviation from linearity
		Irrigation interval (days):						
		7	14	21	28			
Ettinger	1968-69	377.9	323.8	317.2	324.7	9.6	0.01	NS
	1969-70	303.3	311.2	321.0	287.8	5.0	0.01	NS
	1970-71	271.3	261.7	264.2	249.8	5.0	0.01	NS
	1971-72	293.6	285.6	269.8	245.5	6.1	0.01	NS
	1972-73	364.6	341.5	344.7	315.3	5.9	0.01	NS
	1973-74	341.4	326.9	330.7	321.7	5.1	0.05	NS
	Average	325.4	308.4	307.9	290.8	4.3	0.01	NS
Fuerte	1968-69	331.8	335.7	332.3	333.0	8.2	NS	NS
	1969-70	329.7	327.9	315.8	328.0	6.0	NS	NS
	1970-71	321.9	304.5	313.7	293.0	5.8	0.01	NS
	1971-72	289.0	298.1	265.3	267.2	6.1	0.01	NS
	1972-73	314.5	293.3	294.4	310.4	5.0	0.05	NS
	1973-74	276.5	292.5	282.2	278.3	4.7	0.05	NS
	Average	310.6	308.7	300.6	301.6	3.0	0.05	NS
Hass	1968-69	221.8	207.4	197.0	191.9	8.0	0.05	NS
	1969-70	207.8	205.3	210.6	206.5	5.2	NS	NS
	1970-71	234.1	200.2	204.9	195.3	4.9	0.01	NS
	1971-72	195.7	186.8	198.0	181.2	3.9	0.01	NS
	1972-73	238.1	236.5	267.1	248.1	4.3	0.01	NS
	1973-74	168.4	155.0	162.6	136.1	3.0	0.01	NS
	Average	210.9	198.5	206.7	193.2	2.5	0.01	NS





Fig. 3. Comparison of the effects of 7-day and 28-day irrigation intervals on the typical cumulative growth curve of the diameter of cv. Hass fruit (Aug., Sept. 1972). Arrows represent irrigations.

Fig. 4. Effect of irrigation interval on growth of Hass fruit (daily average, 1972). ○ 7 days; ▲ 28 days.

### Fruit growth

Fig. 3 illustrates fruit growth in August–September 1972. A considerable growth difference was registered between the two extremes of irrigation intervals (7 and 28 days). After 28 days, the difference in the fruit circumference was 1.3 mm. Fig. 4 represents fruit growth of cv. Hass throughout the summer of 1972. In June, growth was uniform, as a result of water being available in the soil following winter precipitation. The greatest growth differences were recorded in July–August, the period of most vigorous fruit growth, whereas in September growth was again uniform. In autumn a growth advantage was registered with fruit irrigated at the 28-day interval.

### Oil content

Oil content is largely affected by yields and by the size of fruit. It is difficult, therefore, to evaluate the effect of irrigation interval on the percentage of oil in the fruit. Oil content was determined in the years 1968–1970, when yields of cvv. Ettinger and Fuerte were relatively uniform. Results point to a tendency towards a decrease in oil percentage with an increase in irrigation interval (Table 4).

Table 4. Effect of irrigation interval on avocado fruit oil content

Each figure is the mean for five plots

Cultivar	Date of sampling	Average fruit weight (g)	Oil content (%)				SE of means	Signif. of linear component	Deviation from linearity
			Irrigation interval (days):						
			7	14	21	28			
Fuerte	8.xi.68	445	14.6	16.9	15.3	15.2	1.02	NS	NS
	25.xi.69	352	13.8	12.7	12.6	13.3	0.34	0.05	NS
	15.xi.70	261	14.8	15.2	15.5	13.5	0.43	0.05	NS
	15.xi.70	364	16.3	15.8	15.3	14.5	0.56	0.05	NS
Ettinger	10.x.70	298	10.7	10.3	9.4	9.9	0.23	0.01	NS
		277	10.4	10.0	10.1	9.3	0.17	0.01	NS
		234	9.0	9.5	8.9	9.2	0.22	NS	NS
	Average	270	10.0	9.9	9.5	9.5	0.11	0.01	NS

### Discussion

The relationship between growth of canopy and fruit-bearing capacity is known in a wide range of fruit trees (Zahner 1968), including avocado trees (Richards *et al.* 1962). Heavy crops suppress vegetative growth. The influence of alternate bearing of avocado trees on tree growth, especially of the Hass cultivar, has been well documented (Lahav and Kalmar 1972; Lahav *et al.* 1975).

The effect of irrigation regime on the rate of avocado tree growth has a cumulative character. Irrigation treatments influencing growth may, in the course of years, affect the productivity potential of the tree.

A close correlation was established between the tree volume and yield, especially with the Fuerte cultivar (Fig. 5), which was in fact less prone to alternate bearing than the Ettinger and Hass cultivars. With all three cultivars an average decrease in yields was registered (13%), with a lengthening of irrigation interval from 7 to 28 days, compared with a more substantial decrease (26%) in the tree volume

(Kalmar and Lahav 1977). Consequently a large tree, obtained with closely timed interval irrigations, bore correspondingly more fruits. However, as previously pointed out, the differences in yields were not significant. Possibly the influence of the irrigation regime was relatively small, because the weight of the crop depends on the number of fruits rather than on their size (Tables 1, 2). The differential irrigation treatments started with the termination of fruit set, and therefore they had only a slight effect on the number of fruits.

Because of the alternate bearing habit of the avocado tree, the effect of irrigation interval on yield could be determined only from data obtained by pairing of years. Although a constant drop in yields was registered for the Ettinger and Fuerte cultivars with an increase in irrigation interval, no significant difference concerning fruit number or weight was established. The average yield decrease for the 6-year period, for trees irrigated once a month as against those irrigated once a week, amounted to 4 kg/tree only (Table 1). This decrease is accounted for by the influence of irrigation regime on fruit size (Table 3).




Fig. 5. Relationship between tree canopy volume at end of experiment and the average yield (1969-1974).

Regression equations:

Ettinger:  $Y = -54.39 + 2.15x$ ;  $r = 0.674$

Fuerte:  $Y = -52.92 + 1.94x$ ;  $r = 0.889^*$

Hass:  $Y = 12.62 + 0.48x$ ;  $r = 0.754$ .

With the Hass cultivar an average decrease of 10 kg/tree was registered for trees irrigated at 28-day *v.* 7-day intervals. The decrease in yield of 17% only is surprising in light of the well-known sensitivity of the Hass cultivar to water supply (Honski 1969), and considering extremes of irrigation intervals of 21 and 28 days. The decrease in yields for cv. Hass was expressed in fruit number and size. An analysis of 5-year yields (without 1968-69) of this cultivar gives comparative figures of three 'on years' with the 7-day interval against only two 'on years' with the 21-day interval. Alternate bearing continued also in the year following the termination of the experiment (1974-75), although the entire plantation was irrigated that summer at a 21-day frequency. The yields for the 6-year period (1969-70 to 1974-75) showed no differences between 7-day and 21-day irrigation intervals (Table 5). Only at the 28-day interval was a significant decrease in productivity found.



A decrease in the seasonal water regime increased significantly the efficacy of its utilization for fruit production in all three cultivars. The highest capacity of water utilization for fruit production was found in the Hass cultivar. An addition of 1000 m<sup>3</sup> water/ha (above 5000 m<sup>3</sup>) resulted in yield increases of 380 kg/ha for Ettinger, 320 kg/ha for Fuerte, and 820 kg/ha for Hass. The relatively low utilization capacity of Fuerte—a very important cultivar in both Israel and California—should be noted, especially as regards the two basic parameters, canopy and fruit production. This may explain the tendency to plant Hass rather than Fuerte trees (Rock 1971–72).

A distinct influence of irrigation interval was expressed in the increase of individual fruit size with increased frequency of irrigation. The influence was greatest with the Ettinger cultivar: fruit picked from trees irrigated every 7 days was 35 g larger than that of trees irrigated every 28 days (6-year average). With Fuerte the difference was 9 g, and with Hass it was 18 g. The smaller effect on the fruit size of the Fuerte and Hass cultivars may be explained by the fact that they were picked well after the start of the rainy season, which was not the case with the Ettinger cultivar. The Fuerte and Hass cultivars demonstrated considerable fruit growth with 'dry' treatments in autumn after rainfall, especially with the high temperatures still prevailing then. The increased fruit growth in autumn of trees irrigated at 28-day intervals can be explained by retarded growth during the summer. The same could be observed with trunk growth (Kalmar and Lahav 1977). Autumn irrigation was often followed by an average daily growth of 0.4 mm or more in fruit diameter.

Table 5. Effect of irrigation interval on the yield of Hass avocado trees in the last 5 years of experimentation (1969–70 to 1973–74) and in 1974–75

Each figure is the mean for five plots

Period	Yield (kg/tree)				SE of means	Signif. of linear component	Deviation from linearity
	Irrigation interval (days):						
	7	14	21	28			
5 years, 1969-70 to 1973-74	67.0	60.2	53.2	51.6	2.2	0.05	NS
1974-75	11.8	41.8	95.8	62.4	4.3	0.01	NS
6 years, 1969-70 to 1974-75	57.8	57.1	60.3	53.4	1.8	0.05	NS

It can be concluded that, to achieve optimal fruit size with the maximum saving in water, irrigation should be frequent in the summer but withheld in the spring (after fruit set), when it would be wasted, as water reserves in the soil are still plentiful after the winter rains. Also in autumn, with the termination of the vigorous growing stages of the fruit, the effect of irrigation treatments is relatively small.

A decrease in irrigation intervals brought about a slight increase in the oil percentage (Table 4). Since the oil percentage in the fruit constitutes an important indicator for determination of harvest date, it should be possible to advance somewhat the harvesting date for the Ettinger and Fuerte cultivars by frequent irrigations to increase the oil percentage and fruit size.

In view of the similar yields obtained at all intervals, and the predominance of the Hass cultivar, it can be inferred that for avocado plantations, on heavy soil and under sprinkler irrigation, the preferable regime is approximately 6680 m<sup>3</sup>/ha (plus

6000 m<sup>3</sup> of winter rains), given at 21-day intervals. Under this treatment, water was more effectively utilized than with the other irrigation treatments tested. More frequent irrigation is preferable only when the objective is to advance the harvest date of Ettinger fruit, or to increase the size of Hass fruit in an 'on year.'

## References

- Gazit, S., and Spodheim, R. (1969). Fruit ripening. In 'The Division of Subtropical Horticulture, 1960-1969', pp. 83-5. (Volcani Institute of Agric. Res.: Bet Dagan, Israel.)
- Homski, S. (1969). Avocado growing. Div. Agric. Publs. Minist. Agric., Haquiria, Tel Aviv, Israel, Publ. No. 84 (in Hebrew).
- Kalmar, D., and Lahav, E. (1977). Water requirements of avocado in Israel. I. Tree and soil parameters. *Aust. J. Agric. Res.* **28**, 859-68.
- Lahav, E., and Kalmar, D. (1972). The value of physiological indicators for planning irrigation of the avocado tree. Prelim. Rep. Agric. Res. Organ., Volcani Center, No. 725 (Hebrew, with English summary).
- Lahav, E., Kalmar, D., and Zamet, D. (1975). Daily fluctuations in the growth of avocado trunks and fruits and the effects of irrigation and meteorological conditions on them. Pamphl. Agric. Res. Organ., Volcani Center, No. 149 (Hebrew, with English summary).
- Richards, S. J., Warneke, J. E., and Bingham, F. T. (1962). Avocado tree growth response to irrigation. *Yearb. Calif. Avocado Soc.* **46**, 83-7.
- Rock, R. C. (1971-72). A projection of California avocado acreage and production to 1977. *Yearb. Calif. Avocado Soc.* **55**, 37-40.
- Zahner, M. (1968). Water deficits and growth of trees. In 'Water Deficits and Plant Growth', ed. T. T. Kozlowski, Vol. 2, pp. 191-254. (Academic Press: New York.)