Taxonomy and Management of Andisols

Proceedings of the

SIXTH INTERNATIONAL SOIL CLASSIFICATION WORKSHOP

Chile and Ecuador 9 to 20 January 1964

Part 3: Tour-Guide for Ecuador

FIELD TRIF BACKGROUND SITE AND PEDON DESCRIPTIONS ANALYTICAL DATA

PREPARED BY:

SOIL MANAGEMENT SUPPORT SERV.CES, USDA SOIL CONSERVATION SERVICE WASHINGTON, DC SOCIEDAD ECUATORIANA DE LA CIENCIA DEL SUELO

Correct Citation:

F.H. Beinroth, W. Luzio L., F. Maldonado P., and H. Eswaran. (1985). Proceedings of the Sixth International Soil Classification Workshop, Chile and Ecuador. Part III: Tourguide for Ecuador. Sociedad Chile na de la Ciencia del Suelo, Santiago, Chile.

Copies of this publication may be obtained from:

Program Leader
Soil Management Support Services
USDA-Soil Conservation Service
P.O. Box 2890
Washington, D.C. 20013
U.S.A.

Sociedad Chilena de la Ciencia del Suelo Pontificia Universidad Católica de Chile Facultad de Agronomía Casilla 6177 - Santiago CHILE



SOCIEDAD CHILENA DE LA CIENCIA DEL SUELO



SOCIEDAD ECUATORIANA DE LA CIENCIA DEL SUELO



UNIVERSIDAD DE PUERTO RICO



SOIL MANAGEMENT SUPPORT SERVICES

CONTENTS		
Preface	1	
Workshop program		
Introduction		
General aspects		
Physical aspects		
Physical characteristics of the excursion route	39	
Analytical methods		
Literature cited		
E. Maldonado, R. Maldonado and C. Zebrowski: The volcanic ash soils of Ecuador	61	
Pedon descriptions and analytical data	77	
Ecuador N° 1	79	
Ecuador N° 9	85	
Ecuador N° 6	91	
Ecuador N° 4	97	
Ecuador N° 5	103	
Ecuador N°12	109	
Ecuador N°13	115	
Appendix I: Supplemental analytical data and descriptions	121	
Appendix II: Micromorphological descriptions, Dr. R. Honorato	153	
Appendix III: Soil moisture and temperatures regimes	157	
Appendix IV: Data sheet explanation	163	
Appendix V : Supplemental data sheets	175	

PREFACE

6TH INTERNATIONAL SOIL CLASSIFICATION WORKSHOP ECUADOR

Steering Committee:

F. H. Beinroth

R. W. Fenwick

National Coordinator, SMSS, SCS, Washington, P.C.

Dr. W. Luzio I.

University of Chile, Pres., Chilean Society of
Soil Science

Dr. F. Maldonado

USAID, Quito, Ecuador, Pres., Ecuadorian Society
of Soil Science

Dr. M. Leamy ICOMAND, Chairman, New Zealand

Host Organizing Committee:

Ecuador Steering Committee

Dr. F. Maldonado, Chairman USAID/Quito Ecuador
G. del Posso M. Soils Department of Ecuador
C. Yanchapoxi Soils Department of Ecuador

Technical Commission

C. Zebrowski

G. del Posso M., Chairman Soil Survey Department of Ecuador (PRONAREG)

C. Luzuriaga Soils Laboratory of Ecuador

Organizational Commission

G. Yanchapoxi, Chairman Soils Survey Department of Ecuador (PRONAREG)

I. Cordova Soil Conservation Department of Ecuador (INIAP)

ORSTOM Mission in Fcuador

W. Garces Soil Conservation Department of Ecuador (PRONACOS)

P. Toledo Remote Sensing Department of Ecuador

<u>lntroduction</u>

The Sixth (VI) International Soil Classification Workshop is one of a series of workshops and training activities of the Soil Management Support Services (SMSS). SMSS is a program of international technical assistance in soil survey, soil classification and correlation, and use and management of soils, primarily in tropical and subtropical areas. The program is funded by the U.S. Agency for International Development (USAID), and is carried out by the U.S. Dept. of Agriculture Soil Conservation Service (USDA/SCS). The previous workshops that have been held are:

	Country	Year	International Committee (ICCM)
I	Brazil	1977	1 COMLAC
ΙΊ	Malaysia/Thailand	1978	ICONTAC/ICOMO.
]]]	Syria/Lebanon	1980	1COMMGRT/1COMID
1 V	Rwanda	1981	ICONLAC/ICOMOX/ICOMMORT
V	Sudan	1982	1COMERT/ICOM1D/ICOMMORT
VI	Chile/Ecuador	1984	ICOMAND

These workshops are an integral part of the activities of SMSS and provide a medium for soil scientists from all over the world to discuss the mandates of the International Committees. These committees are working to refine Soil Taxonomy with respect to use and application in intertropical areas and as a means for agrotechnology transfer.

The VI International Soil Classification Workshop is organized as a joint endeavor of the Sociedad Chilena de la Ciencia del Suelo and the Sociedad Fcuatoriana de la Ciencia del Suelo, both acting in behalf of various universities and national institutions, the SMSS, the University of Puerto Rico (UPR), and USAID. This cooperation and collaborative effort has been the prime reason for the success of our previous workshops. The present workshop also promises to be most successful.

This tour guide is a compilation of some basic information on the history, climate, soils, geology, vegetation, and agriculture in Ecuador. It has been prepared and written by Mr. G. del Posso, Mr. C. Luzuriaga, Mr. C. Zebrowski, and translated from Spanish to English by Dr. F. Maldonado. This contribution is under the auspices of the Ecuadorian Soil Science Society. The soils were described by T. D. Cook of SMSS, SCS and G. del Posso and others of the Soil Survey Department. The analyses of the soils were performed by the University Laboratory in Ecuador and the National Soil Survey Laboratory (NSSL) of the SCS at Lincoln, Nebraska. This workshop and tour guide could not have been accomplished without the assistance, contributions and cooperation of many people and organizations. To them we are most grateful.

The assistance of the Director General and his staff of the National Agrarian Regionalization Program (PRONAREG), and the National Institute for Agricultural Research (INIAP), and National Soil Conservation Program (PRONACOS), and the National Council of Science and Technology (CONACYC) is greatly appreciated. G. del Posso and F. Maldonado have also given of their time and energy to make this workshop a success.

We would like to thank Dr. Steve Holzhey, Head, NSSL, and his staff for performing all the analyses in a timely manner. Special analyses were run by Dr. John Kimble, Soil Chemist, SMSS, that are required for the proposed orders of Andisols. He also efficiently coordinated and compiled the data and information for the tour guides. We are grateful for his cooperation. Thin sections and micromorphological descriptions were made by Prof. Ricardo Honorato. His work and contributions are greatly appreciated. We also express our gratitude to Dr. Koji Wada, Kyushu University, for X-ray and mineralogical analysis.

We also appreciate the advice and suggestions from Dr. F. Colmet-Daage, Dr. Mike Leamy, Dr. A. Van Wambeke and Dr. K. Flach.

To make an international meeting like this successful, many Ecuadorean colleagues and organizations have contributed untold hours of extra work beyond their normal duties. We acknowledge their assistance and support. This workshop, like the previous ones, is a model of international cooperation and achievement. In this same spirit, we anticipate beneficial and productive discussion at this workshop to advance and improve Soil Taxonomy and to exchange knowledge on soil classification and agrotechnology transfer between scientists.

RICHARD W. FENWICK
National Coordinator
SMSS

SIXTH INTERNATIONAL SOIL CLASSIFICATION WORKSHOP

- Taxonomy and Management of Andisols

CHILE AND ECUADOR

9 to 20 January 1984

PROGRAM FOR ECUADOR

Saturday, 14 January 1984

- Participants arrive at Quito airport; transfer to and check-in at Hotel Inter-Continental Quito
- 1500 Registration Free afternoon and evening in Quito; night in Quito

Sunday, 15 January 1984

Free day or optional sightseeing in Quito; night in Quito

Monday, 16 January 1984

0800 Registration of Ecuadorian participants, Hotel Inter-Continental Quito

OPENING CEREMONY

Chairman: J. Delgado

Venue: Pichincha Room, Hotel Inter-Continental Quito

0900 National anthem of Ecuador

Welcome -- F. Maldonado

Official inauguration -- H. Ortiz, Undersecretary of Agriculture

Introductory remarks by: J. Goodwin, USAID/Quito R.W. Arnold, SCS/SMSS M.L. Leamv. ICOMAND

Monday, 16 January 1984 (cont'd)

1000 Refreshments

Technical Session VII: RATIONALES FOR TAXONOMIC CRITERIA

FOR ANDISOLS: SUBORDERS

Chairman: R. Tavernier
Rapporteur: R. W. Fenwick

1030 S. W. Buol: Use of soil temperature regimes in Soil Taxonomy

- 1100 A. Van Wambeke: Soil climatic regimes as criteria for establishing sub-order of Andisols.
- 1120 S. Shoji: The case for recognizing a sub-order of non-allophanic Andisols
- 1140 Discussion
- 1200 Lunch

Technical Session VIII: RATIONALES FOR TAXONOMIC CRITERIA
FOR ANDISOLS: GREAT GROUPS AND
SUBGROUPS

Chairman: S. W. Buol Rapporteur: A. Alvarado

- 1330 M. Otawa: Criteria for great groups of Andisols
- 1400 F. Colmet-Daage: Criteria for subgroups of Andisols
- 1430 Discussion
- 1500 Refreshments

Technical Session IX: PROBLEMS OF ANDISOLS Chairman: R. L. Parfitt
Rapporteur: V Neall

- 1530 J. M. Kimble and W. D. Nettleton: Analytical characterization of Andisols
- 1550 T. D. Coca: Field of identification and mapping of Andisols
- 1610 R. W. Arnold: Criteria for intergrades between Andisols and other orders
- 1630 Discussion

Monday, 16 January 1984 (cont'd)

Technical Session X: FIELD TRIP BACKGROUND Chairman: C. O. Scoppa

- 1700 E. Maldonado and F. Maldonado: Distribution of volcanic ash soils in Ecuador
- 1730 G. Yanchapaxi and H. Serrano: The Machachi soil sequence
- 1800 G. del Posso and C. Luzuriaga: The Santo Domingo soil sequence
- 1830 Announcements
- 1845 Adjourn

Night in Quito

Tuesday, 17 January 1984

Field Trip I: QUITO-MACHACHI-SANTO DOMINGO-QUITO

0730 Dep. Hotel Inter-Continental Quito

0900 Arr. Pedon ECU 01

Discussion Leader: F. Maldonado Rapporteur: A. Cortes Refreshments

1030 Dep. Pedon ECU 01 1100 Arr. Pedon ECU 09

> Discussion Leader: G. del Posso Rapporteur: F. Maldonado

1215 Dep. Pedon ECU 09

1300 Arr. INIAP Experiment Station Santa Catalina for lunch

1345 Dep. Cafeteria

1400 Arr. Pedon ECU 06

Discussion Leader: W. Luzio Rapporteur: M. Mendoza

1530 Dep. Pedon ECU 06

1545 Arr. auditorium of Santa Catalina station, refreshments

Tuesday, 17 January 1984 (cont'd)

Technical Session XI: EROSION, CONSERVATION AND MANAGEMENT OF ANDISOLS

Chairman: F. Maldonado Rapporteur: W. Luzio

1600 G. de Noni and J. F. Nouvelot: Erosion and conservation of volcanic ash soils in the highlands of Ecuador: A. case study

Discussion with INIAP staff about soil management
Review of pedons ECU 01, 09 and 06

1730 Dep. Santa Catalina station

1900 Arr. Quito

Night in Quito

Wednesday, 18 January 1984

Field trip V: QUITO-SANTO DOMINGO

0730 Dep. Hotel Inter-Continental Quito

0930 Arr. Pedon ECU 04

Discussion Leader: A. Alvarado Rapporteur: C. O. Scoppa Refreshments

1045 Dep. Pedon ECU 04

1330 Arr. INIAP Experiment Station Santo Domingo for lunch

1430 Dep. Cafeteria

1445 Arr. Pedon ECU 05

Disucssion Leader: F. Colmet-Daage Rapporteur: G. del Posso

1615 Dep. PedonECU 05

1630 Arr. meeting room of Santo Domingo station
Refreshments
Discussion with INIAP staff about soil management
Review of pedons ECU 04 and 05

1730 Dep. Santo Domingo station

1900 Arr. Santo Domingo

Night in Santo Domingo, Hotel Zaracay

Thursday, 19 January 1984

Field Trip VI: SANTO DOMINGO-QUEVEDO-SANTO DOMINGO-QUITO

0730 Dep. Hotel Zaracay

0900 Arr. Pedon ECJ 12

Discussion Leader: A. Cortes Rapporteur: C. Zebrowski

Refreshments

1015 Dep. Pedon ECU 12

1115 Arr. Pedon ECU 13

Discussion Leader: A. Van Wambeke Rapporteur: A. Alvarado

1230 Dep. Pedon ECU 13

1300 Arr. INIAP Experiment Station Pichilingue for lunch

1400 Discussion with INIAP staff about soil management

Review of pedons ECU 12 and 13 and field trips in Ecuador

1500 Dep. Pichilingue station

1630 Arr. Santo Domingo, refreshments

1700 Dep. Santo Domingo

2000 Arr. Quito

Night in Quito, Hotel Inter-Continental Quito

Friday, 20 January 1984

Technical Session XII: EVALUATION OF THE ANDISOL PROPOSAL IN DIFFERENT ENVIRONMENTS

Chairman: W. G. Sombroek Rapporteur: J. M. Kimble

- OBOO H. Ikawa: A critical evaluation of the placements of the Andepts of Hawaii according to the proposed key for Andisols
- 0830 F. N. Muchena: A critical evaluation of the placements of the Andepts of Kenya according to the proposed key for Andisols
- 0900 B. Clayden: A critical evaluation of the placements of the Adnepts of New Zealand according to the proposed key for Andisols.
- 0930 Discussion

Friday, 20 January 1984 (cont'd)

1000 Refreshments

Technical Session XIII: | COMAND SUMMARY DISCUSSION

Chairman: M. L. Leamy
Rapporteur: B. Clayden

1030 Discussion

1230 Lunch

Technical Session XIV: RECOMMENDATIONS OF THE WORKSHOP

Chairman: F. H. Beinroth Rapporteur: T. D. Cook

1400 Discussion and adoption of commendations and recommendations

CLOSING SESSION
Chairman: R. Tavernier

1530 The workshop in retrospect -- W. G. Sombroek

1545 Workshop impact in Chile -- W. Luzio

1600 Workshop impact in Ecuador -- F. Maldonado

1615 Workshop impact on 1COMAND -- M. L. Leamy

1630 Vote of thanks -- A. Van Wambeke

1645 Adjourn

Refreshments


2000 Closing Dinner, Hotel Inter-Continental Quito

Saturday, 21 January 1984

Participants depart Quito

Optional pedologic or touristic programs

Guide Book for Field Trip



Organized by:

SOIL MANAGEMENT SUPPORT SERVICES
Soil Conservation Services U.S. Department of Agriculture
co-sponsired by University of Puerto Rico

TOUR GUIDE

SIXTH INTERNATIONAL SOIL CLASSIFICATION WORKSHOP TAXONOMY AND MANAGEMENT OF ANDISOLS CHILE AND ECUADOR

January 9-20, 1984

International Organization: USDA, Soil Conservation Service

National Coordination: Sociedad Ecuatoriana de la Ciencia del Suelo

Sponsors:

Soil Conservation Service (SCS)

Agency for International Development (AID)

National Agrarian Regionalization Program (PRONAREG) National Institute of Agricultural Research (INIAP) National Council of Science and Technology (CONACYT)

Prepare by:

Guillermo del Posso, Ing. Agr. M.S.*

Carlos Luzuriaga, Ing. Agr. M.S.*

Translated by:

Fausto Maldonado, Ing. Agr. Ph. D.**

^{*} Members of the Soils Department, PRONAREG
** Member of the Ecuadorian Soil Science Society

INTRODUCTION

This tour guide has been prepared for the field tour in Ecuador for the Sixth International Soil Classification Workshop which will be held in Chile and Ecuador from January 9 to 20, 1984.

The tour will show the distribution and types of volcanic ash soils, starting in the highland "Sierra" and going down to the tropical lowlands. These Andisols are the result of the interaction of different tropical temperature and moisture regime and vegetation on the volcanic ash of different ages.

This guide includes general information on the country, descriptions of relevant physical aspects of the profiles and their surroundings, and the morphological descriptions and analytical data for each pedon.

The guide was prepared by the Technical Committee, Mr. Guillermo del Posso, Mr. Carlos Luzuriaga, and Mr. Claude Zebrowski and translated by Fausto Maldonado, of the Ecuadorian Soil Science Society.

The authors acknowledge the support and assistance received from the Director and from Departments of the Ministry of Agriculture, National Agrarian Regionalization Program (PRONAREG).

Finally, the Technical Commission received several suggestions and advice from Dr. Hari Eswaran and Terry Cook of the Soil Management Support Services, Soil Conservation Service, United States Department of Agriculture and Dr. Fred Beinroth, University of Puerto Rico.

1. GENERAL ASPECTS

1.1 History

A "Homo sapiens" of Asiatic origin lived in the country about 20,000 years ago; several stone tools found close to Quito prove his existence.

Around 1470 A.D. the Incas conquered the area; both the oral tradition and written documents testify the important role they played.

Tupac Yupanqui, his son Hupanqui, his son Huayna Capac and this latter's children, Atahualpa, and Huascar, who kept a permanent antagonism had a decisive role in the Inca empire history.

The Spanish conquest started in 1526 when, on September 21 in the second of the three expeditions that left Panama for the southern countries, Bartolome Ruiz discovered what is now Ecuadorian territory.

Sebastian de Benalcazar founded Quito, the present capital city of Ecuador, on December 6, 1534. Other cities were founded later: Portoviejo and Guayaquil, Loja and Cuenca.

On August 10, 1809, Quito declared independence from Spain although the independence process was only completed in 1822. After a short period as part of Gran Colombia, Ecuador was established as an independent republic in 1830.

1.2 Population

Ecuador has currently more than 8 million inhabitants according to the last estimate of 1983; population density is about 32 persons/km 2 ; 64.7 percent of them live in the coastal region, 56.6 percent in the highlands (Sierra) and only 2.1 percent in the Amazoniann region.

The Interandean Valley, a series of basins support a rural population mostly in scattered houses. Population density frequently exceeds 100 inhabitants/km². This high population density is the consequence of historical and geographic phenomena. On the slopes of the Andean Cordilleras, inhabited sites go up to 3,500 m above sea level, the very top level of crops, although some local variations exist due to microclimates.

In the Coastal area, the population distribution is related to the regional agricultural development while in the Amazoriann region, population is localized close to the rivers. Colonization in recent years has developed at the Cordilleras' foothills and along the roads leading to oil exploiting areas.

The Galapagos Islands have about 6,000 inhabitants or less than $1/km^2$ only four of the islands are inhabited. The remaining make up part of the world known natural reserve.

Like other andean countries, Ecuador has a high percentage of native Indian's mainly in the highlands; their best known language is quichua. Native tribes also live in the Amazonian region, although since oil exploitation, a strong colonization current has developed with people from the highlands moving into this area.

The Costal Plain shows a great population diversity including indigenous, black, asiatic, and white people of Spanish origin.

1.3 Political Division and Government (Figure 1)

Ecuador is administratively divided into 20 Provinces. Five in the Coast, ten in the Sierra, four in the Amazonian Plain, and the Galapagos Islands. The Provinces are divided in "cantones" (126), these into "rural" (720) or "urban" (227) "parroquias." This division is the result of a centenary evolution in function of the different stages of space occupation. The Provinces cover areas of different dimension, depending on populated or partly populated areas, or if they have a recent or older development.

Government

According to the Constitution of January 15, 1979, Ecuador is a democratic republic with a President elected by the people on an absolute majority basis. The Vice President, elected with the President, is the Head of the National Development Council, institution in charge of preparing the development plans for the country. The National Chamber of Representatives is the legislative power. The Judicial Power is formed by the High Provincial Courts and the Supreme Court as the top judicial authority based in Quito.

Figure 1. ECUADORIAN LAZGGY **PROVINCES** 2Bolivar 7 3 Cañar 10 4 Chimborazo 5 Cotopaxi 17 6El Oro 15 7 Esmeraldas 13 8 Galápagos 9 Guayas 18 12 10 imbabura 16 II Loja 9 12 Los Rios 13 Manabi 14 14 Morona Santiago 15 Napo 16 Pastaza н 19 17 Pichincha 18 Tungurahua 19 Zamora Chinchipe 20 Carchi

President

The President of Ecuador is Dr. Osvaldo Hurtado, who assumed the power on May 24, 1981. Ecuador will have new elections in January 1984.

1.4 Economic Overview

Ecuador is a small developing country whose economy has been traditionally based on agriculture. About half of its population obtain their main income from agriculture, livestock rearing, or by exploiting other renewable resources--forests and fisheries.

The Andes range, here divides into two main branches, with a high valley in the middle. This produces a wide variety of climates and the geologic configuration adds to facilitate the production of tropical crops (banana, cacao, coffee, cotton, sugar cane, rice, etc.) as well as temperate climate crops and tropical crops of the highlands like potatoes, maize, etc.

Although improved technology has reached several production areas, traditional agricultural techniques are still being used in many areas. In these areas, poor or at least lower than average yields are obtained. The introduction of improved technology has resulted in labor underemployment and, as a consequence rural to urban migration, with all the associated problems.

Economic development has been based, during the last decade, mainly on oil production from the Amazonian region. This led to continuing development when many other countries were suffering economic problems. The GNP constantly increased, there was a solid international monetary reserve, a reduced inflation rate (10 to 15 percent), and a stable money value. This led to large investments in the infrastructure, to the introduction of heavy subsidies to some imported products like wheat, and to the establishment of political prices. Agricultural production has not received the necessary support from the government and, at the beginning of the eighties, the reduction in oil production and its prices have led to the same economic situation found in most of the developing countries.

Nevertheless, Ecuador has a great potential in its natural renewable resources which are a source of new income, like the shrimp production industry which now represents the second largest export product.


1.5 COMMUNICATIONS (Figure 2)

Roads: A total of 50,000 km of roads make up the road infrastructure of which 8,000 km are paved (asphalt), 14,000 km are other permanent roads, and 18,000 are only for seasonal use.

Railways: Two railways exist--one going from Guayaquil to Riobamba, Quito, Ibarra, and San Lorenzo; and the other joining Riobamba to Cuenca.

Maritime Transportation: This is very important because about 95 percent of the export and import products are carried by ship. Ecuador has a national company for common freight (TRANSNAVE) and another for oil derivatives transportation (FLOPEC).

Air Transportation: There are two international airports and 42 small airports or airstrips. Ecuador has an international airline, Ecuatoriana de Aviacion, serving the Western South American counties, Panama, and the United States of America. Other small airlines serve the internal needs of the country.



Telecommunications: Both phone and telex services permit good communication within the country and with the rest of the world.

2. PHYSICAL ASPECTS

2.1 Physiography: Geographical Setting and Relief Geographical Setting.

Ecuador covers an area of $275,032 \text{ km}^2$ and is located on the Western coast of South America (Figure 3). It is bounded on the north and northeast by Columbia, on the east and south by Peru, and on the west by the Pacific Ocean. Its population, geography, history, and development make up part of that great geographic Region of the Andes Range, which goes along the western side of the continent, here between the Pacific Ocean and the Amazonian Plain. Also belonging to Ecuador, the Galapagos Islands located about 1,000 km from the contine cal seashore. The scientific value of these islands has been known for a longtime and has led to the recognition of the islands as a Humankind Heritage.

PANAMA

PODUTOS

Galapagos

Islands

OCEAN

PERU

Andes mountain range


The name of Ecuador has its origin in the equator which crosses the country about 30 $\,\mathrm{km}$ north of Quito.

Relief

The major relief is the Andes Range which divides the country into three natural regions: The Coastal Plain (the Costa), the Andean mountain ranges, the Interandean Valley (the Sierra), and the Eastern plains which belong to the Amazon Basin (the Orients of Amazonian in Region). (Figures 4 and 5)

Figure 4.


NATURAL REGIONS



The Sierra

The Andes Range form a mountainous barrier 100 to 120 km wide with external slopes of more than 4,000 m in two dominant directions: NE-SW on the north from Quito and south of Alausi and N-S in the middle. From the Colombian border in the north to the town of Alsusi in the south, there are two parallel ranges (West and East Cordillers) with altitudes between 4,000-4,500 m and cool or cold climates; several snow peaks stand out of these cordilleras, the highest being Chimborazo Mt. at about 6,300 m.

The central part of the Sierra is about 50 km wide and has a series of intermountainous valleys separated by minor transverse chains called "nudos," which with the main chains form an association similar to a ladder. The valleys are flat or rolling, and they range in altitude from 1,600 to 3,000 m. The whole area is covered by volcanic ash.



South from Alausi to the town of Zaruma, there are no high volcanoes and the mountains' altitude range from 3,600 to 4,700 m. Further south to the Peruvian border, the mountains are not higher than 3,500 m and no volcanic ash covers them.

The Costa

The Costa is the area west of the Western Andes Range from 600~m to sea level. It is widest in the area of Guayaquil (180 km) and is about 100~km in the south.

To the west and close to the ocean, there is a small range which goes from Esmeraldas southward to Portoviejo and Jipijapa and to the east of Guayaquil.

Between the Andes and the Coastal Range, there is a large plain about 80 km wide, slightly dissected to the north, close to Santo Domingo (600 m) and gently sloping to the south These alluvial plains and valleys are drained by large rivers. The seashore shows different landscapes; straight line beaches are found as well as cliffs and small bays and mangrove stands.

Amazonian Region or Oriente

Occupying almost 50 percent of the country, the Amazonian region is located to the east of the Cordillera Oriental and belongs to the Amazon Basin. Average altitudes are 500 m or less and the topography is formed by level plateaus which are usually dissected by rivers. The region is still mostly under forest and has a humid tropical climate.

The Galapagos Islands

This insular area of volcanic origin is formed by 13 islands at about 1,000 km west of continental Ecuador. They have different sizes and are formed by volcanoes emerging from the sea and reaching altitudes to 1,600 m.


2.2 Geology and Recent Volcanism Geology (Figure 6)

The present geologic structure of Ecuador reflects the interaction of various orogenic and paleogeographic events among which the most important is the uprise of the Andes Cordillera. While this occurred, the Andes were under an intense erosion process, which led to the formation of continental sediments (of detrital origin; i.e., the mechanical alteration and destruction of rocks). Through the different geologic eras, these sediments were deposited either in the external slopes or in the graven between the cordilleras. Conglomerate, breccia, sandstones, clays, and shales are among these rocks.

On the other side, several oceanic transgressions and regressions, have produced the deposition of marine sediments. Among these limestones, clays, and sands are found.

Volcanism produced both acid and basic lava (rhyolitic, andesitic, and basaltic) and pyroclastic, both coarse and fine.

There are also sediments of mixed origin: volcano-sedimentary formations resulting from the volcanic activity in marine environments and sediments of the fly ash type.



Some groups of crystalline and sedimentary terrains, under strong pressures resulted in the formation of metamorphic rocks (gneiss), slates, marble, quartizes, etc.

Finally, as a consequence of breaking produced by tectonic movements, certain formations were crosscut by liquid crystalline rocks which led to the formation of intrusive massifs of granit, diorite and granodiorite.

Volcanism, tectonic movements and sedimentary deposits are the foundations of Ecuador geology, but the physiography was remodeled during the Quaternary period. In fact, late volcanic activity has covered a large area of the Andes with a thick pyroclastic layer, especially a mixture of dust and volcanic ask, frequently hardened and known as "cangahua." On the other side glaciations, mass movements, and erosion by rivers have carried a high amount of detrital materials, contributing to the formation of moraines, alluvial fans, and spreading materials to form the eastern plateaus and river valleys.

Recent Volcanism

Volcanism is important in Ecuador due to the relief formed and the thickness of and area covered by the products emitted. In fact, volcanism has met here the appropriate conditions, probably due to plate tectonics, where plate expansion has produced abundant faults and fractures through which volcanic products were ejected to the surface. Recent volcanism plioquaternary, is formed in the following regions:

In the Andean area where two volcanic regions can be identified, the first one from the Chiles to the Sangay Volcanoes is characterized by stratovolcanoes. On the western cordillera--necks and calderas predominate. The altitude of the whole area is less than 5,000 m. Only Iliniza, Carihuairazo, and Chimborazo Mountains are higher and are permanently snowcapped. On the eastern cordillera, snowcapped mountains are more frequent. There are large volcanic cones (Cotopaxi, Tungurahua and Sangay) with slopes of about 70 percent and with hundred-meter-wide craters. These volcanoes have erupted andesitic lava of the aa type and mainly pyroclastics. An unresolved problem in the presence of the vast volcanic formation known as "cangahua" which might be an eolian deposit.

Volcanic activity has moved from the western to the eastern cordilleras and may currently be seen in the Cotopaxi, Tungurahua, and the active Reventador and Sangay volcanoes.

Going south, the second region from Sangay volcano to Zaruma, the large stratovolcanoes disappear and a large plateau is found with cliffs, locally resembling organ pipes. This large plateau is made of a mass of rhyolitic lava flows covered by a volcanic ash layer of above 50 cm.

In the subandean zone, (east from the main cordilleras) volcanism is reduced to three volcanoes, located on the Napo anticline. These are similar to the volcanoes in the eastern cordillera, but are more Stromboli type and their lava is basaltic rather than andesitic.

In the Galapagos Islands, the 14 main islands are formed from volcanoes which have emerged from a submarine mountain range. Isabela Island for instance, the largest one in the archipelagos, is formed by six volcanoes. Volcanism here is closer to the Hawaiian-type characterized by shield-like

cones with very wide calderas. Most common are the basaltic lava, very fluid of the pahoehoe type. The highest cones $(1,000\ \text{to}\ 1,700\ \text{m})$ and the widest calderas $(5\ \text{to}\ 9\ \text{km})$ are in the Fernandian and Isabela Islands.

2.3 Climate (Figure 7)

Being on the equator and with the high relief of the Andes, Ecuador has a great range of climates even over very short distances.

Main Factors Influencing Climate

Climate is influenced by two main factors: the general atmospheric circulation and the local air masses resulting from the relief. There are few other minor factors like the ocean currents which play an important role in the Coastal zone climate.

The General Atmospheric Circulation

Ecuador, as a whole, is located within an equatorial belt characterized by low atmospheric pressures, high moisture, and high temperatures. As a consequence of the tilting of the earth's rotation axis in relation with its orbit around the sun, this belt periodically moves towards the Southern Hemisphere. On the other side close to the 30° north and south, there are two subtropical anticyclones with a less moist, less warm air, and with high atmospheric pressures; this leads to a displacement to the lower pressure area of the equatorial belt, the Intertropical Convergence Zone. Air masses coming from the two hemispheres come in contact to generate an intertropical front, unstable zone with atmospheric disturbance, as the air masses try to overcome each other.

During the year, the intertropical front will follow the same movements as the equatorial belt:

--From February to July the front will move to the north, influencing the western area of the American continent. Ecuador is covered by temperate and rather dry air masses coming from the south and producing a stable and rather agreeable weather.

--From September to December the front enters the Southern Hemisphere influencing the eastern zone of the continent. Ecuador is under warm and moist air masses coming from the north resulting in a temperature increase, atmospheric instability, and stormy weather.


Local Air Masses

Topographic contrasts produce four types of air masses each with different properties.

--Warm air masses of oceanic origin are located over the Pacific Ocean. They are very humid and move towards the continent. Close to the Cordillera (West) the warm air rises uphill due to convention and, as it cools, it forms clouds and rain. Most of this thick cloudy layer stays onsite, but a part enters the interandean valley, although diminished of part of its moisture.

--Warm air masses of continental origin are located over the Amazonian region. Also rising by convection, they cool on the eastern cordillera and produce heavy rains on that side, sometimes entering the Interandean Valley. This type of continental air masses also exist over the coast but are not as important as over the Amazonian in area.

CLIMATE



- --Temperate air masses are located over most of the Andes between 2,000 and 3,000 m above sea level.
- --Cold air masses, rather small, are ${f located}$ around the tops of the highest mountains.

Oceanic Currents

The Ecuadorian Coast and particularly the Santa Elena Peninsula, are under the influence of the cold Humboldt current most of the time, whose origin is the Antartic Ocean. When it is close to the coast, usually between May and October, the cool air masses cause a lowering of the temperature. Although these masses are moisture-saturated, no heavy rains occur, only a light mist and drizzle, locally called "garua."

A counter current is formed, coming from the west, opposed to the Humboldt current. It reaches the Ecuadorian coast from December to April. This results in oceanic warm winds and storm formation with temperature increases over the coast.

A still unexplained phenomenon is the southward movement of the current, reaching even to the north of Chile, which facilitates the movement of warm weather (25-29°C) coming from Panama. This phenomenon is known as "El Nino" (Child Jesus) because it starts commonly in December. On the average, it is produced every 7 years. Its consequences are a mixed blessing: on one side comes the abundant rain to a dry area; on the other fish and aquatic birds are killed, or at least pushed to colder areas.

All the above influences are compounded to give the great variety of Ecuadorian climates.


Main Climate Classes (Figure 8)

The factors mentioned above combine to produce different kinds of climate in Ecuador. Eight classes have been identified and their names show both the amount and distribution of rainfall during the year (tropical or equatorial, having one or two rainfall peaks, respectively) and the average temperature.

- --Megathermal rainy. With a mean annual temperature around 25° C., total rainfall exceeds 3,000 mm and may go up to 6,000 mm per year with a very regular distribution, slightly decreasing from December to February. Relative humidity is 90 percent or higher. This class is found in the extreme northern area (Esmeraldas) and in the Amazonian region.
- --Tropical megathermal moist. This covers the outer hill-slopes of the two cordilleras. Annual rainfall exceeds 2,000 mm and may reach 5,000 mm, mostly in one rainy season. Temperature ranges from 15 to 24° C according to changes in altitude. Relative humidity is about 90 percent.
- --Subhumid tropical megathermal. Rainfall ranges between 1,000 and 2,000 mm, and it is concentrated in one period (December to April), with the remaining months dry. Average temperature is 25°C and relative humidity ranges between 70 and 90 percent. This class covers an 80 km-wide belt in the coast from the northern border to Guayaquil.

DISTRIBUTION OF CLIMATES

(Tipos de climas)



- --Dry tropical megathermal. Rainfall averages between 500 to 1,000 mm concentrated from January to April. The remaining months are very dry and average temperature is very high. The area influenced by this class is about 60 km wide west of the previous class and from the equator to about 4°30' S.
- --Subarid tropical megathermal. Mean annual rainfall in less than 500 mm and mean temperatures range between 20 and 26° C and relative humidity is about 80 percent. The rainy season is characterized by a cloudy sky, mists and very low intensity rain. Low mountains more than 300 m high are a little moister. This class covers small areas at the coast in Manta, Santa Elena, and the coastal belt of the Galapagos Islands.
- --Subhumid equatorial mesothermal. It is the most common over the Interandean area, except for areas higher than 3,000 to 3,200 m and some valleys. Total annual rainfall ranges from 500 to 2,000 mm. Mean temperatures from 10 to 20° C, following the altitude and relative humidity goes from 65 to 85 percent.
- --Dry equatorial mesothermal. Temperatures range between 18 and 22° C with almost no change from one season to another. Two dry seasons separate two rainy seasons which total less than 500 mm rainfall. Relative humidity ranges from 50 to 80 percent and clear skies are common.
- --High altitude equatorial climate always affects areas over 3,000 m. Mean annual temperature is about 8° although variations go from 0° to 20° in some cases. Rainfall ranges between 1,000 and 2,000 mm according to altitude and aspect of hillsides. Relative humidity is almost always higher than 80 percent.


Fauna

Ecuador is an area of exceptional interest concerning its flora and fauna. In spite of its relatively small area comparatively in South America, a high percentage of species are found in the country. For instance, 1,500 out of the 2,600 bird species in this part of the world are found in Ecuador. This is of course due to the ecological diversity going from the coastal desertic areas to the Amazonian jungle. The Andes Cordillera has contributed to create differences in flora and fauna too.

2.4 Land Use (Figure 9)

Agricultural production diversity in Ecuador is exceptional, including tropical as well as temperate climate products. In the area close to the equator, the highlands of the A.ides Cordillera have crops typical of four season areas (except that day length does not change during the year). Changes in temperature are due to altitude combined with differences in moisture too.

Land use changes also according to plots size. Land distribution shows the known pattern of large areas belonging to few owners: 67 percent of the landholdings are less than five hectare, and occupies 7 percent of the agricultural land area, while 2 percent of the landholdings are of more than 100 hectares, and occupying 48 percent of the area. Crops occupy most of the minifundia (78 percent of land) while pastures (47 percent) are most common in the large haciendas (37 percent of land). Large haciendas benefit from irrigation water, more than small landholdings.



The introduction of mechanization has resulted in a decreased use of agricultural labor, which causes unemployed. This has caused many people to migrate to the cities.

All the above compounds to produce less than the country needs. Wheat, fats (oils and animal fats), and milk represent 32.5 percent of all agricultural imports; paper and wood products amount to 22 percent and the line of agricultural inputs; and machinery (33 percent) show how dependent the national agriculture is on outside inputs. Nonetheless, the balance of the sector within the external trade is rather positive. By 1980 there were \$392 million imports versus \$701 exports. Four tropical products make 91 percent of all agricultural exports: banana (34 percent), cacao (30 percent), coffee (20 percent), and sugar (7 percent).

The Sierra

The upper limit for agricultural use ranges from 3,200 to 3,800 m, depending on local drought, wind, or mist exposure. Few potato plots can reach the 4,000 m contour line. Above it only a natural grass (paramo) vegetation is found whose only use is for sheep (rarely cattle) or llama grazing.

Each crop is localized at a specific altitude level following the most suitable combination of temperature and moisture conditions. This results in the formation of well characterized belts of the various crops. The top level includes barley, broad beans, andean tubers like oca, (Tropaeolum sp.) and potato (Solanum tuberosum).

Above 3,800 m a very harsh climate and the frost danger prevent the growth of crops.

The middle level from 2,400 to 3,000 m is the maize area. Maize is mostly associated with beans. When irrigation water is available horticulture becomes important, as around Ambato. This level also slows pasture development, either natural in few areas or cultivated. It is at this level that wheat is being produced, although its production has decreased due to high production costs in comparison with imported wheat.

The lower level is the one of sugar cane production (below 2,200 to $2,400\,$ m). This crop is still found in the large haciendas, while the small farmer prefers to have tomato, beans, and fruit crops like avocado. These crops grow only with irrigation.

In the southern province of Loja, the cordilleras are lower than in the north, and the area has a seasonal agriculture based on the rather irregular rainy season. Maize (hard, high protein) is there associated with another species of bean, the sarandaja (Dolichos sp.). Some tropical crops such as cassava and peanuts are also grown.

On the outer hillslopes of the cordilleras, land has been cleared by a slash and burn system of cultivation. The older the occupation of land, the more common is the presence of pastures and less of other crops.

The Sierra region shows the highest population pressure. Minifundia of less than 5 hectares make up 78 percent of all landholdings, but cover only 11 percent of the area. Only 52 percent of a small farmer income originates

in agriculture. The remaining 48 percent is obtained from other sources, especially from various jobs performed in the cities, as construction workers, etc.

The Coast

Moisture changes from southwest to northeast influence the tropical agriculture of the coast.

Desertic Southwest. The Santa Elena Peninsula and the area close to the Peru border are occupied by natural vegetation of dry areas like cacti, and show very unfavorable conditions for agricultural use.

Moist Northwest. Both accessibility, topography, and the high rainfall which have produced a tropical jungle and have prevented an early development or any kind of exploitation. Current use is reduced to pastures and few perennial or semipermanent crops (Cassava, plantain, coffee, and to a lesser degree, cacao) along the main roads and rivers. To the south of this area, colonization projects are being started.

The Center and Southeast. The tropical moist climate, with a dry season, provides favorable conditions for agricultural use. Two areas can be identified: the coastal hills and the eastern plains.

The coastal hills. The presence of clouds above 300 m has encouraged the use with tropical tree crops: coffee, cacao, and citrus are often mixed with subsistence crops and pastures in the small landholdings. In the zone of low hills and moderate relief, most of the area is under pasture. Cattle rearing is mainly for meat production.

The Plains. With the exception of the west central part whose unfavorable soil conditions allow only the development of natural pastures for extensive grazing, the plains area is formed on sedimentary materials of high fertility, where the main industrial and export crops are established.

--Cacao is more sensitive to moisture than coffee and grows in the southern part. There are about 1 million hectars of single crop and 150,000 mixed with coffee. Cacao yields are usually low due to the lack of a good plantation management and because most plantations are old. Annual production is about 90,000 tons.

--Coffee is grown all over the plains (C. robusta) where it has been cultivated more recently than in Manabi and the south of the Andes (C. arabica). It often grows mixed with cacao, bananas, citrus, and even with mango or other fruit crops in the small farms. As a single crop, it occupies about 120,000 hectares. Annual production ranges between 75,000 and 80,000 tons.

--Bananas are cultivated under two systems (1) small (family) farms where they grow associated to other fruit crops, this fills most of the country needs; (2) large farms produce export bananas. Their production started back in the fifties in Los Rios and Guayas Provinces and more recently in El Oro Province. Over production and international commercialization problems have brought about a decrease in the plantation area (about 200,000 hectares in 1972, 110,000 hectares in 1980).

--Sugar Cane. In small plantations is mostly for alcohol and brown sugar production while large enterprises work with associated sugar mills. The latter are to the east of Guayaquil. Sugar production of about 3,800,000 tons is slightly over the country's needs.

--Rice is mostly cultivated in the flood plain with a total area of 130,000 hectares. Present yields could be improved with a better water use and with improved techniques. Annual production of 145,000 tons does not meet the country's needs.

--Oil palm, of comparatively recent introduction, is cultivated in the more moist humid northern area along the main roads. Oil palm production increased from 21,000 tons in 1970 to 208,000 tons in 1980. It is one of the main raw materials to make edible oil, but its production has not met the country's needs yet. Soybeans are produced mainly in Los Rios Province and complement oil palm in the oil industry.

In spite of the importance that all these products have, the west lowland plains are under utilized as compared to their potential capability. Topography and soil conditions permit an intensification of all the crops and justify high investments to improve irrigation and drainage.

The Amazonian Region

Long before the Spanish conquest, aborigine people in this area were using the land under a shifting cultivation system based on forest clearing of the jungle. This, along with hunting and fishing, provided a balanced diet. Shifting cultivation has been, with slight variations from site-to-site, the method used all over the Amazonian region. Based on the same starting need: that being low natural fertility of the equatorial forests' soils, their rational use implies a rotational cultivation. Indigenous people to the Ecuadorian Amazonian cultivate up to 40 different species in one vegetable garden or orchard; the most common are cassava, plantain, Colocassia sp., and other tubers. The production period of a plot varies according to soil fertility from 3 to 10 years. A plot abandoned for 25 years can not be distinguished from the surrounding forest.

In the foothill areas of the cordillera where colenization fronts are moving forward, shifting cultivation is being changed to extensive cattle raising introduced by the colonizers. Instead of having small isolated plots, there is a whole area of forest clear and converted to pastureland. This results in a further impoverishment of already low fertility soils, with no compensation for the lost organic matter.


Also present in the area are a few industrial enterprises which started with tea production and more recently the introduced oil palm.

A balanced development of the Amazonian region requires a difficult compromise between the aspirations of those who chose to settle in these areas and the native people presently there who want their ways respected.

2.5 Soils* (Figure 10)

The interaction of climate and parent materials results in the formation of several soils which can be included in four groups:

^{*}Ecuador has adopted as official soil classification system the USDA, SCS "Soil Taxonomy."



12

2.5a Alluvial Soils

- l. Little developed or nonhydromorphic soils which are found in the western valleys of the coastal region, mainly between Babahoyo and Machala, although they are also present in Manabi (north and northeast of Portoviejo) and northeast of Guayaquil. They are mostly Fluvents and they are either water saturated a short time in the year or not at all. Soil textures are coarse for those soils closer to the cordillera and finer in those away from the mountains. These are young soils with no diagnostic horizons, except for an Ochric epipedon. They are usually fertile soils and well suited to a wide range of crops.
- 2. Andic soils. These have developed on volcanic ash deposits; they are well drained and they are mostly found in the Northern Amazonian region. They belong to the Andic Dystropepts or Typic Dystropepts subgroups in "Soil Taxonomy." (Soil Survey Staff, 1975) They have volcanic or allophanic minerals; are moderately deep and clayey; they have a rather low fertility, but with a good management and fertilization could produce all kinds of crops suited to the area.
- 3. Hydromorphic soils. They are poorly drained soils where biologic activity is limited. Artificially drained, these soils can be used and have good fertility. They are Aquepts or Fibrists.
- 4. Saline water saturated soils. They are found in the mangrove areas and are mostly Sulfaquepts.

2.5b Soils on Recent Volcanic Ejects

They are formed on deposits of lapilli or ashes which have been transformed by climatic influence to crystalline or amorphous clays (allophane). They can be distinguished of each other by their age or the climate under which they have developed.

- 1. Sandy soils. They are found around recent volcanoes or in dry volcanic areas. They have a coarse (sand) texture, low organic matter content, and low water retention capacity. They have been classified as Psamments (?) or Vitrandepts when they have a high content of volcanic glass or pumice.
- 2. Loamy soils. They are the result of a more advanced weathering of the sandy soils, due to moister climate. They are rich in organic matter and nutrients, the highest fertility being shown by soils in the Interandean Valley from Tulcan to Azogues. They are mainly Ustolls or Udolls, with some Eutrandepts.
- 3. Allophanic soils. They are found at higher altitudes in the volcanic Sierra, moister areas where allophane is the secondary material instead of crystalline clays. Water retention capacity goes from 50 to 100 percent. They are richer in organic matter than the previous ones, but with lower fertility. Poor climatic conditions (cold) and steep slopes are the main constraints to their use. They are Dystrandepts and have the known problem of high phosphorus fixation.
- 4. Perhumid allophanic soils. These soils are found in areas of high rainfall along the outside slopes of the cordilleras, or in areas with older ash deposits. With the high allophane content, there is a

higher water retention capacity (more than 100 percent) in these soils and they are almost completely nutrient-depleted; their fertility is very low and their agricultural use and management represent a difficult problem. In few areas, these soils are covered by less humid allophanic layers developed from recent volcanic deposits. They are Hydrandepts, characterized by an irreversible dehydration resulting in hard coarse aggregates when the soil profile is exposed to air drying.

2.5c Soils on Ancient Materials

Whatever they have originated, the most important factor of these soils' formation is climate, which results in crystalline clay formation, sometimes rich in silica.

Montmorillonite soils. They are found where climate action is reduced, so rock weathering is not too advanced; silica and bases are little leached. These are montmorillonite-rich soils with high nutrient content. They are in general good soils for agriculture, nevertheless, there are certain differences between them:

- 1. Heavy clay soils, extremely plastic when wet but very hard and with deep cracks when dry. These soils have developed on clayey sedimentary rocks; carbonate deposits decrease as rainfall increases. This relationship corresponds to soil classification as follows:

 - a. Vertisols with high CaCO₃: Torrerts and Usterts
 b. Vertisols with low CaCO₃: Usterts
 c. Vertisols without CaCO₃: Uderts
- 2. Soils on flat areas: They are Planosols, characterized by a sandy or silty top soil overlying a more clayey layer. For the "Soil Taxonomy" (Soil Survey Staff, 1975) they are Paleustalfs or Paleargids. The former are Alfisols with an argillic horizon, with more than 35 percent base saturation; they have developed under an ustic regime. The latter are Aridisols with an argillic horizon.
- 3. Soils without vertic characteristics. They do not show cracks, even during the dry season there is no contraction or expansion. In the "Soil Taxonomy" (Soil Survey Staff, 1975) they are:
 - a. Clay soils of the dry area: Ustropepts
 - b. Clay soils of the moist area: Eutropepts
- 4. Well structured black soils. They have a Mollic epipedon, easily identified in the field. They have high content of organic matter and are found in the uphills of the Coastal Mountain range where high moisture allows organic matter accumulation. They are Mollisols in the "Soil Taxonomy," (Soil Survey Staff, 1975) either Ustolls or Udolls.

2.5d Kaolinite Soils

These soils are found in areas with very high rainfall and temperature factors which have led to kaolinite formation after bases and silica have been leached. These soils are grouped as follows:

1. Saturated soils: They are found in areas of high rainfall but have more than 50 percent base saturation and are nutrient-rich soils. They have been classified as Oxic Eutropepts or Oxic Tropudalfs, the former being more frequent.

- 2. Unsaturated soils. They are poor soils of limited agricultural potential. They have low base saturation and have been grouped as unsaturated compact soils, unsaturated eroded soils, and unsaturated clayey soils. These have been classified according to the "Soil Taxonomy" (Soil Survey Staff, 1975) as:
- a. Unsaturated compact soils: Oxic Dystropepts,
 b. unsaturated eroded soils: Oxic Dystropepts and
 Entic subgroups soils.
 - c. unsaturated clay soils: Oxic Dystropepts and Haplortoxs

2.5e Very Shallow Soils

They are grading to no soil (rock outcrops) and are located either over recent lava flows, on the mountain crests or in highly eroded dry areas. They are mostly Orthents.

- 3.2 Physical Characteristics of Excursion Route* (Figures 11 and 12)
 In order to make a brief description of the different zones where
 the soil profiles selected for the field trip are located, the excursion route
 was subdivided into three parts. Each well defined according to its major
 differences: physiography, geology, climate, land use and soils.
 - 3.2.1 Interandean Valley (from Quito to Machachi) Physiography and Geology

This part starts in Quito City $(2,850\ m)$ and follows for Los Chillos Highway in direction to the Interandean Valley $(2,450\ m)$, where it is possible to experiment some decrease in altitude and after an increase in the Machachi Valley $(3,000\ m)$, with this difference we concluded that the north part is lower than the south part.

The Interandean Valley is dominated by Andean Chain which is integrated by two branches. East and west to each side of the valley. The altitude of the branches ranges between 3,200 m at the level of hillock (collado: lowest part of the chain) and 5,000 m where there are some volcanoes as Pichincha (4,749 m), El Corazon (4,788 m) and Cotopaxi (5,897 m). The typical characteristics of this zone are the presence of pyroclastic materials in all the area.

This material is from the last eruptions of the Pichincha and Cotopaxi volcanoes, which have formed some layers of great thickness (from 1 m to 10 m or more), characterized by alternative layers of Lapilli (2 to 40 mm of diameter) and alternated layers from successive emissions from volcanoes.

Climate

In this part if is possible to observe a climatic sequence in two senses:

- 1. North-South.--In this sense exists basic differences in humidity. In the south, it is higher than in the north, this means that Machachi valley has more precipitation than Calderon (north part of Quito).
- 2. East-West.--In this case the differences are higher than the last one because with the increase in the altitude from valley to the higher part of the Andean Chain in both sides of the valley, the quantity of precipitation increases significantly (Figure 13).

^{*}This section was prepared by G. de Passo, Chief of Soils, Department of PRONAREG.

ROUTE EXCURSION





Figure 12

Precipitation Data

Sense	Place	Mean Annual Precipitation (mm)
North	Calderon	600.0
	Quito	1,236.4
	St. Catalina	1,400.0
South	Machachi	1,500.0

Source: PRONAREG, 1972-1982; prepared by G. de Posso.

Figure 13.



Prepared by G. del Posso.


Land Use

According to the climate sequence and in relation to the altitude, the land use of this zone is the following:

The upper limit of the land use for agriculture is between 3,200 m to 3,800 m depending on local conditions. Over this limit is located the coldest natural grassland called "Paramo."

Each crop is located in a defined level of altitude according to its requirements (basically temperature and humidity). The following scheme permits to observe this distribution: (Figure 14)





Prepared by G. del Posso

There is a change in land use in this zone according to the size of the farms. In general, in the valley area exists large farms (more than 100 hectare called Latifundios) and a few small farms (less than 5 hectares called Minifundios). In a few cases some large farms start in the valley and go to the boundary of the Paramo.

The majority of the small farms (Minifundios) are located generallly on high lands characterized by slopes of more than 40 percent.

The dominated land use in the big farms (Latifundios) is pasture, but in the Minifundios, the normal land use is crops (specially high andean crops) which is the typical food for the Indians:

```
--Melloco (Ullucus tuberosus)
--Oca (Oxalis tuberosa)
--Chocho (Lupinus mutabilis)
--Mashua (Tropaeolum tuberosum)
--Haba (Vicia Faba)
--Potatoes (Solanum tuberosum)
--Maiz (Zea maya)
```

The pastures in the big farms are:

```
--Rye grass and leguminous
```

--Kikuyo

--Oat and leguminous

The cattle dominant are:

```
--Holstein F. (imported from U.S.A.)
```

Source: INIAP.

⁻⁻Holstein F. plus criollo cattle (cattle from Ecuador)

Soils

The distribution of the soils in this part of the route is related with climatic sequence according to the altitude. The following scheme has summarized the principal factors in this sense:

First part of excursion route from Quito to Machachi

ALTITUDE	2,400	3,200	3,600
(m)	-,	3,200	3,000

Precipitation (mm)

Dry season (mm)

Temperature (°C)

Temperature Regime (°C)	Isothermic	Isomesic	Isofrigid
Principal crops	maize, pasture, wheat, vegetables	barley, potatoes, and high andean crops	natural grassland of Paramo
Soils	Mollisols Eutrandepts	Eutrandepts Dystrandepts	Cryandepts

Source: Soil Department--PRONAREG: prepared by G. del Posso.

3.2.2 Counterfort of the West Branche of Andean Chain in direction to the Coastal area (from hillock to Santo Domingo).

For this second part of the excursion route, the principal physical characteristics are summarized in the following scheme between hillock (collado) to Tandapi and Tandapi--Santo Domingo:

Second part of excursion route

Physical Hillock (collado) Tandapi to characteristics to Tandapi Sto. Domingo Physiography -Strong slopes (50%) -Very strong slopes and -Area covered by pro-(70%), but near to Geology clastic materials Sto. Domingo City, exempt the eroded the relief becomes places, where there more soft and in are outcrops of old Sto Domingo is volcanic rocks. practically flat. -Area also covered by volcanic ash, where the aerial projection are finer and have less thickness then the last one. Climate -Very humid zone -Very humid zone (Udic M.R.) (Perudic M.R.) -The temperature -The temperature in increase according to this zone is practithe altitude (decrease cally uniform and more or less 2° C by there is small each 200 meters). increase until Sto. -From isofrigid to Domingo. isohyperthermic limit. -Isohyperthermic (more 22° C. Altitude 3,700 to 1,600 m 1,600 m to 500 m Land Use Below grassland of -Sugar cane, banana, Paramo: barley, pasture, oil palm, potatoes, habas, maiz, coffee, rubber, pinecold humid forest, subapple, manila, cocoa, tropical forest, subetc. vegetation. Soils Dystrandepts Dystrandepts Hydranpets

Source: Soil Depto. PRONAREG: prepared by G. del Posso.

3.2.3 Guayas River Basin (Sto. Domingo--El Empalme)

This third part is included in the Guayas River Basin and presents the following physical characteristic:

Third part of excursion

route SANTO DOMINGO EL EMPALME

Physical Characteristics

Altitude (m) 550 80

Physiography
and
Geology

-In general the relief of this area is flat.
-There is a gradual decrease in altitude from Sto.
Domingo to El Empalme.
There are two volcanic layors in all the area.

There are two volcanic layers in all the area which belong to two emissions separated by a

genetic period.

Climate Sto. Domingo Quevedo El Empalme

Temperature uniform: more than 22° C

Soil temperature: isohyperthermic

Precipitation

Dry season

Land Use -Oil palm -Short cycle crops

-Rubber -Maize

-Coffee and Cocoa -Coffee and Cocoa

-Banana -Banana

-Tropical pasture -Tropical pasture

Soils Dystrandepts Eutrandepts Hydrandepts Tropudalfs

(Sto. Domingo) (El Empalme)

Source: Soil Department--PRONAREG: prepared by G del Posso.

ANALYTICAL METHODS

The following is a brief description of the procedures and methodology used by the National Soil Survey Laboratory. If a more detailed description of any procedure is needed, the following report should be consulted. "Soil Survey Investigations Report No. 1, Soil Survey Laboratory Methods and Procedures for Collecting Soil Samples," USDA, SCS, Revision, August 1982. For equivalent procedures but ones of a less automated and instrumented nature, the 1972 revision should be consulted.

Method codes are included in the column headings on the data printouts and can be used as a key to the methods and procedures.

Some general comments on how the samples are handled:

All samples are air-dried and crushed using a wooden roller and sieved to remove fragments greater than 2 mm in diameter. All determinations are made on sieve samples and reported on an ovendry weight basis unless otherwise stated. The ovendry weight basis is determined by ovendrying a sample at 105°C.

CONSIDERATIONS IN THE USE OF LABORATORY DATA

In consideration of the soil characterization data, be aware of the ways your concepts of specific properties compare with the same properties defined operationally. Each of us has mental pictures and conceptual definitions to in visualizing properties and processes. These frequently go far beyond the information supplied by any particular analysis, and we are tempted to give numbers on paper credit for connotations that are strictly our own. this light, there are two pitfalls. One is the temptation to mix observations with concepts. Rather than first laying observations and then building inferences, it tempting to describe and interpret in terms of predetermined inferences. If, at some later date concepts change, there may be no way to untangle the observations from the outdated concepts to piece together new inferences.

The second pitfall is the temptation to remain unaware the operations (analyses) involved in an observation or measurement. Results differ with different methods, but the nature of any particular operation remains the same so long as the same method is used. To compare one bit of data another, it is important to know how both bits gathered. This leads to operational definitions (methods). The operational definition is a device to describe a soil property in terms of the operations used to measure it. operational and conceptual definitions as we for discuss the following laboratory analyses. Then, as we use field and laboratory data to classify soils, we can attempt the difficult task of separating observations from concepts.

SYNOPSIS OF ANALYTICAL METHODS

PARTICLE SIZE DISTRIBUTION ANALYSIS (PSDA) AND COARSE FRAGMENTS--RECORDED AS WEIGHT PERCENT.

The fractions coarser than 20 derived from mm are screening and weighing in the field or from field volume estimates. Be sure to include volume estimates of the 20-75 mm and 75-250 mm fractions when sending samples to the laboratory. The 2-20 mm fraction is obtained by sieving in the laboratory. During sample preparation, the soil is crushed so all but the coarse fragments pass through a 2-mm sieve. In soil survey, coarse fragments are split the 2-5 mm, 5-20 mm, and 20-75 fractions and reported as percent of the less than 75 mm, material. Fragments larger 75 mm but smaller than 250 mm are included in the column labeled > 2 mm, percent of the whole soil (really percent of the < 250 mm material).

For PSDA, a portion of the fraction passing through the sieves treated with H₂O₂ to remove organic matter and is candle filtered to remove soluble salts. It is dispersed in water with sodium hexametaphosphate, a dispersing agent, and with a specified physical agitation. The dispersed soil is poured through a sieve which catches the sands and allows the silt and clay to pass through. various sand fractions are further separated by dry shaking in a nest of sieves. The silt fractions and clay are determined by pipetting known volumes from suspensions remaining after known settling times. Fine (< 0.0002 m) is determined by centrifugation and pipetting. The necessary settling and centrifugation times are based calculations which treat the particles as perfect spheres with consistent, designated specific gravity (Stokes These calculations are used to determine the precise depths and times at which a pipetting will capture exactly the size fraction desired. Operationally then, the clay fraction is not the array of particles smaller array of particles that behaves in 0.002 mm. It is the suspension in water as though it consists of perfect spheres of a designated specific gravity and a diameter of 0.002 mm. The actual dimensions may be uncertain and quite different from the conceptual perfect spheres. It is possible to calculate sedimentation rates for other shapes and specific gravities, but this is only done in special cases. standard operation usually separates the more active fraction (clay), which is dominated by clay minerals, from the more inert fraction (silt) which is dominated by nonclay minerals. Thus, the operation would serve its purpose even if we had no concept of particle shape and specific gravity. fact, the concept of clay as a distinct kind of soil material predates the development of modern concepts about clay-size particles and clay minerals.

Clay-sized carbonates are determined by treating the pipetted sample of clay with HCl and measuring the pressure of CO.

The specific chemical and physical treatments and calculations are indicated by codes in the headings. Detailed procedures are listed by these codes in Soil Survey Investigations Report No. 1.

ORGANIC CARBON--REPORTED AS WEIGHT PERCENT

Organic carbon is determined by the Walkley-Black method which is a wet combustion procedure. The sample is treated with a strong oxidizing agent (potassium dichromate) and digested in sulfuric acid. An empirical recovery factor is used in calculating percent organic carbon. The NSSL uses a recovery factor of 0.77 as proposed by Walkley. This organic carbon represents decomposed soil organic matter. It normally excludes relatively fresh plant residues, roots, charcoal, and carbon of carbonates. It can be multiplied by 1.72 for an approximation of organic matter content.

The standard procedure is not used when organic carbon content exceeds roughly 8 percent. Loss on ignition at 400°C is an acceptable substitute in such cases.

LOSS ON IGNITION--REPORTED AS WEIGHT PERCENT

This is a measure of organic matter content and generally used for samples high in organic carbon (> 8%). Ovendried samples are held overnight at 400°C, and the weight loss is measured. Some of the weight loss is sometimes caused by dehydration of clay minerals, but normally the weight loss is mostly from burning of organic matter. When there are large amounts of amorphous material or gypsum present, this weight loss by dehydration is more of a problem. Remember, this approximates organic matter not organic carbon.

NITROGEN

Nitrogen is determined by Kjeldahl digestion with ammonium being steam distilled into boric acid and titrated with HCl using an automatic titrator.

CITRATE--DITHIONITE EXTRACTABLE IRON, ALUMINUM, AND MAN-GANESE--REPORTED AS WEIGHT PERCENT

This procedure was originally developed for extraction of iron from soils. Some authors have designated the extracted iron as "free iron" reflecting beliefs about the nature of this component. This method extracts and measures amorphous forms and some or the crystalline forms but not total soil iron. Aluminum extracted during the rather severe chemical treatment is also recorded, and the two values together are used in the spodic horizon definition. Manganese extracted by this method is usually less than

l percent. Some researchers use this Mn and relate it to internal soil drainage properties. Iron extracted by this procedure is often closely related to clay distribution with depth. Dithionite-citrate-extractable iron, aluminum, and manganese are determined by extracting a soil sample with an excess of sodium dithionite and sodium citrate in an 8-ounce bottle which is shaken overnight. Superfloc is added and an aliquot of the supernate is removed for the determination of iron, aluminum, and manganese by atomic absorption.

BULK DENSITY--REPORTED IN GRAMS PER CENTIMETER CUBED

Bulk densities are determined at two or more moisture contents. In coarse and moderately coarse-textured soils, they are determined at 1/10-bar moisture (moisture held against a 1/10 atmosphere pressure difference across a semipermeable membrane), and ovendry (moisture after drying at 110°C). In soils of medium and finer texture, they are determined at 1/3 bar and ovendry.

Most of the bulk densities are obtained equilibration of Saran-coated clods at these suctions. Clods for this purpose are selected in the field and are dipped in Saran dissolved in methyl ethyl ketone. dipped clods are dried by hanging them in air until the methyl ethyl ketone evaporates. The dry Saran then remains as a thin, tough film surrounding and supporting the clods for transport to the laboratory. This film is reinforced by added dipping in the laboratory, and supports the clods through wetting, equilibration at the desired suctions, and ovendrying. It can stretch and contract, thus allowing for changes in bulk density as moisture contents change. Bulk density is determined by weighing the clods in air, then weighing them in water. The first measurement is weight, and the difference between weight in air and weight in water is volume.

The bulk density as reported on the data sheet is calculated to represent the < 2 mm material.

COEFFICIENT--OF LINEAR EXTENSIBILITY (COLE)

COLE is a <u>calculated</u> value. It is based on the shrinkage of the natural soil clod between 1/3-bar moisture content (1/10 for sandier soils) and ovendryness. Therefore, bulk densities at these moisture contents are used to calculate COLE as follows:

COLE =
$$l(\frac{dry}{moist}, \frac{bulk}{bulk}, \frac{density}{density})^{1/3} - 1$$

The cubed root of the ratio of the two bulk densities is used to correct the clod shrinking in three directions. The value obtained from the equation is corrected for coarse fragments, hence, COLE is on a whole soil base. COLE

multiplied by 100 is called Linear Extensibility (LE) and is simply COLE in percent.

1/3-BAR WATER

The 1/3-bar water is determined by equilibrating natural fabric clods at 1/3 bar and determining the weight percent moisture as a percent of the ovendry weight of the soil.

15-BAR WATER

The 15-bar water is determined by pressure membrane extraction; < 2 mm soil material is used. The water retained is expressed as a percent of the ovendry weight of the soil.

WATER RETENTION DIFFERENCE--WRD

WRD, like COLE, is also a derived value based on moisture contents at 1/3 bar (1/10 bar for sandier soils) and 15 bar. The 1/3 bar or 1/10 bar moisture contents are often referred to as field capacity while the 15-bar content is often called the wilting point. WRD is the difference between 1/3 bar (1/10 bar) and 15 bar divided by 100 then multiplied by the moist bulk density to put it on a volume base. If the soil contains coarse fragments, the aforementioned value is corrected to represent the coarse fragment content, hence, WRD is on a whole soil base.

EXTRACTABLE BASES--REPORTED AS MEQs/100 g SOIL

These are the major exchangeable bases, magnesium, sodium, and potassium. They are extracted by displacement from the cation exchange complex by another cation, ammonium (ammonium acetate buffered at pH 7 is the The term extractable rather than extraction solution). exchangeable bases is used because any additional source of soluble bases will influence the results. In soils soluble salts or carbonates, the soluble cations must be measured separately and the results subtracted from the determination of exchangeable bases extractable bases for (Exchangeable = extractable - soluble). Exchangeable Na can be computed with acceptable accuracy unless salt contents exceed approximately 20 mmhos/cm at 25°C. Exchangeable Na (extractable Na) - (Na in saturation extract) (saturation percentage, percent water in the saturated paste) divided by (1,000). Exchangeable Na can be derived with greater accuracy than the other cations in the presence of gypsum or carbonates. Assuming exchangeable K negligible compared to exchangeable Ca and Mg, then exchangeable = CEC (NH,OAc) exchangeable Na. suitably reproducible for comparison approximation is between soils and for classification. Exchangeable Mg can be computed in the same fashion as exchangeable Na. Results

are not as satisfactory for exchangeable Ca when computed in the presence of carbonates or large amounts of gypsum. To prevent misuse of the Ca values, NH₄OAc extractable Ca is omitted from the data sheet when the carbonates are thought to significantly influence the results.

EXTRACTABLE ACIDITY--REPORTED REQS/100 g SOIL

This is the acidity released from the soil by a barium chloride-triethanolamine solution buffered at pH 8.2. It includes all the acidity generated by replacement of the hydrogen and aluminum from permanent and pH dependent exchange sites.

EXTRACTABLE ALUMINUM--REPORTED MEQS/100 q SOIL

This measures exchangeable aluminum, which is constituent only in acid soils (pH < 4.5). Aluminum will if the pH rises precipitate above 4.5 to 5.0 analysis. The Αl extracted by 1N KCl approximates exchangeable Al. Because KCl is an unbuffered salt, usually effects the soil pH only one unit or less, so the extraction is made near the pH of the soil. In acid soils, the sum of bases plus Al extracted by 1N KCl is close to the effective CEC of soil in the field.

CATION EXCHANGE CAPACITY -- REPORTED AS MEQs/100 q SOIL

Several different reagents and pH levels are used to measure the CEC. It may be determined by summing all the exchangeable cations, by summing bases and extractable aluminum, or by saturating the exchange complex with one kind of cation, replacing it with another cation, and measuring the replaced cation. Ammonium in neutral ammonium acetate is the ion used for the later method.

The CEC-sum of cations is calculated by summing the extractable bases and extractable acidity. It represents the pH dependent charge. CEC-sum of cations is not calculated if soils contain soluble salts. The bases + aluminum CEC value is calculated by summing the extractable bases and KCl extractable aluminum. It is commonly called the effective CEC for acid soils since it more closely represents CEC at the soil pH. It also is not calculated for soils that have soluble salts.

The CEC-NH $_4$ OAc is an analytically determined value. It is the value used in calculating CEC/clay ratios. The CEC-sum of cations - CEC NH $_4$ OAc is the pH dependent charge from pH 7 to pH 8.2.

CEC measurements at other pH or by other cations may each yield somewhat different answers. It is important to know the operational definition (procedure, pH, and cation used) before evaluating the meaning of the measurement.

BASE SATURATION -- REPORTED AS PERCENT OF CEC

Two values are frequently reported. One is derived from the equation:

Base saturation percentage by NH₄OAc, pH 7 is equal sum of NH₄OAc extractable bases divided by NH₄OAc-CEC times 100.

The other is:

Base saturation percentage by sum of cations pH 8.2 is equal to the sum of NH,OAc extractable bases divided by CEC sum of cations pH 8.2 times 100. This value is not reported if either extractable Ca or extractable acidity is

Differences between these two base saturations depend upon the size of the pH dependent CEC. Class definitions our soil classification specify which is to be used.

The sum of bases is considered equal to exchangeable cations unless gypsum, salts, or carbonates are present. Normally, the bases extracted by this method exceed 100 percent when the more soluble constituents are present. We then assume 100 percent base saturation. The Ca from carbonates is usually much larger than Mg. Extractable Ca is omitted if more than a trace (> 0.4%) of carbonate (as CaCO3) is present or if calculated base saturation exceeds 110 percent.

Other values can be obtained by using CEC derived from operationally differing methods. There are numerous differing methods utilized for numerous specific purposes. Always be sure which method you are dealing with.

EXCHANGEABLE SODIUM

Exchangeable sodium percentage (ESP). Calculated by dividing NH,OAc extractable sodium minus water soluble sodium by NH, ÖAc-CEC and multiplying by 100.

<u>CALCIUM CARBONATE EQUIVALENT--REPORTED AS WEIGHT PERCENT</u>

The amount of carbonate (CO₃) components in the soil are measured by treating the sample with HCl. The evolved CO₂ is measured manometerically. The amount of carbonate is then calculated on a CaCO₃ equivalent no matter what form (Na₂CO₃, MgCO₃, etc.). Calcium carbonate is most commonly reported on the < 2 mm base but, equivalent" in some soils with hard carbonate concretions, carbonates are determined on both the < 2 and 2-20 mm material; and reported on a < 20 mm base.

CALCIUM SULFATE AS GYPSUM--REPORTED AS WEIGHT PERCENT

Calcium sulfate (gypsum) is determined by extracting gypsum in water and precipitation in acetone. Gypsum is reported on both the < 2 mm and < 20 mm base.

pH--WATER, SALT, NaF

The pH is measured by a glass electrode in a soil-water or soil-dilute salt solution. The extent of the dilution is shown in the heading. l:l means one part by weight of water was added for each part by weight of dry soil. This is important in spodic soils because, within the limits of dilutions used, the pH increases markedly with increasing dilution. Measurements in soil-water are usually 1:l and are 2:l for the dilute salt solution.

The use of dilute salt solutions is a popular method for masking seasonal variations in pH. Readings in the dilute salt solutions are usually lower than with distilled The values may be equal or even higher in highly weathered soils of the tropics. These soils would have a anion exchange capacity. Readings in 0.01M CaCl tend to be uniform regardless of time of year. Readings in 1N KCl tend to also be uniform. The former are more popular regions lacking extremely acid soils. The latter are more popular in regions where many soils are quite acid KCl is used to extract exchangeable aluminum. The pH reading then shows the pH at which the aluminum was extracted.

The pH may also be measured in lN NaF. This measurement is done in soils where there is a large amount of amorphous material present. If there is a large amorphous component, this pH will be greater than 9.5. High values for the NaF pH will also be found if there are free carbonates. Therefore, care must be taken in interpreting this data. One gram of soil is placed in 50 ml of lN NaF stirred and the pH read after 2 minutes.

SODIUM PYROPHOSPHATE (pH 10) EXTRACTABLE CARBON, IRON, AND ALUMINUM--REPORTED AS WEIGHT PERCENT C, WEIGHT PERCENT FE AND WEIGHT PERCENT AL

This reagent has been used widely as an extractant of organic matter. It successfully removes much of the organo-metal accumulations in spodic horizons, but is relatively ineffective in extracting many forms of inorganically bound iron and aluminum. For this reason, it is a key to the chemical identification of spodic horizons.

MINERALOGY

The clay, silt, and sand fractions of selected horizons from each profile were analyzed to determine the mineralogical composition. The different techniques employed are briefly as follows:

X-ray diffraction analysis (XRD).

A Phillips diffractometer with a copper tube is used. XRD is run on the untreated clays and also

on K and Mg saturated clays with associated heating or glycerol solvation.

2. Differential thermal analysis (DTA) was done only on the clay fractions. The instrument used was the Columbia Scientific Thermal Analyzer. Semiquantitative estimates are made using the data.

 $\frac{\texttt{PETROGRAPHIC}}{\texttt{This}} \, \frac{\texttt{ANALYSIS--RECORDED}}{\texttt{analysis}} \, \frac{\texttt{AS}}{\texttt{is}} \, \frac{\texttt{NUMBER}}{\texttt{analed}} \, \frac{\texttt{PERCENT}}{\texttt{after}} \, \frac{\texttt{The}}{\texttt{the}}$ the petrographic microscope which is the chief tool for identification and quantitative analyses in the sand and coarser silt fractions. The data are listed supplemental sheet or at the bottom of the newer data sheets. Minerals are grouped under two major headings: resistant and weatherable.

SOLUBLE CATIONS AND ANIONS IN WATER EXTRACTED FROM SATURATED PASTE--REPORTED AS MEQS/LITER OF SOLUTION

Soluble cations and anions are determined in the water extracted from a saturated paste made by adding distilled water to soil until the saturation point is reached. saturation, the soil paste glistens as it reflects light, flows slightly when the container is tipped, and slides freely and cleanly off a spatula except for soils containing large amounts of clay. In calculations of exchangeable sodium percentage, soluble sodium is subtracted from extractable sodium to obtain an estimate of exchangeable sodium.

TOTAL SALTS

Measure of percent total estimated soluble salt in the soil. Calculated from conductivity of the saturated extract.

ELECTRICAL CONDUCTIVITY

Electrical conductivity (EC) of the saturate extracted obtained the saturated paste used in the determination of water soluble anions and cations.

SAR

Sodium adsorption ratio determined by the following equation:

$$SAR = \sqrt{\frac{c_a^{++} + Mg^{++}}{2}}$$

The Ca, Mg, and Na are in meg/l and determined from the saturated extract of the water-soil saturated paste.

PHOSPHATE RETENTION CAPACITY

Determined by shaking 5 g of air-dry soil (< 2 mm) with 25 ml of a phosphate solution (1 ml = 1 mg F) for 24 hours in a 50 ml polypropylene centrifuge tube (Blakemore et al., 1981). Tubes are centrifuged and the P remaining in solution measured using the molybdo-vanadate method. Results are reported as percent, P-retained.

ACID OXALATE-EXTRACTABLE ALUMINUM AND SILICON

Determined by shaking 0.5 g air-dry soil (< 2 mm) with 50 ml acid oxalate reagent for 4 hours in a polyropylene centrifuge tube in the dark (Blakemore et al., 1981). Tubes are centrifuged and Al and Si determined by atomic absorption spectrophotometry.

KOH-Al

Determined by reacting 0.2 g soil with 2 ml 4N KOH for 10 minutes, filtering and determining the Al in the solution.

AVAILABLE PHOSPHOROUS (Bray-1)

Determined by shaking 1 g of air-dry soil (< 2 mm) with 10 ml of extracting solution (0.025N HCl + 0.03N NH₄F) for 5 minutes in a 50 ml Erlenmeyer flask. Extracts are filtered through Whatman No. 2 filter paper. A 2 ml aliquot is added to 8 ml solution of 50 ppm P prepared from reagent grade potassium hydrogen phosphate (KH₄PO₄). Phosphorcus is measured using a colorimeter and reported in parts per million.

OTHER USEFUL INFORMATION

There are a couple ratios placed on the data sheet. These are useful in making internal checks of the data and making other useful determinations.

RELATIONSHIP OF MINERALOGY AND CATION EXCHANGE

Ratio of CEC (NH,OAc, pH 7) to clay can be used to make an estimation, of the clay mineralogy. For family placement, some CEC/clay relationships are as follows: Kaolinitic < 0.2, kaolinitic or mixed 0.2 - 0.3, 0.3-0.5, mixed or montmorillonitic 0.5 - 0.7montmorillonitic > 0.7. When amorphous material is present, these relationships do not hold.

Ratio of CEC (at pH 8.2) to 15-bar water of greater than 1.5 and more exchange acidity than the sum of bases plus KCl extractable Al, would suggest a soil with a high pH dependent charge. This can be used along with bulk density to separate subgroups that have high levels of amorphous materials commonly derived from pyroclastics.

RELATIONSHIP OF 15 BAR TO CLAY

This value is used to give an indication of dispersion in the particle size determination. If the ratio is greater than 0.6 and soil related factors do not adequately explain the situation, incomplete dispersion in particle-size analysis may be the cause. Some soil-related factors that can cause deviation from the 0.4 reference point are: activity clays (kaolinites, chlorites, and some micas) tend to lower the ratio to 0.35 or below. Clay-size carbonate tends to decrease the ratio. Organic matter increases the 15-bar water content. Amorphous mineral materials increase the ratio.

In "Soil Taxonomy," 1975 the inverse of this ratio is sometimes used. A ratio of < 1.25 used in conjunction with bulk density and poor dispersion qualities; i.e., ones high in amorphous materials.

PH DEPENDENT CHARGE
The pH dependent charge as inferred by "Soil Taxonomy," 1975 is the difference between the sum of bases (NH,OAc) plus extractable acidity and sum of bases (NH₄OAc) IN KCl exchangeable Al, divided by measured clay or by 15-bar water times 2.5 under conditions defined in "Soil Taxonomy," 1975.

LITERATURE CITED

- Blakemore, L. C., P. L. Searle, and B. K. Daly. 1981. Soil Bureau Laboratory Methods. A. Methods for Chemical Analysis of Soils. New Zealand Soil Bureau Scientific Report 10A. DSIRO, New Zealand.
- Bray, R. H. and L. T. Kurtz. 1975. Determination of total, organic and available form of phosphorous soil. Soil Sci. 59:39-45.
- Ecuador. Agrarian Regionalization National Program, National Resources of Ecuador Information made by PRONAREG-ORSTOM Quito Agriculture Ministry, 1973-1983.
- Ecuador, PRONAREG Soil Inventory of Ecuador made by Soil Department, Quito Agriculture Ministry, 1973-1983.
- Ecuador, CENTRAL BANK, Atlas of the World, part of Ecuador national information in natural resources prepared by PRONAREG, Quito, 1979-1983.
- Newhall, F. 1972. Calculation of soil moisture regimes from climatic record. Rev. 4. Mimeographed, 17 pages, 3 tables, 7 figures. Soil Conservation Service, USDA, Washington, D.C.
- U.S. Department of Agriculture, Soil Conservation Service. 1972. Soil Survey Laboratory Methods and Procedures for Collecting Soil Samples. Soil Survey Investigations Report No. 1. Government Printing Office, Washington, D.C.
- U.S. Department of Agriculture, Soil Conservation Service. 1975 Soil Taxonomy. Agriculture Handbook No. 436. U.S. Government Printing Office, Washington, D.C.
- U.S. Department of Agriculture, Soil Conservation Service. 1982. Soil Survey Laboratory Methods and Procedures for Collecting Soil Samples. Soil Survey Investigations Report No. 1. Government Printing Office, Washington, D.C.
- Van Wambeke, A. 1981. Calculated soil moisture and temperature regimes of South America. SMSS Technical Monograph No. 2. Soil Conservation Service, USDA, Washington, D.C.

THE VOLCANIC ASH SOILS OF ECUADOR

Edmundo Maldonado* Fausto Maldonado** Claude Zebrowski***

Soil Scientist, PRONAREG (Programa Nacional de Regionalización Agraria).

^{**} Agricultural Specialist, USAID Mission to Ecuador.

*** Advisor Soil Scientist, ORSTOM (Office por la Recherche Scientifique et Technique d'Outre Mer, France).

Introduction

According to Wolf (1) Ecuador is one of the most interesting countries for vulcanologic studies, and before him, Alexander von Humboldt could already understand the importance of Ecuadorean volcanoes and their effects. Reiss and Stubel, German vulcanologists, choose Ecuador for their studies on pyroclastics and volcanic rocks (1870 1874).

In the present century, volcanic ash soils have attracted the attention of researchers who have seen in them interesting characteristics; most visible of all was their very dark, almost black, low chroma, color. This led both to Miller (2) and Frei (3) to call the highland Sierra soils, Suelos Negros Andinos, a name which was locally used for several years.

Later, when elaborating the Soil Taxonomy System, other characteristics were added to these already known, to identify volcanic ash soils, which allowed their inclusion within a specific class of soils.

Several authors have contributed, since 1956, to a better knowledge of volcanic ash soils in Ecuador, with information and data to improve their use and management, and to complete their classification (4, 5, 6, 7, 8).

Since 1973, it has been PRONAREG, the Programa Nacional de Regionalización Agraria, of the Ministry of Agriculture, which has been in charge of the soil survey of the country, with the technical assistance of ORSTOM. During the development of this inventory, several problems have arisen which obliged to modify map legends. These will be discussed during the Ecuador part of the VI Soil Classification Workshop.

1. The Volcanic Terrain in Ecuador

From $1^{\rm O}$ North latitude, at the border with Colombia, to about $2^{\rm O}$ 30' South the Andes Cordillera shows a number of volcanoes, some of them reaching more than 5000 m above sea level.

The Andes Cordillera origin is known as the result of American Continent drift to the west, which caused the uplift of deep magma and the subsequent production of basaltic lavas (in Ecuador, this basic rock formation has been identified with the names of Piñón and Macuchi). These are pillow lavas, which indicate a marine environment for deposition.

On the other side, to the east, the crystalline rocks of the Guyana-Brazil Shield, started to move up too.

Two large areas had already been individualized as geological regions in the area: one with a basic volcanic basement to the west and north, and another of a metamorphic basement to the East and South (present Amazonian Region and Cordillera Oriental or Real).

But it was in the Tertiary when a generalized uplift occurred in the Andes, when both a western and an eastern range were formed. The former was made from volcanic and sedimentary terrains and the latter as an axis of metamorphic compressed rocks in overthrust foldings. At the same time, faulting was active on both sides of each range, thus creating parallel horsts going in a North-east to South west direction, with a grabben in between.

According to SAUER (9), the Andes Cordillera, after a series of vertical movements and erosion phases, at the end of the Tertiary reached only 2000 to 2500 m above sea level. Then, high magnitude movements started in the Pleistocene. These vertical movements affected different areas in the country, but their magnitude was higher to the north of parallel 2° South.

The differential uplifting caused the rupture of rock masses both with faults along the mountain ranges, and transversal to them, thus creating basins which drain either to the East or to the West. Through the lower resistance points in the Earth's crust, along the cordilleras, magma went up to the surface to reestablish volcanic activity. It is at that time that most of the volcanoes we now know were formed and continued growing, in some cases even in the present time, reaching more than 4500 m altitude.

During the Quaternary, volcanic activity shifted form the western to the eastern range where now we find the most active volcanoes (Cotopaxi, Tungurahua, Sangay, Reventador).

According to SAUER (9), the Western range is formed by the "Pacific petrologic Type" of rocks (sialic-lithogenic). These rocks are more acid (dacitic), than the ones in the eastern range, which are of the "Atlantic type" (simic-Hypogenic) and which are usually formed in the most recent phases of volcanic activity.

Volcanoes were most frequently producing andesitic or dacitic materials, and gave place to both lava flows and

pyroclasitc materials of various sizes, although they have not too different composition. Some fine, wind blown materials are probably at the origin of the "cangahua", the consolidated tuff similar to loess, which is found either below more recent deposits or outcropping where the topsoil has been eroded (10).

2. Climate

Being a tropical (equatorial) country, mean temperature is almost constant during the year, so soil temperature regimes are all "iso" regimes. Nevertheless, temperature varies with altitude producing temperature regimes from the isohyperthermic to the isofrigid. Temperature limits for the various regimes have been modified in Ecuador, being adjusted according to the temperature growing limit of crops:


- 10° limit, minimal crop temperature corresponds to "frigid";
- 13^o limit between the "mesic" and the "thermic" regimes, which corresponds to the growing limit of maize;
- 20 22° for "hyperthermic", coincides well with the tropical crops line.

Rainfall shows important variations as it can be seen in the map (Fig. 1). In the Coastal area, rainfall increases from the Southwest, in Salinas (less than 500 mm), to North East (close to 6000 mm) in the East of Esmeraldas Province. In the Amazonian Region, rainfall is more homogeneous (usually higher than 2000 mm per year).

Abundant rainfall to the footslope of the mountain ranges, decreases with altitude, but with lower temperatures, evaporation also decreases, resulting in a compensation for rainfall reduction, and makes more humid the high altitude areas (páramos).

In the Interandean valley, rainfall decreases in the valleys, which have a comparatively dry climate.

More than the total year rainfall, their distribution determines volcanic ash soil evolution. This distribution directly influences the soil moisture regime, whose spatial variations are indicated in the rainfall map.



REGIMENES DE TEMPERATURA Y HUMEDAD TEMPERATURE AND MOISTURE REGIMES ${\tt FIG},\ 2$ LEYENDA frigid mesic thermic hiparthermic humedod udic Hu uStic Н \$ Fuente PRONAREG - ORSTOM oridic Нc

3. Influence of the main genetic factors on soil characteristics

The short range of variation among the various volcanic materials (lapilli, dacitic ashes) cannot explain alone the large evolution differences in soils (Colmet-Daage, 5).

On the contrary, age and particle size have a more obvious influence. Thus, the evolution difference between Amazonian soils (Hydrandepts from Puyo) and the soils of the moist area of the Guayas basin (Dystrandepts of Santo Domingo and Los Bancos) is the result of age differences in the volcanic materials ejected between those regions.

In a similar way, variations in soil characteristics of northern and central soils (close to volcanoes) and the southern soils (far from them), are due to age differences between them, although it is also possible that particle size, finer in the south than in the north will play an important role in soil genesis, explaining the differences noted.

Rainfall and mainly the duration of the humid season, as it is already known (6,17), have a primary role in volcanic ash soils' genesis: in moist areas, pedogenesis leads to individualization and preservation of amorphous materials in the soil, while in drier areas clay minerals are formed (mainly halloysite). So in the Ecuadorean Coastal Area, in a climosequence, the distribution of Dystrandepts, Eutrandepts and Mollisols is a typical example of the influence of the dry season on soils genesis.

Likewise, in the Highland Sierra, Andepts are located in the humid areas, i.e. in the high zones, while in the lower basin areas, which are drier, pedogenesis, with clay minerals formation, leads to the formation of Mollisols. In addition to the rainfall influence on the genesis of clay minerals, it is important to emphasize its role on base leaching and consequently on base saturation, pH values and carbonate accumulation in soils; all characteristics whose changes are strong and rapid, especially in the Sierra, where striking climate changes are seen on rather short distances.

Temperature, in addition to its influence on the chemical reactions and the weathering rate of volcanic materials, also has an important role in the accumulation and composition of organic matter; so humic and fulvic acids show similar contents in the cold areas, while in the warm ones, fulvic acids are more abundant.

4. Soils

The Soil map legend established for the Soil Survey of Ecuador has taken into consideration two important soil features: particle size distribution and mineralogy, which reflect the evolution degree of soils. So, Colmet-Daage et al. (11) have defined four great sets of soils:

- Sandy soils
- Sandy loam soils
- Clayey soils
- Soils with amorphous materials

Additionally, taking into account the subsoil, another set was established for soils developed on "cangahua" (Fig. 3).

a) Sandy soils. With the exception of alluvial soils located in the Eastern plain between Lago Agrio and Coca, sandy soils have developed from wind-blown volcanic ejecta. These are not evolved soils, either because they are young, formed on recent materials, or because they are located in areas where drought reduces weathering. They may be found close to recent volcanoes like Tungurahua, Quilotoa, Cotopaxi, Cuicocha, or in dry areas like San Antonio de Pichincha, Latacunga, Riobamba, or Palmira. In the latter area, as behind the Chimborazo Mountain, soil erosion renewes soil materials and allows the presence of sandy soils.

In addition to a sandy texture ranging from fine sand to gravel, these soils are characterized by a 1/3 bar water retention of less than 20% and their organic matter content.

- Soils with less than 1% Organic matter in the upper 20 cm have been classified as PSAMMENTS.

In addition to the soils on medium to coarse lapilli, close to recent volcanoes (Cotopaxi, Tungurahua) over 3600 m, i.e. under a frigid regime, most of these soils are at lower altitudes under isothermic regimes.

For these soils, pH values range from less than 6 to higher than 8, showing carbonate mycelia in very dry areas.

 Soils with more than 1% organic matuer were classified as VITRANDEPTS. In these soils, there is a beginning of sand weathering, either because deposits

SUELOS SOBRE PROYECCIONES VOLCANICAS SOILS FORMED ON VOLCANIC ASH DEPOSITS Esmeroidos FIG. 3 LEYENDA - Orgánicos Afteramientos rocosos Fuente' Mopas de suelos PRONAREG-ORSTOM

are older or because they are in moister zones, something that contributes to organic matter accumulation.

To this early weathering corresponds the formation of amorphous materials in moist areas and of clay minerals (halloysite or 14 A minerals) in dry areas. pH and base saturation values depend on the climate under which soils are evolving (Table 1).

Table 1. pH Values and Base Saturation (B. S.)
According to (imatic Zones.

Moisture regime rature reg.	Ustic	Udic
Isomesic		pH < 6
Isothermic	pH - 7 B. S. > 50%	B. S. < 50%

b) Loamy soils. These are more evolved soils either located in moister areas, or they are far away of the emission centers of ashes, or even they are developed from older pyroclastic materials.

In the Sierra, they are located in the Interandean valley at altitudes lower than 3600 m a.s.l., mostly between 2200 and 2300 m; in the Coastal area, they are found on recent deposits in the relatively dry area between Quevedo and Balzar.

These are soils whose characteristics are those of a Mollisol and locally of an Eutrandept. Thus, color is dark brown to black, texture is loam to sandy loam. Clay is essentially halloysite with occasionally mont-morillonite, or amorphous products, the latter appearing in the moister areas.

Base saturation is higher than 50%; bulk density values range from .8 to 1.3 and NaF test is mostly negative, except when amorphous materials are present. Water retention at 1/3 bar shows different values, related to texture. For instance, for a first sandy loam subset, water retention is between 10 and 20%, while for a loam subset water retention is higher than 20%.

For each subset, as indicated in the following tables, main soil characteristics vary with climate: so pH, Base saturation, bulk density, MaF reaction, color and organic matter content vary according to moisture regimes, and the two latter characteristics also depend on temperature.

Table 2. <u>Values of Certain Characteristics of Soils According to Climatic Zones</u>

- Sandy loam soils: water retention at 1/3 bar: 10 to 20%

Tempe- regime rature reg.	Ustic	Udic
Isomesic	- strong black color chroma: 0 - 1 NaF negative B.D. ~ 1.2-1.3 - Carbonate mycelia	color chroma: 0 - 1 NaF positive
Isothermic		- dark brown color chroma 1 - 2 - O.M. > 3% - pH = 6

- Loamy soils: water retention 1/3 bar 20%

Moisture		
Tempe- regime rature reg.	Ustic	Udic
	- black soil	- strong black
Isomesic	- NaF negative	- NaF positive
	B.D.≃ 1.2	- B.D89
		- B.S. ≈ 50%
	- O.M. < 3%	- O.M. > 4%
	- pH ≃ 7	- pH ≃ 6
Isothermic	- Carbonate	- B.D. ≃ 1
	mycelia	- NaF negative
	local	- B.S. ≃ 50%
	- pH ≈ 6.5	- pH ≃ 6.5
Isohyperthemic	- B.D.:.8 - 1	- B.D.:.8
	- B.S.>60%	- B.S. 50-60%

c) Clayey soils. These are black soils with a mollic epipedon and base saturation higher than 50%. They are MOLLISOLS, with more than 30% clay (halloysite or montomorillonite).

In the Sierra, they occupy small areas between 2200 amd 3200 m, that is to say they have an isothermic temperature regime. They have developed on old ashes, sometimes hard, cangahua deposits, with moisture regimes

going from ustic to udic. In the Coast, they are found south of El Empalme in an area under ustic regimes; at the Dystrandept-Mollisol transition the most clayey soils of the climosequence are found.

d) Soils with amorphous materials. They are found in the eastern plain and in the moist area of the Guayas basin; in the outside hillslopes and in the high altitude areas of the mountain ranges, i.e. in moist areas under udic or perudic regimes, having as a consequence that soil moisture at any time is higher than water retention at 15 bar pressure.

They are soils with a bulk density lower than .8 in the upper 35 cm, and with base saturation lower than 50%.

Several characteristics, such as water retention and organic matter have permitted the separation of several subsets.

Soils with water content lower than 100% (1/3 bar) are Dystrandepts whose secondary properties, as indicated in the table below, depend on temperature. When they are under isofrigid regime, they are Cryandepts.

Table 3. Soil Properties According to Temperature Regime

Temperature Regime	Isofrigid	Isomesic	Isothermic	Isohyperthermic
Color	- black chroma: 0 - 1	- black chroma: 0 - 1	- dark over 50 cm chroma: 1 - 2	- dark over 10 to 20 cm
Base Saturation	very low		low	
Humic and fulvic acids	H. A. > F. A. Especially in depth			H. A. < F. A.
Нд	pH water > pH KCl			pH water > pH KCl

Among these Dystrandepts, those with a lower water retention (20 to 50%) are younger soils with sandier textures and higher bulk density (close to 1) or soils located in drier areas which have a lower content of amorphous materials.

- Soils with water retention (1/3 bar) higher than 100% have been classified as Hydrandepts or Hydric Cryandepts. They are strongly smeary, with a very low bulk density (0.3 to 0.5). They have a base saturation lower than 10%, and their reaction to NaF is very active. Organic matter content ranges from 10 to 20% in the upper 30 cm. Color, pH and fulvic & humic acids percentages change according to temperature, as in the previous soils.

These Hydrandepts whose water retention can reach 300% have developed from lapilli and old ash deposits in very humid areas, as the eastern zone and the northeast of Esmeraldas province.

Organic soils. These are black soils (chroma 0 - 2) which have all the properties of the previously described soils, but with a higher organic matter content: this is higher than 20% throughout the profile and sometimes it can reach 30% in the top horizon. Water retention is always high (higher than 100%) and base saturation very low (lower than 10%), with a sum of bases lower than 3 meq/100 g soil in the topsoil and 1 meq/100 g soil in depth.

These are soils developed from fine ashes found at very long distances from volcanoes; this explains why they are frequent in the high altitude zones south of Cañar to Saraguro "páramos". They might be classified in a "histic" subgroup of Hydrandepts.

e) Soils on Cangahua. They are very loamy to very fine sandy soils developed on a hard layer (cangahua) which appears at less than 1 m depth.

They are found in the Interandean valley, in the lower parts of the basins and in the internal slopes to altitudes of 3200 to 3500 m. Higher than these altitudes cangahua disappear.

In the lower and drier areas, soils are not deep, because of erosion; in many cases, soil has been lost and the cangahua outcrops. Soil is sandy with very low organic

matter content, pH is higher than 7 and in the drier areas (transition regime from ustic to aridic) there is carbonate accumulation as mycelia. There is an abrupt contact with cangahua and the latter is not weathered. The soil is a DURUSTOLL and cangahua is considered as a duripan.

Little by little, and with the increase in moisture in the higher zones because of a decrease in evapotranspiration (in some cases with increase in rainfall also), soil color becomes darker due to higher organic matter content, and weathering is higher. Then we have clays such as Halloy-site and Montmorillonite, an argillic horizon is formed with coatings of very dark color (black) at the contact zone of cangahua and upper deposits. The presence of this layer is caused by cangahua weathering and clay illuviation as has been shown by the coatings. pH is slightly acid, humic acids content is higher than fulvic acids, especially in the argillic horizon. These soils have been classified as DURIUDOLLS.

Conclusions

The various studies made on volcanic ash soils in Ecuador have permitted to improve the knowledge of these soils which cover about 31% of the country. They have been completely mapped by PRONAREG.

Soil map legends were established taking into consideration many characteristics (texture, mineralogy, water retention, organic matter and sorption complex), whose changes depend on the climate, especially on moisture; age of the materials and the distance to emission centers are of minor importance.

Soil classification, not yet in its final form, was developed, in some cases, using simple criteria of land use but not utilized by the Soil Taxonomy. Water retention at 1/3 bar, for instance has a limit of 100% to separate Dystrandepts from Hydrandepts, as the organic matter content of 1% is the limit between Vitrandepts and PSAMMENTS. These criteria and value, however, must still be discussed and better defined.

REFERENCES

- Wolf, T. 1982 Geografía y Geología del Ecuador, F.A. Brockhaus, Leipzig.
- Miller, E. 1948 Ecuadorean Soils and some of their Fertility Properties. Ph.D. Thesis Cornell University, Ithaca, N.Y.
- Frei, E. 1957 Informe al Gobierno del Ecuador sobre reconocimientos edafológicos exploratorios, FAO, Reporte No. 585, 35 p. Rome.
- Maldonado, F. 1973 Study of a Climo-Toposequence on Volcanic Ash Soils in Ecuador, Ph.D. Thesis, Rijks universiteit, Ghent, Belgium.
- 5. Wright, A.C.S. The "Andisols" or Humic Allophane Soils of South America in Meeting on the Soil Classification and Correlation of Soils from Volcanic Ash, FAO, World Soil Resources Reports No. 14, pp. 9-22.
- 6. Colmet-Daage, F. 1967 Caractéristiques de quelques sols d'Equateur dérivés de cendres volcaniques, I, Essai de characterization des sols des régions tropicales humides. Cah. ORSTOM, Sér. Pedologie, 5 (1): 1-38.
- 7. Colmet-Daage, F. et. al. 1967 Caractéristiques de quelques sols d'Equateur dérivés de cendres volcaniques, II, Conditions de formation et d'évolution. Cas ORSTOM. Sér. Pedologie 5 (4): 353-392.
- 8. Colmet-Daage, F. 1969 Caractéristiques de quelques sols d'Equateur dérivés de cendres volcaniques, III, Comparaison de l'évolution de quelques sols des régions tropicales chaudes et tempérés froides d'altitude. Cas ORSTOM. Sér. Pedologie 7 (4): 495-560.
- 9. Sauer, W. 1971 Geologie von Ecuador, Gebriider Borntralger, Belgin, 316 p.
- 10. Hoffstetter, R. 1956 Ecuador in Léxique Stratigraphique Internationale, 5a. Amerique Latine, Parix.
- 11. Colmet-Daage, et. al. 1976 Leyenda de la Carta Nacional
 de Suelos, PRONAREG.

Pedon Description and Analytical Data

The section contains, for each pedon, a description for the pedon sampled, analytical data in two parts: the first sheet is the standard NSSL laboratory data sheet and the second shows the results of the mineralized analysis.

Pedons that were sampled but not included in the tour

part of the guide are in Appendix II.

If there are questions on the data sheet, Appendix III

should be checked for an explanation.

The pedons are placed in the order that we will visit them. No classifications have been placed on any of them. If you have a chance, it would be useful to try and work through their classifications before we visit them in the field.

The location of each pedon is shown on Figure 11.

Date:

6/83

Series: Ecuador No. 1 Pedon No: S83FN-325-001

Taxonomy: Loamy over cindery, mixed, isomesic cumulic Hapludolls Latitude: 0°31'00" Longitude: 78°37'10"

Location: Tour stop No. 1. Machachi-Aloasi area. Physiography: Hillside in mountains.

Geomorphic Position: Hillslope

Slope and Aspect: 10% ENE convex-convex Elevation: 3200 m M.S.L

Microrelief: none

Air Temp. Summer: 11.0°C Winter: 10.9°C Precipitation: 1200 mm Udic moisture regime. Water Table: > 10 M

Drainage: Well drained Stoniness: nonstony Permeability: moderate

Land Use: Small grains of wheat and barley

Erosion or Deposition: Slight sheet and rill erosion

Parent Material: moderately weathered, ask and unweathered pumice deposits
Described by: T. Cook, G. Del Posso, and Soil Survey Staff.
Remarks: Date described: January 25, 1983. Field Moisture, 0-4 cm dry. 4-115 cm slightly moist. 115-180 cm very slightly moist.

Vegetation: before temperate forest (chilea, fencus, eucalyptus grove).

- Ap 0 21 cm Black (10YR 2/1, rubbed), fine sandy loam; grayish brown (10YR 5/2, dry); plowed soft clods; very friable, non-sticky and non-plastic; common very fine roots; abrupt smooth boundary.
- 47 CM Very dark brown (10YR 2/2) fine sandy loam; weak medium and coarse angular blocky structure; very friable, ron-sticky and slightly plastic; few very fine, fine, and medium roots; many very fine and few fine tubular pores; 2% 2-5 mm pumice fragments; gradual smooth boundary. The upper part of the horizon is a dense compacted tillage pan.
- A2 47-75 cm Black (N 2/0) gravelly sandy loam; weak coarse and very coarse subangular blocky structure parting to weak coarse and very coarse granular; very friable, non-sticky and non-plastic; few very fine, fine, and medium roots; many very fine and common fine and medium tubular pores; coarse fragments, 15% 2-50 mm pumice; abrupt wavy boundary.
- 75 84 cm Very pale brown (10YR 7/4) and dark yellowish brown (10YR 4/4) very gravelly coarse sand; single grain; loose, non-sticky and non-plastic; few very fine and coarse roots; many very fine and fine interstitial pores; coarse fragments, 45% 2-45 mm pumice; abrupt wavy boundary. Pumice strata is discontinuous in the pit and varies from 4-10 cm in thickness.
- 84 115 cm Black (10YR 2/1) fine sandy loam; weak medium and coarse subangular blocky structure parting to weak medium and coarse granular; friable, non-sticky and non-plastic; few very fine roots; common very fine and fine and a few medium tubular pores; 3% 2-10 mm pumice fragments; diffuse wavy boundary.
- 115 150 cm Very dark brown (10YR 2/2) fine sandy loam; weak medium and coarse subangular blocky structure parting to weak medium and coarse granular; friable, slightly sticky and slightly plastic; few very fine roots; common very fine and fine and few medium tubular pores; 2% 2-15 mm pumice fragments; diffuse wavy boundary.
- Very dark brown (10YR 2/2) loam; weak medium and coarse 150 - 182 cm subangular blocky structure parting to weak medium and coarse granular; friable, slightly sticky and slightly plastic; few very fine roots; common very fine and fine tubular and a few medium tubular pores; 2% 2-15 mm pumic fragments; gradual wavy boundary.
- 182 205 cm Dark brown (7.5YR 3/2) and brown to dark brown (7.5YR 4/4) loam; massive; friable, slightly sticky and slightly plastic; few very fine roots; common very fine and fine and few medium tubular and many very fine interstital pores; 2% 2-15 mm pumice fragments; about 50% of the horizon is very firm moist, is weakly cemented and broken in hands with difficulty; material looks like weathered tuff.

PAGE 1 OF 4 PAGES

U. S. DEPARTMENT OF AGRICULTURE

NATIONAL SOIL SURVEY LABORATORY

25.9

1.32 1.34 0.005

9.3 0.20

28.1 12.5 0.20

SOIL CONSERVATION SERVICE

LINCOLN, NEBRASKA 68508

ECUADOR 001

ECUADOR-SMSS

832110

8 0.55

LAB CLASSIF: LOAMY OVER CINDERY, MIXED, ISOMESIC CUMULIC HAPLUDOLL

S 83FN-325 -001

DATE 09/25/84

1.2

0.1

SAMPLE NO. 83P2103-2110 PEDON NO. 83P 459 PROJECT NO. 83P 83

GENERAL METHODS 181A, 2A1, 2B

-3-- -4-- -5-- -6-- -7-- -8-- -9-- -10- -11- -12- -13- -14- -15- -16- -17- -18- -19- -20-(- - -TOTAL - - -)(- -CLAY- -)(- -SILT- -)(- - - - - - -SAND- - - - - -)(-COARSE FRACTIONS(MM)-)(>2MM CLAY SILT SAND FINE CO3 FINE COARSE VF F M C VC - - - - WEIGHT - - - WT LT LT .002 .02 .05 .10 .25 .5 1 2 5 20 .1- PCT 0 .0002 .002 -.02 -.05 -.10 -.25 -.50 -1 -2 -5 -20 -.75 .75 WHOLE --- PCT OF <2MM (3A1) --- --- --- <-- PCT OF <75MM(3B1)-> SOIL SAMPLE HZN DEPTH HORIZON LT .002 .05 LT NO. NO. (CM) .002 -.05 -2 .0002 .002 -.02 -.05 -.10 -.25 -.50 **** SEE FOOTNOTE REGARDING PSDA BELOW **** 832103 18 0- 21 AΡ 4.9 35.4 59.7 21.3 14.1 16.1 20.5 13.8 7.1 2.2 832104 lili 2S 21- 47 1 A1 5.0 34.3 60.7 20.4 13.9 15.2 19.3 14.1 8.6 3.5 4V 832105 49 7 3S 47- 75 A2 2.5 26.1 71.4 14.7 11.4 15.6 16.4 15.4 14.0 10.0 9 6 27V 832106 45 75- 84 2C1 74 42 3.7 95.5 0.8 1.7 2.0 1.3 29.8 7.2 33.7 23.5 11 832107 10 54V 99 75 5S 84-115 3AB: 6.1 42.3 51.6 26.9 15.4 16.2 16.2 10.1 6.9 2.2 832108 6٧ 40 6S 115-150 **3AB2** 7.3 36.9 55.8 23.1 13.8 15.0 20.0 11.6 6.9 2.3 4٧ 832109 44 7S 150-182 3ACB 9.2 32.2 58.6 20.3 11.9 16.0 20.3 13.7 6.8 1.8 1 4V 832110 8S 182-205 47 3C2 6.3 32.1 61.6 18.7 13.4 16.3 20.8 15.5 ORGN TOTAL EXTR TOTAL (- - DITH-CIT - -)(RATIO/CLAY)(ATTERBERG)(- BULK DENSITY -) COLE (- - -WATER CONTENT - -) WRD EXTRACTABLE 15 - LIMITS - FIELD 1/3 OVEN WHOLE FIELD 1/10 SAMPLE HZN 15 WHOLE FE AL MN BAR LL PI MOIST BAR DRY SOIL MOIST BAR BAR SOIL NO. 6A1C 6B3A BAR NO. 6R3A 6C2B 6G7A 6D2A 8D1 8D1 4F1 4F 4A3A 4A1D 4A1H 4D1 4B4 4B1C 4B1C <- - - - - - - - PCT OF <2MM - - - - - -> 4B2A 4C1 PCT <0.4MM <- - G/CC - - -> CM/CM <- - -PCT OF <2MM - -> CM/CM 832103 1 2.66 0.321 0.4 0.2 2.20 1.35 832104 2 2.34 0.184 6.6 0.4 0.2 2.12 1.48 1.23 1.25 0.005 832105 3 2.28 21.8 7.4 0.17 0.2 0.5 4.56 2.72 1.11 1.11 27.8 0.18 832106 6.8 4 0.21 0.3 TR 3.13 3.25 832107 5 1.13 0.5 0.1 1.44 0.92 1.29 1.29 25.8 832108 6 1.15 5.6 0.25 0.6 0.1 1.40 0.85 1.35 1.35 22.5 6.2 0.21 832109 7 0.96 0.1 1.26 1.01 1.23 1.27 0.010

1.79 1.98

^{*} CAUTION SHOULD BE USED WITH PSDA DATA WHEN USED FOR MAKING ENGINEERING INTERPERTATIONS BECAUSE OF INCOMPLETE DISPERSION OF AMORPHOUS MATERIAL. THEY ARE PRESENTED TO AID IN CLASSIFICATION AND ASSESSMENT OF MINERALOGY. REFER TO PEDON DESCRIPTION FOR

```
PAGE 2 OF 4 PAGES
       ECUADOR 001 S 83FN-325 -001 DATE 09/25/84 PEDON NO. 83P 459 NATIONAL SOIL SURVEY LABORATORY
-1-- -2-- -3-- -4-- -5-- -6-- -7-- -8-- -9-- -10- -11- -12- -13- -14- -15- -16- -17- -18- -19- -20-
          (- NH4OAC EXTRACTABLE BASES -) ACID- EXTR (- - - - CEC - - -) AL -BASE SAT- CO3 AS RES. CA MG NA K SUM ITY AL SUM NH4- BASES SAT SUM NH4 CACO3 OHMS
                                                                                                   COND_{-}(- - - - PH - - - 1)
                                                                                                   MMHOS NAF CACL2 H20
SAMPLE HZN 5B5A 5B5A 5B5A BASES
                                                 CATS DAC ++ AL
                                                                              OAC <2MM /CM
                                                                                                  /CM
                                                                                                              0.1M
       NO. 6N2E 602D 6P2B 6Q2B 6H5A 6G9A 5A3A 5A8B 5A3B
  NO.
                                                                 5G1
                                                                         5C3 5C1 6E1G 8E1
                                                                                                   81 8C1D 8C1F 8C1F
           1:2 1:1
832103
            5.9
                 0.9
                       0.1
                                 7.3 24.2
                            0.4
                                              TR 31.5 10.8
                                                                               68
                                                                                                          9.9
                                                                                                               5.2
                                                                                                                     5.4
832104
            6.6
                 1.1
                       0.1
                            0.3
                                  8.1 27.8
                                             -- 35.9 10.6
                                                                         23
                                                                               76
                                                                                                         10.1
                                                                                                               6.0
                                                                                                                    6.3
832105
            6.9
                 1.6
                       0.2
                            0.2
                                  8.9 21.5
                                             TR 30.4 11.4
                                                                         29
                                                                              78
                                                                                                         10.2
                                                                                                               5.9
                                                                                                                    6.2
832106
            1.7
                 0.7
                       0.1
                            0.2
                                  2.7
                                       8.1
                                              -- 10.8
                                                       2.5
                                                                         25
                                                                            100
                                                                                                          8.2
                                                                                                               5.7
                                                                                                                     6.3
832107
            6.7
                                             -- 12.9
                 1.4
                       0.3
                            0.4
                                 8.8
                                       4.1
                                                       8.8
                                                                         68 100
                                                                                                          9.7
                                                                                                                     6.7
832108
            7.5
                       0.3
                 1.5
                            0.4
                                 9.7
                                       6.4
                                             TR 16.1 10.2
                                                                         60
                                                                              95
                                                                                                          9.5
                                                                                                               6.1
                                                                                                                    6.8
832109
            8.1
                 1.7
                       0.3
                            0.6 10.7
                                       6.3
                                             0.1 17.0 11.6 10.8 1
                                                                               92
                                                                         63
                                                                                                          9.4
                                                                                                               6.4
                                                                                                                     6.8
832110
           7.0
                1.7
                       0.4
                            0.8 9.9
                                       6.9
                                            -- 16.8 11.3
                                                                         59
                                                                               88
                                                                                                                     6.6
          (- - - - SPODIC HORIZON CRITERIA - - - - )
                                                                            (------)
          (- -NA PYROPHOSPHATE EXTRACTABLE- -) INDEX
                                                                            (-----)(---)
            C FE AL FE+AL FE+AL AL+C OF
                                                                            (- - - - X-RAY - - - -)(- -DTA - -) TOTAL DOM
SAMPLE HZN
                          (- -DIVIDED BY- -) ACCUM
                                                                            (- - - - - <2U - - - -) RES WEATH
       NO. 6A4A 6C8A 6G10 DI-CI PCT PCT
                                                                             7A21 7A21 7A21 7A21 7A3 7A3 7B1A 7B1A
 NO.
           <- PCT OF <2MM-> FE+AL CLAY CLAY
                                                                             <- RELATIVE AMOUNTS -> <- - - - - PCT - - - ->
832103
                 0.1
                       0.2
                            0.5
                                  0.1
                                             610
832104
                 0.1
                       0.2
                            0.5
                                 0.1
                                             868
                                                                             VR 1 NX 6
                                                                                                                 2 GA39
832105
                       0.2
                 0.1
                            0.4
                                  0.1
                                             816
802106
832107,
832108
                       TR
                                                                             NX 6
                 0.1
                       0.1
                           0.3
                                  TR
                                             305
                 0.1 0.1
                            0.3
                                  TR
                                             436
                                                                             FD 1 NX 6
832109
                       0.1
                            0.1
                                  TR
832110
                       0.1
                           0.1
                                  TR
```

FAMILY CONTROL SECTION: DEPTH 25-100 PCT CLAY 4 PCT .1-75MM 63. SPODIC HORIZON: INDEX OF ACCUMUL CANALYSES: S= ALL ON SIEVED <2MM BASIS V= 75-20MM FROM VOLUME ESTIMATES MINERALOGY: KIND OF MINERAL VR VERMICULITE NX AMORPHOUS GA GLASS AGG FD FELDSPAR

RELATIVE AMOUNT 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

S U P P L E M E N T A L D A T A S H E E T

		EXT P BRAY1	15 BAR MOIST	P RETENT	(AC: AL	ID OXAL SI	ATE) FE	EXTR AL (KOH)	pH KCl
LE	HZN NO.	6S3 PPM	4B2b <	*	+ -PCT OF	* <2MM-	* ·	>	8Clg
03 04 05 06 07 08 09	1 2 3 4 5 6 7 8		6.9 8.8 8.5 2.6 8.9 8.8 12.4 17.5	36 49 50 49 36 34 32 36	0.4 0.5 0.5 0.1 0.3 0.2 0.2	0.2 0.3 0.4 0.1 0.3 0.2 0.2	0.4 0.4 0.4 0.3 0.3 0.3 0.4	0.3 0.3 0.4 <0.1 0.1 0.1 0.1	

W ZEALAND PROCEDURE

CP83FN128

		(PTICAL	,)	() (DTA) (TOT A	
SAMPLE NO.	HZN NO.	FA 7BlA	RE 7Bla	7BlA	7B1A	7B1 A	7BlA	7BlA	7BlA	7BlA	7Bla	7BlA	K20	FE 6C7A
83P2103 83P2104 83P2105	2	VFS	2	GA39	FP24	GS17	FG 7	HN 4	PR 4	HG 2	OP 2	0G<1	VR 1 NX 6 0.5	1.6
83P2106 83P2107	4												NX 6 0.3	1.3
83P2107 83P2108 83P2109	6 7												FD 1 NX 6 0.4	2.5

ANALYSES: S=ALL ON SIEVED < 2mm BASIS

MINERALOGY: FA = FRACTION ANALYZED RE = RESISTANT

KIND OF MINERAL: OP = OPAQUES FP = PLAGIOCLASE FELDSPAR GA = GLASS AGGREGATES GS = GLASS HN = HORNBLENDE PR = PYROXENE

VR = VERMICULITE NX = NON-CRYSTALLINE HG = GLASS COATED HORNBLENDE FG = GLASS COATED FELDSPAR

OG = GLASS COATED OPAQUE FD = FELDSPAR

RELATIVE AMOUNT: 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

MINERALOGY BASED ON SAND/SILT:

MINERALOGY BASED ON CLAY:

FAMILY PLACEMENT:

COMMENTS:

Series: Ecuador No. 9 85 Date: 8/82

Pedon No: S83FN-325-009

Taxonomy: Fine-loamy, mixed, isomesic Andic Humitropept

Latitude: 0°32'15" Longitude: 78°37'10"

Location: This pedon is located about 60 km south of Quito west of Machachi on Corazon Volcano (Aloasi zone pit two of Machu sequence).

Physiography: Mountains

Geomorphic Position: On lower foot slopes

Slope and Aspect: 25% E convex-convex Elevation: 3420 m M.S.L

Microrelief: Smooth

Air Temp. Summer: Winter:

Precipitation: 1400 mm Udic moisture regime.

Water Table: > 10 m

Drainage: Well drained Permeability: moderate

Stoniness: nonstony Land Use: Cropland

Erosion or Deposition: slightly eroded with sheet and rill erosion.

Parent Material: multiple deposits of ash and pumice.

Described by: T. Cook, G. Del Posso, and Soil Survey Staff.

Remarks: Date described: January 25, 1983. Field Moisture, 0-26 cm moist,

some clods dry, 26-220 moist.

Vegetation: before temperate forest

- Ap 0 26 cm Black (10YR 2/1, rubbed), loam; dark gray (10YR 4/1, dry); plowed, soft clods; very friable, non-sticky and non-plastic; weakly Omeary, very moist or wet; common very fine and fine roots; clear wavy boundary.
- Al 26 68 cm Black (10YR 2/1) loam; weak medium and coarse subangular blocky structure parting to moderate medium and coarse granular; friable, slightly sticky and slightly plastic; weakly smeary, very moist or wet; few very fine roots; many very fine interstitial few fine and medium tubular, pores; 2% 2 to 5 mm pumice fragments; a ffuse wavy boundary.

 30 to 50% worm casts in horizon.
- 68 120 cm Black (N 2/0) loam; weak very coarse subangular blocky structure parting to moderate medium and coarse subangular blocky; friable, slightly sticky and slightly plastic; a few thin patchy organic coatings on peds; few very fine roots; many very fine, few fine and medium tubular pores; coarse pumice fragments, 10% 2 to 40 mm; abrupt wavy boundary.

 30 to 50% worm casts in horizon. Fragments increase in amount in the lower part of the horizon.
- 2C 120 138 cm Very pale brown (10YR 7/4) and dark yellowish brown(10YR 4/4) very gravelly coarse sand; single grain; loose, non-sticky and non-plastic; few very fine roots; many very fine and fine interstital pores; coarse pumice fragments, 40% 2 to 40 mm; abrupt wavy boundary.
- 3Abl 138 183 cm Black (10YR 2/1) loam; weak very coarse subangular blocky structure parting to weak coarse subangular blocky; friable, slightly sticky and slightly plastic; few thin patchy dark reddish brown (5YR 3/3) organic coatings on peds; many very fine, few fine tubular pores; 1% 2 to 10 mm pumice fragments; diffuse wavy boundary. 30 to 50% worm casts in horizon.
- 3Ab2 183 220 cm Black (10YR 2/1) loam; weak very coarse subangular blocky structure parting to weak medium and coarse subangular blocky; friable, slightly sticky and slightly plastic; weakly smeary, very moist or wet; few thin patchy dark reddish brown (5YR 3/3) organic coatings on peds; many very fine and few fine tubular, pores. 30 to 50% worm casts in horizon.

ECUADOR 009

ECUADOR-SMSS

LAB CLASSIF: FINE-LOAMY, MIXED. ISOMESIC ANDIC HUMITROPEPT

S 83FN-325 -009

DATE 09/25/84

SAMPLE NO. 83P2091-2096 PEDON NO. 83P 457

PROJECT NO. 53P 83

U. S. DEPARTMENT OF AGRICULTURE SOIL CONSERVATION SERVICE NATIONAL SOIL SURVEY LABORATORY

PAGE 1 OF 4 PAGES

LINCOLN. NEBRASKA 68508

GENERAL METHODS 1B1A, 2A1, 2B

-8-- -9-- -10- -11- -12- -13- -14- -15- -16- -17- -18- -19- -20-(- - -TOTAL - - -)(- -CLAY- -)(- -SILT- -)(- - - - - -SAND- - - - -)(-COARSE FRACTIONS(MM)-)(>2MM CLAY SILT SAND FINE CO3 FINE COARSE VF F M VC - - - - WEIGHT - - - - WT SAMPLE HORIZON HZN DEPTH LT .002 .05 LT LT .002 .02 .05 .10 .25 .5 1 2 20 .1- PCI 0 NO. NO. (CM) -2 .0002 .002 -.02 -.05 -.10 -.25 -.50 .002 -.05 -5 -2 -20 -75 - - - - - PCT OF <2MM (3A1) - -- - - - - - - - - - - > <- PCT OF <75MM(3B1)-> SOIL **** SEE FOOTNOTE REGARDING PSDA BELOW ***** 832091 18 0- 26 AΡ 7.1 36.7 56.2 22.6 14.1 14.1 17.8 14.3 7.8 43 832092 28 26- 68 Α1 3.4 40.8 55.8 26.6 14.2 14.2 18.1 12.5 8.0 3.0 1 TR 5٧ 45 6 832093 35 68-120 A2 2.0 28.6 69.4 17.0 11.6 13.4 16.5 15.9 9.5 14.1 6 20V 71 33 832094 45 120-138 2C 0.1 3.3 96.6 1.8 1.5 0.7 7.1 31.9 35.0 21.9 9 50V 99 832095 5S 138-183 **3AB1** 4.1 46.5 49.4 30.7 15.8 15.4 17.0 9.8 5.8 1.4 36 832096 6S 183-220 **3AB2** 5.4 37.4 57.2 24.7 12.7 13.0 20.3 13.6 7.3 ORGN TOTAL EXTR TOTAL (- - DITH-CIT - -)(RATIO/CLAY)(ATTERBERG)(- BULK DENSITY -) COLE (- - -WATER CONTENT - -) WRD N S - LIMITS - FIELD 1/3 OVEN WHOLE FIELD 1/10 EXTRACTABLE 15 1/3 15 WHOLF SAMPLE HZN FE AL MN CEC BAR LL PΙ MOIST BAR DRY SOIL MOIST BAR BAR BAR SOIL NO. NO. 6A1C 6B3A 6R3A 6C2B 6G7A 6D2A 8D1 8D1 4F1 4F 4A3A 4A1D 4A1H 4D1 4B4 4B1C 4B1C <- - - - - - - PCT OF <2MM - -PCT <0.4MM <- - G/CC - - -> CM/CM <- - - PCT OF <2MM - -> CM/CM 832091 1 6.17 0.420 0.4 0.3 3.00 2.30 0.84 0.99 0.056 50.7 16.3 0.29 832092 4.32 0.253 0.6 0.4 4.68 3.68 0.98 1.07 0.029 46.2 12.5 0.32 832093 3.70 0.229 0.5 0.37.40 5.35 1.05 1.21 0.040 39.1 10.7 0.25 832094 0.23 0.3 TR 20.00 65.00 6.5 832095 2.02 J.6 0.2 3.22 1.98 1.20 1.26 0.016 34.3 0.31 8.1 832096 1.69 0.1 2.54 1.57 1.22 1.28 0.016 31.5 8.5 0.27

^{*} CAUTION SHOULD BE USED WITH PSDA DATA WHEN USED FOR MAKING ENGINEERING INTERPERTATIONS BECAUSE OF INCOMPLETE DISPERSION OF AMORPHOUS MATERIAL. THEY ARE PRESENTED TO AID IN CLASSIFICATION AND ASSESSMENT OF MINERALOGY. REFER TO PEDON DESCRIPTION FOR FIELD TEXTURE.

832092 2 5.1 1.7 0.2 0.4 7.4 20.4 0.2 27.6 15.9 7.6 3 27 47 10.6 5.3	PAGES RATORY -20-
CA MG NA K SUM ITY AL SUM NH4- BASES SAT SUM NH4 CACO3 OHMS MMHOS NAF CACL2 HOSAMPLE HZN 5B5A 5B5A 5B5A 5B5A BASES CATS OAC + AL OAC <2MM /CM /CM .01M NO. NO. 6N2E 602D 6P2B 6Q2B 6H5A 6G9A 5A3A 5A8B 5A3B 5G1 5C3 5C1 6E1G 8E1 81 8C1D 8C1F 8G	
NO. NO. 6N2E 602D 6P2B 6Q2B 6H5A 6G9A 5A3A 5A8B 5A3B 5G1 5G3 5C1 6E1G 8E1 81 8C1D 8C1F 8G 5C2	
832091 1 10.0 2.3 0.1 0.8 13.2 19.3 TR 32.5 21.3 41 62 10.0 5.5 832092 2 5.1 1.7 0.2 0.4 7.4 20.4 0.2 27.6 15.9 7.6 3 27 47 10.6 5.3	8C1F
832092 2 5.1 1.7 0.2 0.4 7.4 20.4 0.2 27.6 15.9 7.6 3 27 47 10.6 5.3	1:1
	5.6 5.9
	6.0
832095 5 8.2 1.2 0.3 0.4 10.1 10.0 20.1 13.2 50 77 9.6 5.9	6.2 6.4
	6.5
(SPODIC HORIZON CRITERIA)	
(SPODIC HORIZON CRITERIA)	·) - j
C FE AL FE+AL FE+AL AL+C OF (X-RAY)(DTA) TOTAL DO SAMPLE HZN (DIVIDED BY) ACCUM (<2U) RES WEA	. DOM WEATH
(SPODIC HORIZON CRITERIA)	7B1A
	,
832091	GA42
832093 3 0.2 0.3 0.6 0.3 1279 NX 6	GA35
832095 5 0.2 0.2 0.5 0.1 812	0/135
832096 6 0.1 0.1 0.2 TR 625 FD 1 NX 6	

FAMILY CONTROL SECTION: DEPTH 25-100 PCT CLAY 3 PCT .1-75MM 56. SPODIC HORIZON: INDEX OF ACCUMUL 2375 ANALYSES: S= ALL ON SIEVED <2MM BASIS V= 75-20MM FROM VOLUME ESTIMATES MINERALOGY: KIND OF MINERAL VR VERMICULITE NX AMORPHOUS GA GLASS AGG FD FELDSPAR

RELATIVE AMOUNT 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

NSSL SUPPLEMENTAL DATA SHEET

		EXT P	15 BAR	P RETENT	(ACID			EXTR	рН
		BRAYL	MOIST	KEIENI	AL	SI	FE	AL (KOH)	KCl
SAMPLE	HZN	6S3	4B2b	*	*	*	*		8Clq
NO.	NO.	PPM	<		-PCT OF <	ZMM-		>	
832091	1		16.2	54	0.5	0.2	0.4	0.4	
832092	2		16.9	67	0.7	0.4	0.5	0.7	
832093	3		14.7	69	0.8	0.4	0.5	0.7	
832094	4			30	0.1	0.1	0.2	0.1	
832095	5		14.8	36	0.5	0.3	0.4	0.3	
832096	6		14.7	47	0.4	0.3	0.5	0.3	

^{*} NEW ZEALAND PROCEDURE

Page 4 of 4

CP83FN128

-----(-----) SAMPLE HZN FA NO. NO. 603A 6C7A <-----> <----PCT------PCT-----> <----RELATIVE AMOUNTS----> <----PCT------> 83P2091 1 83P2092 2 VFS GA42 FP22 GS18 FG 8 HN 4 PR 3 HG 1 FE 1 OG 1 VR 2 NX 6 0.3 1.8 83P2093 3 NX 6 0.2 1.8 83P2094 4 VFS GA35 FP18 FG14 GS11 HG 7 HN 5 FE 4 BT 3 OG 2 FD 1 NX 6 0.2 0.9 83P2095 83P2096 6 FD 1 NX 6 0.3 2.6

ANALYSES: S=ALL ON SIEVED < 2mm BASIS

MINERALOGY: FA = FRACTION ANALYZED RE = RESISTANT

KIND OF MINERAL: FE = IRON OXIDES PP = PLAGIOCLASE FELDSPAR GA = GLASS AGGREGATES GS = GLASS HN = HORNBLENDE

FG = GLASS COATED FELDSPAR OG = GLASS COATED OPAQUE BT = BIOTITE FD = FELDSPAR

RELATIVE AMOUNT: 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

MINERALOGY BASED ON SAND/SILT:

MINERALOGY BASED ON CLAY:

FAMILY PLACEMENT:

COMMENTS:

00

Series: Equador No. 6 91 Date: 6/83

Pedon No: S83FN-325-006

Taxonomy: Fine-loamy, mixed, isomesic Andic Humitropept

Latitude: 0°20'15" Longitude: 78°33'45"

Location: In a pasture on the Santa Catalina INIAP Experiment Station

Physiography: Terrace

Geomorphic Position: On a hillslope Slope and Aspect: 12% E planar-planar

Elevation: 3140 m M.S.L.

Microrelief: smooth

Air Temp. Summer: 10.6°C Winter: 11.1°C Precipiation: 1450 mm Udic moisture regime

Water Table: > 10 meters

Drainage: Well drained Permeability: moderate

Stoniness: nonstony

Land Use: pasture in an experimentation station

Erosion or Deposition: none

Parent Material: ash and pumice deposits

Described by: T. Cook, G. Del Posso, and Soil Survey Staff

Remarks: . Date described: January 21, 1983. Field Moisture, 0-185 cm moist. Worm casts occupy 20 to 50 percent of the pedon. All coarse fragments are proice. Vegetation: Before clearing a native temperate forest, now mostly kikuyu grass

- Al 0 10 cm Black (10YR 2/1) loam; moderate medium coarse and very coarse granular structure; friable, non-sticky and non-plastic; many very fine, common medium and coarse roots; common very fine interstitial and few very fine pores; 1% 2 to 5 mm fragments; clear wavy boundary.
- A2 10 55 cm Black (N 2/0) loam; weak coarse subangular block structure parting to weak coarse and very coarse granular; friable, slightly sticky and slightly plastic; very weakly smeary, very moist or wet; common very fine and fine roots; many very fine and few fine tubular pores; 1% 2 to 5 mm fragments; clear wavy boundary.
- 55 79 cm Black (N 2/0) gravelly loam; weak coarse subangular blocky structre parting to weak coarse and very coarse granular; friable, slightly sticky and slightly plastic; very weakly smeary, very moist or wet; few thin discontinuous organic coatings on peds; few very fine and fine roots; many very fine tubular and few fine tubular pores; coarse fragments, 20% 2 to 60 mm; clear wavy boundary.
- 2Abl 79 117 cm Black (10YR 2/1) loam; weak coarse subangular blocky structure parting to weak coarse and very coarse granular; friable, slightly sticky and slightly plastic; very weakly smeary, very moist or wet; common thin continuous organic coatings on peds; few very fine and fine roots; many very fine tubular and few fine tubular pores; 1% 2 to 5 mm fragments; diffuse wavy boundary.
- 2Ab2 117 151 cm Black (10YR 2/1) silt loam; weak coarse subangular blocky structure parting to weak coarse and very coarse granular; friable, slightly sticky and slightly plastic; weakly smeary, very moist or wet; many thin continuous organic coatings on peds; few very fine and fine roots; many very fine and few fine tubular pores; 1% 2-5 mm fragments; diffuse wavy boundary.
- 2Ab3 151 185 cm Black (N 2/0) silt loam; weak coarse subangular blocky structure parting to weak coarse and very coarse granular; friable, slightly sticky and slightly plastic; weakly smeary, very moist or wet; many thin continuous organic coatings on peds; few very fine and fine roots; many very fine and few fine tubular pores; coarse fragments, 15%; most fragments in layer about 10-15 cm thick, average for whole horizon less than 5%.

PAGE 1 OF 4 PAGES

U. S. DEPARTMENT OF AGRICULTURE

NATIONAL SOIL SURVEY LABORATORY

SOIL CONSERVATION SERVICE

ECUADOR 006

LAB GLASSIF: FINE-LOAMY, MIXED, ISOMESIC ANDIC HUMITROPEPT

S 83FN-325 -006 DATE 09/25/84 SAMPLE NO. 83P2065-2070

ECUADOR-SMSS PROJECT NO 83P 83

CUADOR-SMSS PROJECT NO. 83P 83

GENERAL METHODS 1B1A, 2A1, 2B

-1-- -2-- -3-- -4-- -5-- -6-- -7-- -8-- -9-- -10- -11- -12- -13- -14- -15- -16- -17- -18- -19- -20-

													_				• •	.0	• • •	-7.0-
SAMPLE NO.	HZN NO.		HORIZON	LT .002	.002	.05)(CL FINE LT .0002	LT .002	.002	.02 05	.05	.10 25	.25 50	.5 -1	VC 1 -2	 2 -5	WE 5 -20	ACTIONS IGHT - 20 -75 <75MM(3	.1-	PCT O WHOLF
***** S 832065	EE FO	OOTNOTE R																		
832066 832067 832068 832069 832070	2S 3S 4S 5S	10- 55	A1 A2 A3 2AB1 2AB2 2AB3	3.8 2.4 3.2 2.6 2.6	50.7 50.7 44.6 54.2 55.2 50.5	44.6 45.5 53.0 42.6 42.2 46.9	0.1 0.1		33.2 34.0 29.4 35.8 35.1 29.9	17.5 16.7 15.2 18.4 20.1 20.6	15.7 13.6 15.5 19.5	16.1 16.4 16.4 16.1 13.7 13.0	8.3 8.6 10.0 6.7 4.8 6.0	3.4 3.6 5.9 2.9 2.9	1.3 1.2 7.1 1.4 1.3 2.1	TR TR 7 TR 1	TR 3 TR 1	3V 3V 40V 3V 3V 13V	31 32 70 29 26 37	3 3 50 3 4 15
SAMPLE NO.	HZN	ORGN TOTA C N 6A1C 6B:	3A	6R3A	FE 6C2B	AL 6G7A	MN 6D2A	CEC 8D1	/CLAY) 15 BAR 8D1	- LIR LL 4F1	RBERG) 11TS - Pl 4F 40.4MM	MOIST 4A3A	1/3 BAR 4A1D	OVEN DRY 4A1H	WHOLE SOIL 4D1	FIELD MOIST 4B4	1/10 BAR 4B1C	BAR	15 BAR 4824	WHOLE SOIL
832065 832066 832067 832068 832069 832070 5	2 3 4 5	6.22 0.42 5.68 0.36 5.38 4.21 3.91 4.13			0.8 0.8 0.8 0.9 1.1	0.6 0.6 0.5 0.5 0.4 0.3		4.55 5.55 9.08 6.56 9.46 10.50	3.30 4.18 6.42 4.34 6.00 6.46				0.98 0.87 1.06 1.01	1.07 0.96 1.15 1.13	0.051 0.029 0.025 0.027 0.038 0.054			56.7 47.0 54.6 45.5 50.9 57.2	15.9 15.4 13.9 15.6	0.30 0.26 0.33 0.35

^{*} PSDA IN THIS PEDON SHOULD NOT BE USED FOR MAKING ENGINEERING INTERPERTATIONS BECAUSE OF INCOMPLETE DISPERSION OF AMORPHOUS MATERIAL. THEY ARE PRESENTED TO AID IN CLASSIFICATION AND ASSESSMENT OF MINERALOGY. REFER TO PEDON DESCRIPTION FOR FIELD TEXTURE.

	ECU	JADOR 0			;	S 83FN-	325 -0	06	DA	TE 09/	/25/84		PEDON	NO. 8	3P 453	NAT	TIONAL	SOIL	SURVEY	OF 4 LABOR	RATORY
		-1	-2	-3	-4	-5	-6	-7	-8	-9	-10-	-11-	-12-	-13-	-14-	-15-	-16-	-17-	-18-	-19-	-20-
SAMPLE NO.		CA I 5B5A 6N2E	MG 5B5A 602D	NA 5B5A 6P2B	К 5В5А 6Q2В	SUM	1TY 6H5A	AL 6G9A	SUM CATS 5A3A	NH4- OAC 5A8B	BASES + AL 5A3B		5C3	NH4 GAC	CACO3 <2MM	OHMS		MMHOS /CM	NAF	-PH - CACL2 .01M 3C1F 1:2	8C1F
832065 832066 832067 12068 32069 632070	1 2 3 4 5 6	3.9 4.6 5.1 5.4 9.3 12.4	0.6 0.4 0.7 0.9 1.5 2.1	0.1 0.1 0.1	0.4 0.2 0.2 0.3 0.7	5.3 6.1 6.7 11.7	28.8 28.8 27.1 27.7 25.6 23.0	0.6 0.3 0.3 TR	33.8 34.1 33.2 34.4 37.3 38.7	21.1 21.8 21.0 24.6	6.2 5.9 6.4 7.0	19 10 5 4	15 16 18 19 31 41	23 25 28 32 48 58					10.9 11.0 11.0 10.9 10.2 9.8	4.7 4.9 5.1 5.4 5.6 5.2	5.1 5.6 5.9 6.0 6.3 6.2
SAMPLE NO.	HZN NO.	(NA C 6A4A	FE 6C8A	PHOSPHA AL 6G10	TE EXT FE+AL (D DI-C	CRITERI FRACTAB FE+AL IVIDED PCT CLAY	LE) AL+C BY) PCT	INDEX OF)					((7A21	- X-F	M - CLAY RAY 2U 7A21 AMOUNT	/)() 7A21	DT 7A3) A) 7A3	TOTAL RES 7B1A	DOM WEATH 7B1A
832065 832066 832067 832068 832069 832070	1 2 3 4 5 6		0.4 0.4 0.3 0.3 6.4 0.5	0.5 0.5 0.4 0.4 0.3 0.3	0.6 0.5 0.5 0.5 0.6	0.2 0.2 0.3 0.2 0.3		314 1449 768 1246 1224 1272						NX 6 NX 6							

FAMILY CONTROL SECTION: DEPTH 25-100 PCT CLAY 3 PCT .1-75MM SPODIC HORIZON: INDEX OF ACCUMUL 5878 43. ANALYSES: S= ALL ON SIEVED <2MM BASIS V= 75-20MM FROM VOLUME ESTIMATES MINERALDGY: KIND OF MINERAL NX AMORPHOUS

RELATIVE AMOUNT 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

		EXT P BRAYl	15 BAR MOIST	P RETENT	(ACID	OXAL SI	ATE) FE	EXTR AL (KOH)	рН КС1
SAMPLE NO.	HZN NO.	6S3 PPM	4B2b	*	* -PCT OF <	*	*		8Clg
					-PCT OF <	2MM-	 	>	
832065	1		17.4	80	0.9	0.4	0.6	1.0	
832066	2		25.0	83	1.0	0.5	0.7	1.1	
832067	3		25.5	86	1.2	0.6	0.6	1.3	
832068	4		26.5	86	1.2	0.7	0.6	1.4	
832069	5		29.6	76	1.0	0.7	0.9	0.9	
832070	6		31.7	70	0.8	0.6	1.0	0.6	

^{*} NEW ZEALAND PROCEDURE

CP83FN128

										 MINED	LOGY											 -
		(-	PTICAL ND/SI	-)	(X-F					DTA)	(TOT	ANAL)
SAMPLE NO.	HZN NO.	FA 7Bla	RE 7Bla <	7Bla	7BlA				7Bla	7Bla	7B1A	7A2I	. :	7A2I	7A21	. 7A2	2 I	7 A2I	7A3	7A3 PC1	K20 603A	6C7A
83P2065	1																					
83P2066 83P2067	_	VFS		GA 34	ОТ32	GC22	GS11					NX 6	i								0.3	2.0
83P2068 83P2069 83P2070	5	VFS .		GA37	OT27	GC21	GS15					NX 6									0.2	1.9

ANALYSES: S=ALI, ON SIEVED < 2mm BASIS

MINERALOGY: FA = FRACTION ANALYZED RE = RESISTANT

KIND OF MINERAL: GA = GLASS AGGREGATES GG = GLASS OT = OTHER NX = NON-CRYSTALLINE GC = GLASS COATED GRAIN

RELATIVE AMOUNT: 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

MINERALOGY BASED ON SAND/SILT:

MINERALOGY BASED ON CLAY:

PAMILY PLACEMENT:

COMMENTS:

y

6/83 Date: Series: Ecuador No. 4

Pedon No: S83FN-325-004

Taxonomy: Coarse-loamy over cindery, mixed, isomesi. Typic Humitropept Latitude: 0°28'00" Longitude: 78°42'40"

Location: Aurora Farm Physiography: Mountains Geomorphic Position: Hillslope

Slope and Aspect: 25% N convex-convex

Elevation: 2780 m M.S.L

Microrelief:

Air Temp. Summer: Winter:

Precipitation: 1700 mm approximately. Udic moisture regime.

Water Table: > 10 meters

Drainage: Well drained Permeability: Moderately slow

Stoniness: nonstony

Land Use: Pasture Erosion or Deposition: None to slight

Parent Material: ash and pumice deposits
Described by: T. Cook, G. Del Posso, and Soil Survey Staff.

Remarks: Date described: cm moist, 150-190 cm wet. January 26, 1983. Field Moisture, 0-45 cm wet, 45-150

Vegetation: Native vegetation was a temperate forest. Now cleared and planted to kikuyu pasture grass.

- 0 24 cm Al Very dark brown (10YR 2/2) loam; weak coarse subangular blocky structure parting to moderate medium, coarse, very coarse granular; friable, slightly sticky and slightly plastic; common very fine and few fine and medium roots; many very fine and few fine tubular pores; 1% 2-5 mm pumice fragments; gradual wavy boundary.
- A2 24 45 cm Very dark brown (10YR 2/2) sandy loam; weak medium and coarse subangular blocky structure; very friable, non-sticky and non-plastic; weakly smeary, very moist or wet; few very fine, fine, and medium roots; many very fine interstitial and tubular and common fine tubular pores; 5% 2-20 mm pumice fragments; clear wavy boundary.
- 63 cm Brown to dark brown (10YR 4/3) gravelly coarse sand; single grain; loose; few very fine, fine, and medium roots; many very fine and fine interstitial pores; 30% 2-20 mm pumice fragments; clear wavy boundary. This and next two horizons composed of many strata of pumice from about 100% 10-60 mm pumice to strata composed of pumice all less than 2 mm in size.
- 2Cl 63-83 cm Pale yellow $(2.5Y\ 7/4)$ gravelly co se sand; single grain; loose; few fine and medium roots; many very fine and fine inter-stitial pores; 30% 2-40 mm pumice fragments; clear wavy boundary.
- Light gray (10YR 7/2) very gravelly coarse sand; single 83 - 1.50 cm grain loose; very few very fine roots; many very fine and fine interstitial pores; 50% 2-60 mm pumice fragments; abrupt wavy boundary.
- 150 168 cm Very dark grayish brown (10YR 3/2) and dark brown (10YR 3/3) silt loam; common fine prominent dark reddish brown (5YR 3/3) and few fine prominent dark reddish brown (2.5YR 3/4) mottles; weak very coarse angular blocky structure; friable, slightly sticky and slightly plastic; weakly smeary, very moist or wet, very few very fine roots; common very fine tubular pores; gradual wavy boundary. Some brown (7.5YR 4/4) material in pores that looks like gel deposits (imogolite). The upper 5 cm of this horizon contains a few pumice fragments 5-10 cm in size.
- 168 160 cm Dark grayish brown (10YR 4/2) and brown to dark brown (10YR 4/3) sall loam; common fine distinct brown to dark brown (7.5YR 4/4) mottles; weak coarse angular blocky structure; friable, slightly sticky and slightly plastic; weakly smeary, very moist or wet, very few very fine roots; common very fine tubular pores; gradual wavy boundary.
- 190 207 cm Brown to dark brown (10YR 4/3) and dark brown (7.5YR 4/4) silty clay loam; massive; friable, slightly sticky and slightly plastic; weakly smeary, very moist or wet, very few very fine roots; common very fine and few fine tubular pores; 7% 2-30 mm pumice fragments.

ECUADOR 004

LAB CLASSIF: FINE-LOAMY OVER CINDERY, MIXED, ISOMESIC TYPIC HUMITROPEPT

S 83FN-325 -004

ECUADOR-SMSS

DATE 09/25/84

SAMPLE NO. 83P2119-2126 PEDON NO. 83P 461

PROJECT NO. 83P 83

NATIONAL SOIL SURVEY LABORATORY

LINCOLN. NEBRASKA 68508

SOIL CONSERVATION SERVICE

U. S. DEPARTMENT OF AGRICULTURE

PAGE 1 OF 4 PAGES

GENERAL METHODS 1B1A, 2A1, 2B

-1-- -2-- -3-- -4-- -5-- -6-- -7-- -8-- -9-- -10- -11- -12- -13- -14- -15- -16- -17- -18- -19- -20-(- - -TOTAL - - -)(- -CLAY- -)(- -SILT- -)(- - - - - - -SAND- - - - -)(-COARSE FRACTIONS(MM)-)(>2MM CLAY SILT SAND FINE CO3 FINE COARSE VF F M C VC ---- WEIGHT --- WT HORIZON LT .002 .05 LT LT .002 .02 .05 .10 .25 .5 1 2 5 20 .1- PCT 0 .002 -.05 -2 .0002 .002 -.02 -.05 -.10 -.25 -.50 -1 -2 -5 -20 -.75 .75 WHOLE <---- PCT OF <2MM (3A1) --- --- --- <-- PCT OF <75MM(3B1)-> SOIL SAMPLE HZN DEPTH NO. (CM) NO. **** SEE FOOTNOTE REGARDING PSDA BELOW **** 832119 18 0- 24 A1 2.1 39.1 58.8 24.3 14.8 16.5 20.1 13.5 6.7 2.0 2 3 V 46 6 832120 2S 24- 45 1.8 27.1 71.1 A2 17.0 10.1 13.5 20.7 18.5 11.8 6.6 6 3 11V 66 20 832121 3S 45- 63 2BW 0.6 3.0 96.4 2.4 0.6 3.9 15.5 26.0 29.5 21.5 10 40V 97 59 48 63- 83 832122 2C1 0.9 1.7 97.4 1.7 TR 1.1 8.6 31.2 36.9 19.6 11 39V 99 66 832123 5S 83-150 2C2 0.9 1.3 97.8 1.3 1.0 7.1 33.0 35.9 20.8 9 58V 99 82 832124 6S 150-168 3ABG 3.1 53.4 43.5 35.0 18.4 17.4 16.4 7.1 2.2 0.4 --__ 26 --832125 7S 168-190 3BW1B 3.6 51.4 45.0 34.3 17.1 14.8 17.6 8.1 1.4 1 3.1 1 2 32 832126 8S 190-207 3BW2B 13.0 36.7 50.3 25.7 11.0 13.2 15.3 11.1 3.9 6.8 15V 49 19 ORGN TOTAL EXTR TOTAL (- - DITH-CIT - -)(RATIO/CLAY)(ATTERBERG)(- BULK DENSITY -) COLE (- - -WATER CONTENT - -) WRD C N P S EXTRACTABLE 15 - LIMITS - FIELD 1/3 OVEN WHOLE FIELD 1/10 1/3 15 WHOLE SAMPLE HZN FE AL MN BAR LL PI MOIST BAR DRY SOIL MOIST CEC BAR BAR BAR SOIL NO. 6A1C 6B3A 6C2B 6G7A 6D2A 8D1 8D1 4F1 4F 4A3A 4A1D 4A1H 4D1 4B4 4B1C 4B1C 4B2A 4C1 NO. PCT <0.4MM <- - G/CC - - -> CM/CM <- - - PCT OF <2MM - -> CM/CM 832119 1 4.26 0.358 1.0 0.7 7.24 5.95 1.02 1.08 0.019 41.4 12.5 0.29 832120 2 2.51 0.190 0.8 0.5 5.28 4.83 1.02 1.07 0.015 37.5 8.7 0.27 832121 3 0.29 0.020 0.3 0.1 2.67 5.83 3.5 832122 4 0.17 0.3 0.1 2.11 4.00 3.6 832123 5 0.14 0.3 TR 1.56 4.44 4.0 6 832124 1.09 0.8 0.2 2.68 2.84 1.09 1.16 0.021 43.6 8.8 0.38 832125 7 1.07 2.50 2.69 0.9 0.2 1.09 1.17 0.024 43.5 9.7 0.37 832126 8 0.49 1.9 0.2 0.90 1.33 1.09 1.17 0.022 42.6 17.3 0.25

^{*} CAUTION SHOULD BE USED WITH PSDA DATA WHE! USED FOR MAKING ENGINEERING INTERPERTATIONS BECAUSE OF INCOMPLETE DISPERSION OF AMORPHOUS MATERIAL. THEY ARE PRESENTED TO AID IN CLASSIFICATION AND ASSESSMENT OF MINERALOGY. REFER TO PEDON DESCRIPTION FOR FIELD TEXTURE.

SPODIC HORIZON: INDEX OF ACCUMUL 764

	ECUAD	OR 004	4	_3	-11	83FN-	325 - 0	04	DA	TE_09/	25/84		PEDON	NO. 8	3P 461	NA'	TIONAL	COLL	CHRUCK	OF 4 LABOR	ATORIC
					-4		-0	-,	-6	-9	-10-	-11-	-12-	-13-	-14-	-15-	-16-	-17-	-18-	-19-	-20-
SAMPLE NO.	HZN 51 NO. 61	CA B5A 5 N2E 6	MG 5B5A 502D	NA 5B5A 6P2B	5B5A 6Q2B	SUM BASES	1ТҮ э ч5А	AL 6G9A	SUM CATS 5A3A	NH4- OAC 5A8B	BASES + AL 5A3B	SAT 5G1	SUM	NH4 OAC	CO3 AS CACO3 <2MM 6E1G	OHMS /CM		MMHOS /CM	NAF	-PH - CACL2 .01M 8C1F 1:2	H20
832119 832120 832121 832122 832123 832124 832125 832126	2 3 4 5 6 7	0.4 0.6 0.5 2.9	0.5 0.3 0.1 0.1 0.6 0.9	0.1 0.1 0.1 0.1 0.1 0.2 0.3	0.1 TR TR TR 0.4 0.4	4.0 2.2 0.6 0.8 0.7 4.1 5.1 6.5	27.3 18.5 3.0 1.7 1.3 10.6 9.8 10.2	0.2 0.3 0.1 TR 0.1 0.1 TR	20.7 3.6 2.5 2.0 14.7	15.2 9.5 1.6 1.9 1.4 8.3 9.0 11.7	4.2 2.5 0.7 5.8 4.2	5 12 14 12 2	13 11 17 32 35 28 34 39	26 23 38 42 50 49 57 56					11.3 11.2 10.4 10.2 9.7 10.1 10.0 9.6	5.4 5.6 5.6 5.3 5.7 5.7	6.0 5.9 6.1 6.1 6.2 6.2
SAMPLE NO.	(- HZN NO. 6A	-NA F C \4A 6	PYROP. FE 5C8A	HOSPHA AL 6G10	TE EXT FE+AL (Di DI-CI	RITERIA RACTABI FE+AL VIDED I PCT CLAY	LE) AL+C BY) PCT	INDEX OF)				(((7A21	X-F - X-F - X21 LATIVE	- CLAY RAY S2U 7A21	(:)(:) 7A21	DT) A) 7A3	TOTAL RES 1	DOM WEATH 781A
832119 832120 832121 832122	1 2 3 4		0.4	0.6 0.5 0.1 0.1	0.6 0.6 0.3 0.3	0.5 0.4 0.2 0.1		726 416						NX 6							
832123 832124 832125 832126	5 6 7 8		0.2 0.2 0.1	0.1 0.2 0.2 0.1	0.3 0.4 0.4 0.1	0.1 0.1 0.1 TR		237 288 173						NX 6 NX 6 KH 1	NX 6						

RELATIVE AMOUNT 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

V= 75-20MM FROM VOLUME ESTIMATES

KH HALLOYSITE

FAMILY CONTROL SECTION: DEPTH 25-100 PCT CLAY 1 PCT .1-75MM 90.

ANALYSES: S= ALL ON SIEVED <2MM BASIS V= 75-20MM FROM VOLUME ESTIMATES

MINERALOGY: KIND OF MINERAL NX AMORPHOUS

Page 3 of 4

ECUADOR 004 S83FN-325-004

NSSL SUPPLEMENTAL DATA SHEET

		EXT	15	P	(ACI	D OXAL	ATE)	EXTR	рН
		P	BAR	RETENT	AL	SI	FE	AL	KC1
		BRAYl	MOIST					(KOH)	NOI
SAMPLE	HZN	6 S 3	4B2b	*	*	*	*	(1(011)	8Clg
NO.	NO.	PPM	<		-PCT OF	<2MM		>	ocig
									_
832119	1		23.8	86	1.2	0.5	0.7		
832120	2		16.0	75	1.1	0.5	0.6		
832121	3		4.5	34	0.3	0.2	0.2		
832122	4		4.4	27	0.2	0.2	0.3		
832123	5		4.5	25	0.1	0.2	0.3		
832124	6		17.5	56	0.5	0.4	0.5		
832125	7		19.1	5 2	0.4	0.4	0.6		
832126	8		26.1	5 4	0.3	0.3			
				J4	0.3	0.3	0.7		

^{*} NEW ZEALAND PROCEDURE

CP83FN128

CPOSENI	20																	
		(0	PTICAL)	(X - RA Y)	-		(TOT	
SAMPLE NO.	HZN NO.	(FA 7BlA	RE 7BlA <	7BlA	7BlA	7B1 A	7B1A	7Bla	7Bla	7BlA	7BlA	7BlÁ	7A2I 7A2I 7A2I 7A2I <relative amounts<="" th=""><th>7A2I</th><th>7A3</th><th>7A3</th><th>K20 6Q3A</th><th>FE 6C7A</th></relative>	7A2I	7A3	7A3	K20 6Q3A	FE 6C7A
83P2119 83P2120 83P2121	2	VFS		GA 32	от27	GS24	GC17						NX 6				0.0	1.9
83P2122 83P2123 83P2124 83P2125 83P2126	5 6 7												NX 6 NX 6 KH 1 NX 6				0.3 0.2 0.3	0.5 2.4 3.0

ANALYSES: S=ALL ON SIEVED < 2mm BASIS

MINERALOGY: FA = FRACTION ANALYZED RE = RESISTANT

KIND OF MINERAL: GA = GLASS AGGREGATES GS = GLASS OT = OTHER NX = NON-CRYSTALLINE GC = GLASS COATED GRAIN

KH = HALLOYSITE

RELATIVE AMOUNT: 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

MINERALOGY BASED ON SAND/SILT:

MINERALOGY BASED ON CLAY:

PAMILY PLACEMENT:

COMMENTS:

Date: 6/83

Series: Ecuador No. 5

Pedon No: S83FN-325-005

Taxonomy: Medial, isohyperthermic Entic Dystrandept

Latitude: Longitude:

Location: In an Oil Palm Grove at INIAP Experiment Station at La Concordia.

Physiography: Terrace

Geomorphic Position: stable summit

Slope and Aspect: 1% planar-planar Elevation: 280 m M.S.L

Microrelief: smooth

Summer: 22.9°C Winter: 24°C Air Temp. Precipitation: 3200 mm Udic moisture regime.

Water Table: 2 meters

Drainage: Somewhat poorly drained Stoniness: nonstony Permeability: Moderately rapid

Land Use: Oil palm plantation and pasture for grazing

Erosion or Deposition: none

Parent Material: several ash deposits
Described by: T. Cook, G. Del Posso, and Soil Survey Staff.

Remarks: Date described: January 27, 1983. Field Moisture, 0-110 cm wet, 110-190 cm free water. Medium and coarse decayed roots and pores to 142 cm, mostly above 110 cm.

Vegetation: before native tropical humid forest.

- 0 -7 cm Dark brown (10YR 3/3) and black (10YR 2/1) loam; weak coarse subangular blocky structure; friable, non-sticky and non-plastic; common very fine and few fine roots; common very fine to fine tubular pores; clear irregular boundary. About 1% fragments of brick and artifacts.
- Brown to dark brown (10YR 4/3) silt loam; weak medium 7 - 27 cm subangular blocky structure; very friable, slightly sticky and slightly plastic; weakly smeary, very moist or wet; few very fine and fine roots; many very fine and fine tubular very fine interstitial pores; gradual wavy boundary. About 18 fragments of brick and artifacts.
- Bw2 27 54 cm Brown to dark brown ($10\,\mathrm{YR}$ 4/3) and dark yellowish brown ($10\,\mathrm{YR}$ 4/4) silt loam; weak coarse angular blocky structure parting to weak medium subangular blocky; friable, non-sticky and non-plastic; weakly smeary, very moist or wet; few very fine and fine roots; common very fine and fine tubular pores; gradual wavy boundary. About 1% fragments of brick and artifacts.
- 54 79 cm Very dark grayish brown (10YR 3/2) and dark yellowish brown (10YR 4/4) loam; weak coarse angular blocky structure parting to weak medium subangular blocky; friable, non-sticky and non-plastic; weakly smeary, very moist or wet; few very fine roots; common very fine and fine tubular and many very fine interstitial pores; gradual wavy boundary.
- 79 100 cm Brown to dark brown (10YR 4/3) and very dark grayish brown (10YR 3/2) loam; weak coarse angular blocky structure parting to weak medium subangular blocky; friable, slightly sticky and slightly plastic; weakly smeary, very moist or wet; few very fine roots; common very fine and fine tubular and many very fine interstitial pores; clear wavy boundary.
- 100 110 cm Variegated gray (5Y 5/1) and olive gray (5Y 5/2) fine sandy loam; common fine and medium distinct and priminent dark yellowish brown and dark brown (10YR 4/4, 5/4, 4/3) and dark grayish brown (2.5Y 4/2) mottles; structureless; friable, non-sticky and non-plastic; moderately smeary, very moist or wet; few very fine roots; many very fine interstitial and common very fine tubular pores; abrupt irregular boundary.
- 110 142 cm Very dark grayish brown (10YR 3/2) and dark yellowish brown (10YR 3/4) silt loam; structureless; friable, slightly sticky and slightly plastic; strongly smeary, very moist or wet; few very fine roots; many very fine interstitial and tubular pores; clear wavy boundary. Very thixotropic free water can be squeezed by hand.
- 3Bwb 142-190 cm Dar', grayish brown (10YR 4/2) silt loam; many fine distinct and prominent dark yellowish brown (10YR 4/4) and brown to dark brown (7.5YR 4/4) mottles; structureless; very friable, slightly sticky and slightly plastic; strongly smeary, very moist or wet; few very fine roots; many very fine interestitial and bubular pares. Very thixotropic free water can be squeezed by hand.

ECUADOR 005

LAB CLASSIF: MEDIAL, ISOHYPERTHERMIC ENTIC DYSTRANDEPT

S 83FN-325 -005

ECUADOR-CASS

DATE 09/25/84

SAMPLE NO. 83P2151-2158 PEDON NO. 83P 465

PROJECT NO. 83P 83

U. S. DEPARTMENT OF AGRICULTURE SOIL CONSERVATION SERVICE NATIONAL SOIL SURVEY LABORATORY LINCOLN, NEBRASKA 68508

PAGE 1 OF 4 PAGES

GENERAL METHODS 181A, 2A1, 2B

-1-- -2-- -3-- -4-- -5-- -6-- -7-- -8-- -9-- -10- -11- -12- -13- -14- -15- -16- -17- -18- -19- -20-(- - -TOTAL - - -)(- -CLAY- -)(- -SILT- -)(- - - - - -SAND- - - - - -){-COARSE FRACTIONS(MM)-)(>2MM CLAY SILT SAND FINE CO3 FINE COARSE VF F M C VC ---- WEIGHT --- WI HORIZON LT .002 .05 LT LT .002 .02 .05 .10 .25 .5 1 2 5 20 .1- PCI C .002 -.05 -2 .0002 .002 -.05 -.10 -.25 -.50 -1 -2 -5 -20 -.75 .75 WHOLE SAMPLE HZN DEPTH 20 1- PC1 C NO. NO. (CM) **** SEE FOCTNOTE REGARDING PSDA BELOW ***** 832151 18 0- 7 A 4.0 56.6 39.4 38.4 18.2 20.3 18.2 0.7 0.2 -- TR 3 7- 27 832152 25 BW1 2.0 53.0 45.0 33.6 19.4 23.5 20.7 0.8 --V 21 --832153 3S 27- 54 BW2 1.7 42.5 55.8 25.6 16.9 26.0 25.6 3.9 0.3 -- TR 3 V 32 3 832154 48 54- 79 2AB 1.2 35.6 63.2 17.9 17.7 25.6 33.5 4.1 38 --832155 1.1 41.7 57.2 5S 79-100 2BWB 23.4 18.3 23.8 29.3 4.0 0.1 -- TR 33 --832156 6S 100-110 2CG 1.9 44.3 53.8 25.1 19.2 22.6 28.6 2.6 31 --832157 75 110-142 3AB 2.2 35.5 62.3 19.4 16.1 24.9 33.6 3.6 37 --832158 85 142-190 3BWB -- 16.2 83.8 7.5 8.7 21.3 43.8 14.8 3.6 0.3 TR 62 --ORCH TOTAL EXTR TOTAL (- - DITH-CIT - -) (RATIO/CLAY) (ATTERBERG) (- BULK DENSITY -) COLE (- - -WATER CONTENT - -) WRD C N P S EXTRACTABLE 15 - LIMITS - FIELD 1/3 OVEN WHOLE FIELD 1/10 1/3 15 WHOLE SAMPLE HZN FE AL MN CEC MOIST BAR DRY SOIL MOIST BAR BAR LL PI BAR EAR SOLL NO. 6A1C 6B3A 6C2B 6G7A 6D2A 8D1 8D1 4F1 4F 4A3A 4A1D 4A1H 4D1 4B4 4B1C 4B1C 4B2 4C1 NO. <- - - - - - - - PCT OF <2MM - - - - - -> PCT <0.4MM <- - G/CC - - -> CM/CM <- - -PCT OF <2MM - -> CM/CM 832151 1 5.71 0.466 0.7 0.4 5.90 5.28 0.80 0.94 0.055 61.7 21.1 0.32 2.68 0.239 832152 0.7 0.5 6.00 6.50 0.73 0.84 0.048 55.7 13.0 0.31 832153 3 2.23 0.200 0.9 0.5 7.47 8.24 0.84 0.95 0.041 58.8 14.0 0.37 832154 4 1.78 0.9 0.5 8.25 10.17 0.85 0.95 0.038 53.1 12.2 0.35 832155 5 1.00 6.0 0.4 6.82 10.00 0.88 0.97 0.033 49.9 11.0 0.34 832156 6 0.57 0.7 0.3 3.53 5.42 1.05 1.09 0.013 32.6 10.3 0.23 832157 7 2.12 0.8 1.9 9.45 9.50 0.77 0.89 0.049 58.5 20.9 0.29 832158 8 1.45 2.5 0.9 0.60 0.85 0.123 77.9 19.4 0.35

^{*} PSDA IN ANDEPTS SHOULD NOT BE USED FOR MAKING ENGINEERING INTERPERTATIONS BECAUSE OF INCOMPLETE DISPERSION OF AMORPHOUS MATERIAL. THEY ARE PRESENTED TO AID IN CLASSIFICATION AND ASSESSMENT OF MINERALOGY. REFER TO PEDON DESCRIPTION FOR FIELD TEXTURE.

	ECU		05 -2	-3											83P 465 -14-			SOIL	SURVEY		RATORY
SAMPLE NO.	HZN	CA 535A 6N2E	MG 5B5A	NA 5B5A 6P2B	K 5B5A 6Q2B	SUM	1 T Y 6 H 5 A	Al. 6G9A	SUM CATS 5A3A	Nri4- OAC 5A8B	BASES + AL 5A3B	SAT 5G1		44N 0AC	<2MM 6E1G	OHMS /CM		MMHOS /CM	NAF	-PH - CACL2 .01M 8C1F 1:2	H20 [′]
832151 832152 832153 832154 832155 832156 832157 832158	1 2 3 4 5 6 7 8	15.5 3.8 1.3 0.5 0.3 0.5 1.7 3.0	1.9 1.0 0.4 0.2 0.1 0.1 0.3	0.2 0.2 0.2 0.2 0.2 0.1 0.1	0.2 0.1 0.1 TR TR 3.1 1.8 2.5	17.8 5.1 2.0 0.9 0.6 0.8 3.9 6.3	43.4 21.6 26.1 25.3 17.7 17.7 33.0 32.6	TR 0.2 0.1 Tk 0.1 0.1	18.3 18.5	23.6 12.0 12.7 9.9 7.5 6.7 20.8 22.8	2.2 1.0 0.9 4.0	9 10 11 2	29 19 7 4 3 4 11	75 42 16 9 8 12 19 28					10.4 10.9 11.1 11.1 11.0 10.8 11.0	5.7 5.6 5.4 5.3 5.8 5.5 5.2	6.0 6.1 5.8 5.6 5.6 5.8 6.1 6.0
SAMPLE NO.	HZN	(NA C 6A4A	FE 6C8A	PHOSPHA AL 6G10	TE EXT FE+AL (DI Di-CI	RITERI RACTAB FE+AL VIDED PCT CLAY	LE) AL+C BY) PCT	INDEX OF)					(· (· (· 7A21	7A21	- CLA RAY - <2U - 7A21	Y)) 7A21	(DT) A) 7A3	TOTAL RES 781A	DOM WEATH 7B1A
832151 832152 832153 832154 832155 832156 832157 832158	1 2 3 4 5 6 7 8		0.3 0.2 0.1 0.1 0.1 0.1	0.4 0.5 0.4 0.3 0.2 0.4	0.6 0.6 0.4 0.3 0.3 0.3	0.2 0.4 0.3 0.3 0.3 0.2		414 514 736 602 373 175 1146						NX 6 NX 6	NX 6						

FAMILY CONTROL SECTION: DEPTH 25-100 PCT CLAY 1 PCT .1-75MM 34. SPODIC HORIZON: INDEX OF ACCUMUL 373 ANALYSES: S= ALL ON SIEVED <2MM BASIS V= 75-20MM FROM VOLUME ESTIMATES MINERALOGY: KIND OF MINERAL VC VERM-CHLOR NX AMORPHOUS MI MICA

RELATIVE AMOUNT 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

Page 3 of 4

NSSL SUPPLEMENTAL DATA SHEET

		EXT P BRAYl	15 BAR MOIST	P RETENT	(ACI	D OXAL	ATE) FE	EXTR AL (KOH)	pH KCl
SAMPLE NO.	HZN NO.	6S3 PPM	4B2b <		+ -PCT OF	* <2MM	*	>	8Clg
832151	1		28.3	68	0.7	0.3	0.4	0.8	
832152	2		24.7	61	1.0	0.5	0.4	1.3	
832153	3		32.1	90	1.4	0.8	0.4	2.2	
832154	4		28.5	94	1.9	1.2	0.4	2.2	
832155	5		24.3	89	1.6	1.2	0.4	1.8	
832156	6		18.6	77	1.2	0.8	0.3	1.4	
332157	7		38.7	97	2.7	1.7	1.0	2.2	
832158	8		58.9	97	2.1	1.5	0.8	2.0	

^{*} NEW ZEALAND PROCEDURE

ECUADOR 005 S83FN-325-005

_	_
_	_
ς	_
	_

Page 4 of 4

		(0	PTICAL)	(X-RA	Y)	(ī	TA)	(TOT	ANAL)
	HZN NO.	(FA 7BlA	RE 7BlA <	7BlA	7BlA		7BlA	7B1A	7B1A	7BlA	7BlA	7BlA	7A2I	7A2I RELATI	7A2I	7A2I	7A2I	7A3	7A3	K2O 6Q3A	
33P2151 33P2152	1 2	VFS		C 3 5 0	CC17	OT14	CS10						vc 1	NX 6						0.1	1.2
	3	413		GAJJ	GCI7	0114	GDIU						•••	1171 0						· • •	
33P2154	4	VFS		GA65	OT16	GC14	GS 5						NX 6							0.0	1.1
33P2155	5	VFS		GA58	OT24	GC15	GS 3						NX 6							0.0	1.3
33P2156	6																				
83P2157	7	VFS		GA78	OT14	GC 6	GS 2						MI 1	NX 6						0.4	2.2
83P2158	8																			4	

ANALYSES: S=ALL ON SIEVED < 2mm BASIS

MINERALOGY: FA = FRACTION ANALYZED RE = RESISTANT

KIND OF MINERAL: GA = GLASS AGGREGATES GS = GLASS VC = VERMICULITE-CHLORITE OT = OTHER NX = NON-CRYSTALLINE

GC = GLASS COATED GRAIN MI = MICA

RELATIVE AMOUNT: 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

MINERALOGY BASED ON SAND/SILT:

MINERALOGY BASED ON CLAY:

FAMILY PLACEMENT:

COMMENTS:

Ecuador No. 12 Series:

Pedon No: S83FN-325-012

Taxonomy: Fine-loamy, mixed, isohyperthermic Typic Argiudoll

Latitude: 0°50'00" Longitude: 79°28'50"

Location: This pedon is located about 90 km south of Santo Domingo near Buena

Date: 8/82

Fe (1 km before this town). Physiography: Fan Terrace. Geomorphic Position: Summit.

Slope and Aspect: 1% pl0nar-planar Elevation: 120 m M.S.L

Microrelief: smooth.

Summer: 23.4°C Air Temp. Winter: 24.9°C Precipitation: 2200 mm Udic moisture regime.

Water Table: > 10 meters

Drainage: Well drained Permeability: Moderately slow

Stoniness: nonstony

Land Use: Forest, ungrazed Erosion or Deposition: none

Parent Material: multiple recent ash deposits over a very old ash deposit with a strongly developed argillic horizon.

Described by: T. Cook, G. Del Posso, and Soil Survey Staff.

Remarks: Date Described: January 28, 1983. Field Moisture, 0-170 cm wet. To a depth of 130 cm there are a few (1%) artifact fragments of broken, brick and stone, tcols.

Vegetacion: Tropical forest in closely with cocoa trees.

- 0 10 cm Very dark brown (10YR 2/2) loam; moderate coarse subangular blocky structure parting to strong medium and coarse granular; friable, slightly sticky and slightly plastic; few very fine, fine, medium and coase roots; many very fine, fine and medium tubular and many very fine and fine interstital pores; clear wavy boundary.
- 10 52 cm Dark brown (7.5YR 3/2) loam; moderate medium and coarse subangular blocky structure parting to weak coarse granular; friable, slightly sticky and slightly plastic; weakly smeary, very moist or wet; few very fine, medium, and fine, coarse roots; many very fine, fine, and medium tubular pores; clear wavy boundary.
- 2Ab 52 83 cm Very dark brown (10YR 2/2) loam; weak coarse angular blocky structure; very friable, slightly sticky and slightly plastic; moderately smeary, very moist or wet; few very fine, medium, and fine coarse roots; many very fine, fine, and medium tubular; gradual wavy boundary.
- 3Btbl 83 102 cm Very dark grayish brown (10YR 3/2) and dark grayish brown (10YR 4/2) silt loam; moderate medium and coarse subangular blocky structure; very friable, sticky and slightly plastic; strongly smeary, very moist or wet; few very fine, fine, and medium, and coarse roots; many very fine, fine, and medium tubular pores; few thin patchy stress oriented coats on peds; gradual wavy boundary.
- 3Btb2 102 - 130 CM Dark brown (7.5YR 3/2) clay loam; weak coarse subangular blocky structure parting to weak coarse granular; friable, sticky and plastic; moderately smeary, very moist or wet; few fine, medium, and coarse roots; many very fine, fine, and medium tubular pores; many thin continuous clay films on peds and in pores; few pressure faces; diffuse wavy boundary.
- 130 150 cm Dark reddish brown (5YR 3/2) clay; moderate medium and coarse prismatic structure parting to moderate medium and coarse angular blocky; firm, sticky and plastic; weakly smeary, very moist or wet; many very fine, fine, and medium tubular pores; many thin continuous stress oriented coats on faces and many thin clay films on peds and in pores; gradual wavy boundary.
- 156 170 cm 4Btb4 Dark reddish brown (5YR 3/4 exterior ped faces) and dark brown (7.5YR 3/4 interior) clay; moderate medium prismatic structure parting to moderate medium and coarse angular blocky; firm, sticky and very plastic; weakly smeary, very moist or wet; common very fine, fine, and medium tubular pores; many thin continuous stress oriented coats on faces of peds and many thin clay fims in pores and on peds, few moderately thick clay films in pores.

ECUADOR 012

ECUADOR-SMSS

LAB CLASSIF: FINE-LOAMY, MIXED, ISOHYPERTHERMIC TYPIC ARGIUDOLL

1B1A, 2A1, 2B

S 83FN-325 -012

GENERAL METHODS

DATE 09/25/84

SAMPLE NO. 83P2167-2173 PEDON NO. 83P 467

PROJECT NO. 83P 83

U. S. DEPARTMENT OF AGRICULTURE SOIL CONSERVATION SERVICE NATIONAL SOIL SURVEY LABORATORY LINCOLN, NEBRASKA 68508

PAGE 1 OF 4 PAGES

-9-- -10- -11- -12- -13- -14- -15- -16- -17- -18- -19- -20-

									. •	• •			- 14-	-15-	-10-	1 / -	-18-	-19-	-20-
SAMPLE NO.	HZN NO.	DEPTH (CM)	HORIZON	LT .	IAL ILT SAN 002 .05 .05 -2	LT	LT .002	.002	.02	.05	. 10 25	M	.5 -1	VC 1	2	WE 5 -20	ACTIONS IGHT - 20 -75 <75MM(3	 .1- 75	PCT 0 WHOLE
**** S	EE FO	OOTNOTE R	REGARDING	PSDA BELO	w ****													, ,	
832167 832168 832169 832170 832171 832172 832173	15 25 35 45 55 65	0- 10 10- 52 52- 83 83-102 102-130 130-150 150-170	A1 BW 2AB 3BTB; 3BTB2 4BTB3 4BTB4	7.0 53 3.3 40 3.4 48 13.0 51 27.1 40 52.3 27	3.5 39.1 3.1 56.0 3.0 48.0 1.8 35.1 3.2 32.1 7.7 20.0 1.5 13.1	5 5 5 2 7		30.6 23.0 29.1 33.5 25.8 17.9 14.3	22.9 .17.1 18.9 18.3 14.4 9.8 7.2	21.0 23.0 23.0 19.3 14.3 9.1 5.4	15.5 30.0 23.9 15.0 17.7 10.1 6.3	2.3 3.3 1.5 0.7 0.7 0.7	0.7 0.3 0.2 0.1 0.1 0.3	0.1	 TR TR TR TR		V V 3V 3V 2V	18 34 28 18 20 11	3 3 2
																		-	
SAMPLE NO.	HZN NO.	ORGN TOT C N 6A1C 6B		6R3A 60	E AL	MN 6D2A	CEC	7/CLAY) 15 BAR' 8D1	LL 4F1	BERG) ITS - PI 4F 0.4MM	MOIST 4A3A	BAR	DRY 4A1H	SOIL 4D1	MOIST 4B4	1/10 BAR 4B1C	1/3 BAR	15 BAR 4B2A	WHOLE SOIL 4C1
832167 832168 832169 832170 832171 832172 832173	2 3 4 5 6	4.33 0.4 1.65 0.1 1.65 0.1 0.58 0.24 0.15 0.19	70	1 1 2	.5 0.4 .8 0.2 .9 0.2 .7 0.3		3.91 5.88 5.35 0.92 0.34 0.25 0.24	2.89 5.52 5.09 1.18 0.59 0.51 0.52				0.91 0.78 0.82 0.88 1.40 1.21	0.83 0.87 0.94 1.51 1.49	0.039 0.021 0.020 0.022 0.025 0.072 0.084			47.1 50.4 52.4 52.4 26.8 38.8 44.2	20.2 18.2 17.3 15.3 16.0 26.6 34.2	0.24 0.25 0.29 0.32 0.15 0.15

^{*} CAUTION SHOULD BE USED WITH PSDA DATA WHEN USED FOR MAKING ENGINEERING INTERPERTATIONS BECAUSE OF INCOMPLETE DISPERSION OF AMORPHOUS MATERIAL. THEY ARE PRESENTED TO AID IN CLASSIFICATION AND ASSESSMENT OF MINERALOGY. REFER TO PEDON DESCRIPTION FOR FIELD TEXTURE.

```
PAGE 2 OF 4 PAGES
       ECUADOR 012 S 83FN-325 -012 DATE 09/25/84 PEDON NO. 83P 467 NATIONAL SOIL SURVEY LABORATORY
-1-- -2-- -3-- -4-- -5-- -6-- -7-- -8-- -9-- -10- -11- -12- -13- -14- -15- -16- -17- -18- -19- -20-
          (- NH4OAC EXTRACTABLE BASES -) ACID- EXTR (- - - -CEC - - -) AL -BASE SAT- CO3 AS RES. COND.(- - - -PH - CA MG NA K SUM ITY AL SUM NH4- BASES SAT SUM NH4 CACO3 OHMS MMHOS NAF CACL2 N 5B5A 5B5A 5B5A 5B5A BASES CATS OAC + AL OAC <2MM /CM .01M .6N2E 602D 6P2B 6Q2B 6H5A 6G9A 5A3A 5A8B 5A3B 5G1 5C3 5C1 6E1G 8E1 31 8C1D 8C1T
                                                                                                         COND.(- - - - PH - - -)
                                                                                                         MMHOS NAF CACL2 H20
SAMPLE HZN 5B5A 5B5A 5B5A BASES
        NO. 6N2E 602D 6P2B 6Q2B 6H5A 6G9A 5A3A 5A8B 5A3B 5G1
                                                                                                         91 8C1D 8C1F 8C1F
            1:2 1:1
                       0.1
832167
        1 18.2
                  3.9
                              0.9 23.1 16.3
                                                -- 39.4 27.4
                                                                                                               10.2
                                                                                                                     6.1
                                                                                                                           6.5
832168
            8.9
                                                -- 29.9 19.4
                  0.9
                        0.1
                             1.1 11.0 18.9
                                                                             37
                                                                                   57
                                                                                                               10.6
                                                                                                                     5.8
                                                                                                                           6.6
832169
        3
            7.7
                  0.4
                        0.2
                              0.9
                                   9.2
                                        22.0
                                                -- 31.2 18.2
                                                                                  51
                                                                                                               10.6
                                                                                                                     6.1
                                                                                                                           6.4
        4
                                                -- 18.0 11.9
832170
            5.8
                  0.6
                        0.2
                             1.6
                                   8.2
                                         9.8
                                                                                  69
                                                                                                                9.8
                                                                                                                     5.8
                                                                                                                           6.7
832171
                              0.5
                                                -- 12.4 9.2
            4.6
                  1.1
                        0.3
                                   6.5
                                         5.9
                                                                                  71
                                                                                                                9.5
                                                                                                                    6.3
                                                                                                                           6.8
                                                -- 15.9 12.9
832172
            5.8
                                   8.6
                 1.7
                        0.4
                              0.7
                                        7.3
                                                                                   67
                                                                                                                     6.2
                                                                                                                            6.6
832173
            6.3 2.2
                       0.4
                             1.0
                                   9.9
                                        8.1
                                                -- 18.0 15.9
          (- - - - SPODIC HORIZON CRITERIA - - - - )
                                                                                 (------)
           (- -NA PYROPHOSPHATE EXTRACTABLE- -) INDEX
                                                                                (-----)
            C FE AL FE+AL FE+AL AL+C OF
                                                                                 (--- X-RAY - - - -)(- -DTA - -) TOTAL DOM
       HZN (- -DIVIDED BY- -) ACCUM
NO. 6A4A 6C8A 6G10 DI-CI PCT PCT
SAMPLE HZN
                                                                                (- - - - - <2U - - - )(- -<2U - -) RES WEATH
 NO.
                                                                                 7A21 7A21 7A21 7A3 7A3 7B1A 7B1A
           <- PCT OF <2MM-> FE+AL CLAY CLAY
                                                                                 832167
832168
                                                                                 MI 1 NX 6
832169
                  0.1 0.3 0.2
                                   0.1
                                               914
                       0.2
832170
        4
                  0.1
                             0.2 TR
                                               218
                                                                                 KH 1 FD 1 NX 6
                                                                                                         KH16
832171
        5
                         TR
832172
                         TR
                                                                                 KH 3 MI 1 CL 1 FD 1 KH59
832173
                         TR
```

FAMILY CONTROL SECTION: DEPTH 25-100 PCT CLAY 6 PCT .1-75MM 28. SPODIC HORIZON: INDEX OF ACCUMUL O ANALYSES: S= ALL ON SIEVED <2MM BASIS V= 75-20MM FROM VOLUME ESTIMATES MINERALOGY: KIND OF MINERAL MI MICA NX AMORPHOUS KH HALLOYSITE FD FELDSPAR CL CHLORITE

RELATIVE AMOUNT 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

ECUADOR 012 S83FN-325-012

Page 3 of 4

NSSL
SUPPLEMENTAL DATA SHEET

EXT 15 P (ACID OXALATE) EXTR PH P BAR RETENT AL SI FE AL KC1 BRAY1 MOIST (KOH)	
SAMPLE MAN 654 ADA6 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4	_
NO. NO. PPM <	í
832167 1 27.2 70 0.8 0.4 0.6 0.6	
832168 2 31.6 85 0.9 0.6 0.7 1.2	
832169 3 25.0 05	
832170 4 21 5 42	
832171 5 20.4 46	
832172 6 25.6 53	
832173 7 42.0 55 0.1 0.1 0.5 0.2	
832173 7 43.8 55 0.1 0.1 0.4 0.2	

^{*} NEW ZEALAND PROCEDURE

Page 4 of 4

CP83FN128

									_										
		(((MINERALOGY-										 X-RA	Y	(DTA) (TOT A) ANAL)	
SAMPLE NO.	HZN NO.	FA 7Bla		Bla 7Bla	7B1A	7B1A	7B1A	7BlA 7	Bla 7Bla	7BlA	7A2I	7A2I	7A2I	7A2I	7a2 I	7A 3	7A 3	K20	FE 607A
83P2167 83P2168 83P2169	2	VFS	0	T62 GA25	GC 7	GS 6		-			MI 1	NX 6						0.3	2.6
83P2170 83P2171	_										кн 1	FD 1	NX 6			KH16		0.5	3.9
83P2172 83P2173	-										КН 3	MI 1	CL 1	FD 1	NX 6	KH59		0.3	4.4

ANALYSES: S=ALL ON SIEVED < 2mm BASIS

MINERALOGY: FA = FRACTION ANALYZED RE = RESISTANT

KIND OF MINERAL: GA = GLASS AGGREGATES GS = GLASS MI = MICA OT = OTHER NX = NON-CRYSTALIJINE GC = GLASS COATED GRAIN

FD = FELDSPAR KH = HALLOYSITE CL = CHLORITE

RELATIVE AMOUNT: 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

MINERALOGY BASED ON SAND/SJLT:

MINERALOGY BASED ON CLAY:

FAMILY PLACEMENT:

COMMENTS:

Series: Ecuador No. 13 115 Date: 8/82

Pedon No: S83FN-325-013

Taxonomy: Fine, mixed, isohyperthermic Pachic Paleustoll

Latitude: Longitude:

Location: This pedon is located 20 km west of Quevedo about 6 km from El

Empalme.

Physiography: Fan Terrace Geomorphic Position: Summit

Slope and Aspect: 1% planar-planar Elevation: 100 m M.S.L

Microrelief: summit

Air Temp. Summer: 23C Winter: 25C (Est from nearby stations.)

Precipitation: 2000 mm Ustic moisture regime.

Water Table: > 10 m

Drainage: Moderately well drained Fermeability: Moderately slow

Stoniness: nonstony

Land Use: Forest, ungrazed Erosion or Deposition: none

Parent Material: multiple recent ash deposits over a very old ash deposit with a strongly developed argillic horizon.

Described by: T. Cook, G. Del Posso, and Soil Survey Staff.

Remarks: Date described: January 28, 1983. Field Moisture, 0-165 cm moist.

About 1% broken brick and stone artifacts from 0 to 77 cm.

Vegetation: native tropical forest of quaua, cocoa, coffee, and other broad leaf trees.

- Al 0 30 cm Black (10YR 2/1, moist) loam; moderate medium and coarse subangular blocky structure; very friable, non-sticky and non-plastic; weakly smeary, very moist or wet; common medium and coarse roots; very fine, common fine, and medium tubular pores; gradual wavy boundary.
- Bw 30 55 cm Dark brown (7.5YR 3/2, moist) and very dark brown (10YR 2/2, moist) silt loam; moderate fine and medium subangular blocky structure; very friable. slightly sticky and slightly plastic; weakly smeary, very moist or wet; few medium and coarse roots; many very fine, fine, and medium, and few coarse tubular pores; diffuse wavy boundary.
- 2Btbl 55 77 cm Very dark brown (10YR 2/2, moist) silty clay: loam, strong medium and coarse subangular blocky structure; friable, slightly sticky and slightly plastic; few fine and medium roots; many very fine and fine tubular pores; few thin patchy stress oriented coats on peds; few thin clay films on peds and in pores; diffuse wavy boundary.
- 2Btb2 77 107 cm Dark brown (7.5YR 3/4, moist) clay; moderate coarse prismatic structure parting to moderate coarse angular blocky; firm, sticky and plastic few fine and medium roots; common very fine and fine, and few medium tubular pores; many thin continuous stress oriented coats on peds; many thin clay films on peds; gradual wavy boundary.
- 2Btb3 107 165 cm Variegated dark yellowish brown (10YR 4/4, moist), red (2.5YR 4/8, moist) and light olive gray (5Y 6/2, moist) clay: moderate coarse and very coarse prismatic structure parting to moderate medium and coarse angular blocky; firm, sticky and plastic; few very fine and fine roots; common very fine and fine tubular pores; many thin continuous stress oriented coats on peds; many thin clay films on peds.

ECUADOR 013

LAB CLASSIF: FINE, MIXED, ISOHYPERTHERMIC PACHIC PALEUSTOLL

S 83FN-325 -013

DATE 09/25/84

SAMPLE NO. 83P2181-2185 PEDON NO. 83P 469

ECUADOR-SMSS

PROJECT NO. 83P 83

U. S. DEPARTMENT OF AGRICULTURE SOIL CONSERVATION SERVICE NATIONAL SOIL SURVEY LABORATORY LINCOLN, NEBRASKA 68508

GENERAL METHODS 131A, 2A1, 2B

-1-- -2-- -3-- -4-- -5-- -6-- -7-- -8-- -9-- -10- -11- -12- -13- -14- -15- -16- -17- -18- -19- -20-(- - -TOTAL - - -)(- -CLAY- -)(- -SILT- -)(- - - - - - -SAND- - - - - -)(-COARSE FRACTIONS(MM)-)(>2MM CLAY SILT SAND FINE CO3 FINE COARSE VF F M C VC ---- WEIGHT --- WT LT .002 .05 LT LT .002 .02 .05 .10 .25 .5 1 2 5 20 .1- PCI 0 .002 -.05 -2 .0002 .002 -.02 -.05 -.10 -.25 -.50 -1 -2 -5 -20 -.75 .75 WHOLE SAMPLE HZN DEPTH HORIZON .002 -.05 -2 .0002 .002 -.02 -.05 -.10 -.25 -.50 -1 -2 -5 -20 -75 75 WHOLE NO. NO. (CM) **** SEE FOOTNOTE REGARDING PSDA BELOW ***** 832181 18 0- 30 A1 10.0 59.5 30.5 37.1 22.4 20.8 8.5 0.6 0.6 832132 28 30- 55 12 BW 13.1 56.2 30.7 36.0 20.2 19.1 10.9 0.5 0.2 13 832183 38 55- 77 **2BTB1** 36.5 43.5 20.0 28.4 15.1 11.5 7.6 0.5 0.30.1 TR 10 2 832184 77-107 2B1B2 70.0 19.7 10.3 13.0 6.7 4.2 3.7 1.4 0.7 0.3 TR 832185 58 107-165 6 2BTB3 84.8 11.8 3.4 6.8 5.0 2.0 1.2 0.2 -- TR ORGN TOTAL EXTR TOTAL (- - DITH-CIT - -)(RATIO/CLAY)(ATTERBERG)(- BULK DENSITY -) COLE (- - -WATER CONTENT - -) WRD C N P S EXTRACTABLE 15 - LIMITS - FIELD 1/3 OVEN WHOLE FIELD 1/10 1/3 15 WHOLE SAMPLE HZN BAR LL PI MOIST BAR DRY SOIL MOIST BAR FF AL MN CEC BAR BAR SOIL NO. NO. 6A1C 6B3A 6R3A 6C2B 6G7A 6D2A 8D1 4F 4A3A 4A1D 4A1H 4D1 4B4 4B1C 8D1 4F1 4B1C 482A 4C1 PCT <0.4MM <- - G/CC - - -> CM/CM <- - - PCT OF <2MM - -> CM/CM 832181 1 3.36 0.344 2.0 0.4 3.01 2.06 0.82 0.94 0.046 50.5 20.6 0.24 832182 1.20 0.113 2.2 0.3 1.33 1.38 1.21 1.27 0.016 33.0 18.1 0.18 832183 0.33 2.9 0.3 0.30 0.59 1.35 1.46 0.026 28.0 21.7 0.08 832184 4 0.30 5.2 0.5 0.21 0.51 1.14 1.42 0.076 46.0 35.6 0.12 832185 5 0.27 5.1 0.5 0.17 0.49 1.04 1.37 0.096 52.9 41.2 0.12

^{*} CAUTION SHOULD BE USED WITH PSDA DATA WHEN USED FOR MAKING ENGINEERING INTERPERTATIONS BECAUSE OF INCOMPLETE DISPERSION OF AMORPHOUS MATERIAL. THEY ARE PRESENTED TO AID IN CLASSIFICATION AND ASSESSMENT OF MINERALOGY. REFER TO PEDON DESCRIPTION FOR FILED TEXTURE.

	ECUADOR -1	013	-3	-4	S 83FN- -5	·325 -0 -6	13 -7	AD -8-	TE 09/ -9	/25/84 -10-	-11-	PEDON -12-	NO. 8	3P 469 -14-	NA -15-	TIONAL -16-	SOIL -17-	CHRVE	2 OF 4 Y LABOF -19-	VATABV
SAMPLE NO.	CA HZN 5B5A NO. 6N2E	5B5A	NA 585A 6P2B	К 5В5А бQ2В	SUM BASES	ITY 6H5A	AL 6G9A	SUM CATS 5A3A	NH4- OAC 5A8B	BASES + AL 5A3B	SAT 5G1	SUM 5C3	NH4 OAC 5C1	CACO3	OHMS /CM		MMHOS	S NAF	PH - CACL2 .01M 8C1F 1:2	H20
832181 832182 832183 832184 832185	1 19.3 2 10.4 3 6.3 4 6.4 5 7.6	1.2 1.6 2.4	0.2	0.7 0.5 0.3	23.2 12.6 8.6 9.5 11.0	14.3		42.0 26.9 16.3 19.7 18.3	17.4			55 47 53 48 60	77 72 79 66 74					9.5 9.9 9.4 9.3 9.6	5.9 6.2 6.2 5.6 6.3	6.3 6.6 6.7 6.1 6.7
SAMPLE NO.	(N C HZN NO. 6A4A	SPOR A PYROR FE 6C8A	PHOSPHA AL 6G10	ATE EXT FE+AL DI-CI	RACTAB FE+AL VIDED PCT	LE) AL+C BY) PCT	INDEX OF)	~ • • • • • •				((7A21	- X-I	- CLA RAY - <2U - 7A21	Y)()(7A21	 (D1 (<2 7A3) [A] [U] 7A3	TOTAL RES 7B1A	DOM WEATH 7B1A
832181 832182 832183 832184 832185	1 2 3 4 5	0.2 0.1 	0.1	0.2 TR 0.1	TR 		1110						KH 2	FD 1 FD 1 CR 2	CR 1	_	KH17 KH44 KH67			

FAMILY CONTROL SECTION: DEPTH 25-100 PCT CLAY 37 PCT .1-75MM 10. SPODIC HORIZON: INDEX OF ACCUMUL O ANALYSES: S= ALL ON SIEVED <2MM BASIS V= 75-20MM FROM VOLUME ESTIMATES MINERALOGY: KIND OF MINERAL KH HALLOYSITE FD FELDSPAR NX AMORPHOUS CR CRISTOBALITE

6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

RELATIVE AMOUNT

S U P P L E M E N T A L D A T A S H E E T

		EXT P BRAY1	15 BAR MOIST	P RETENT	(ACID	OXAL SI	 ATE) FE	EXTR AL	pH KCl
SAMPLE NO.	HZN NO.	6S3 PPM	4B2b <	*	* -PCT OF <	* 2 MM - ·	*	(KOH) >	8Clg
832181 832182	1 2		28.4 23.6	74 64	0.6 0.4	0.5 0.4	1.1	0.5	
832183 832184	3 4		34.1 44.4	50 69	0.1 0.2	0.1	0.8 0.5 0.2	0.4	
832185 	5 		53.0	60	0.2	0.1	0.1	0.2 0.2	

^{*} NEW ZEALAND PROCEDURE

CP83FN128

Crosini	20					
				mineralogy)
		(() (CLAY-	(DTA) (TOT ANAL)
SAMPLE NO.	HZN NO.	FA 7BlA	RE 7BlA 7BlA <		7A2I 7A2I 7A2I 7A2I 7A2I	K2O FE 7A3 7A3 6Q3A 6C7A
83P2181 83P2182 83P2183 83P2184 83P2184	2 3 4	VFS	OT65	GA26 GC 5 GS 4	KH 1 FD 1 NX 6 KH 2 FD 1 CR 1 NX 6 KH 3 CR 2 NX 6	KH17 0.2 4.2 KH44 0.3 4.9 KH67 0.2 5.0

ANALYSES: S=ALL ON SIEVED < 2mm BASIS

MINERALOGY: FA = FRACTION ANALYZED RE = RESISTANT

KIND OF MINERAL: FD = FELDSPAR GA = GLASS AGGREGATES GS = GLASS OT = OTHER KH = HALLOYSITE NX = NON-CRYSTALLINE

GC = GLASS COATED GRAIN CR = CRISTOBALITE

RELATIVE AMOUNT: 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

MINERALOGY BASED ON SAND/SILT:

MINERALOGY BASED ON CLAY:

FAMILY PLACEMENT:

COMMENTS:

Appendix I

Extra Data and Descriptions

This appendix contains analytical data and pedon descriptions for pedons that were sampled but not used on the tours because of constrains of time.

Series: Equador No. 2 122 Date: 6/83

Pedon No: S83FN-325-002

Taxonomy: Ashy, iscresic Mollic Vitrandept Latitude: 2°07'20" Longitude: 78°43'50" Location: About 5-6 km south of Palmira. Physiography: In uplands in eplian dunes

Geomorphic Position: On a slope

Slope and Aspect: 13% W planar-convex Elevation: 3200 m M.S.L

Microrelief: undulating
Air Temp. Summer: Winter:

Precipitation: 500 mm Ustic moisture regime.

Water Table: > 10 m

Drainage: Well drained Permeability: Moderately rapid

Stoniness: nonstony

Land Use: Grassland or grazing land

Erosion or Deposition: severe wind erosion

Parent Material: ash over unrelated tuff breccia

Described by: T. Cook, G. Del Posso, and Soil Survey Staff.

Remarks: Date described: January 29, 1983. Field Moisture, 0-126 moist. Vegetation: clumps of stipa sp. separated by bare soil.

- Al 0 53 cm Dark yellowish brown (10YR 4/4, rubbed dry), loamy fine sand; black (10YR 2/1, moist); structureless; very friable, non-sticky and non-plastic; few very fine and fine roots; few very fine tubular and many very fine interstital pores; diffuse smooth boundary.
- A2 53 90 cm Very dark brown (10YR 2/2, moist); loamy fine sand, structureless; very friable, non-sticky and non-plastic; few very fine roots; few very fine tubular and many very fine interstital pores; gradual smooth boundary.
- 90 110 cm Very dark gray (10YR 3/1, moist); loamy fine sand, structureless; friable, non-sticky and non-plastic; very weakly cemented; few very fine roots; few very fine tubular and many very fine interstitial pores; 1% 2-5 mm rock fragments; clear wavy boundary.
- 2C 110 120 cm White (10YR 8/1, moist); gravelly loamy fine sand, structureless; firm, non-sticky and non-plastic; weakly to moderately cemented; few very fine tubular and few very fine interstitial pores; coarse fragments, 30% 2-30 mm; common fine white (N8/0, moist) filaments; abrupt wavy boundary.
- 3R 120 126 cm unweathered bedrock, tuff breccia. There is a thin (0-3 cm) discontinuous dark gray (10 YR 4/1, moist) layer just above the hard rock.

PAGE 1 OF 4 PAGES

U. S. DEPARTMENT OF AGRICULTURE

18.8

16.9

23.8

24.3

4.1 0.21

4.7 0.18

5.3 0.26

5.7 0.17

SOIL CONSERVATION SERVICE

ECHADOR 002

1 0.60 0.050

2 0.37 0.026

3 0.26

4 0.27

FOR FIELD TEXTURE.

832191

832192

832193

832194

LAB CLASSIF: ASHY, ISOMESIC MOLLIC VITRANDEPT

S 83FN-325 -002 DATE 09/25/84 SAMPLE NO. 83P2191-2195 PEDON NO. 83P 471

0.5 0.1

0.1

0.1

0.1

0.6

0.6

0.6

ECUADOR-SMSS PROJECT NO. 83P 83 NATIONAL SOIL SURVEY LABORATORY
LINCOLN, NEBRASKA 68508

GENERAL METHODS 1B1A, 2A1, 2B

-1-- -2-- -3-- -4-- -5-- -6-- -7-- -8-- -9-- -10- -11- -12- -13- -14- -15- -16- -17- -18- -19- -20-(- - -TOTAL - - -)(- -CLAY- -)(- -SILT- -)(- - - - - -SAND- - - - - -)(-COARSE FRACTIONS(MM)-)(>2MM CLAY SILT SAND FINE CO3 FINE COARSE VF F M C VC - - - - WEIGHT - - - - WT SAMPLE HZN DEPTH HORIZON LT .002 .05 LT NO. NO. (CM) **** SEE FOOTNOTE REGARDING PSDA BELOW ***** 832191 1S 0-53 0.9 30.1 69.0 1.6 A1 8.3 21.8 44.7 22.6 0.1 832192 2S 53- 90 1.0 27.7 71.3 5.9 21.8 51.3 17.7 -- TR A2 1.9 0.4 20 --3S 90-110 A.3 0.9 29.4 69.7 8.6 20.8 44.7 19.2 1.2 1 TR 27 3 832193 3.1 1.5 832194 45 110-120 2C 1.2 31.4 67.4 9.7 21.7 42.1 16.7 3.0 2.7 3 43V 63 51 832195 5\$ 120-126 3R ORGN TOTAL EXTR TOTAL (- - DITH-CIT - -) (RATIO/CLAY) (ATTERBERG) (- BULK DENS!TY -) COLE (- - -WATER CONTENT - -) WRD C N P S EXTRACTABLE 15 - LIMITS - FIELD 1/3 OVEN WHOLE FIELD 1/10 1/3 15 WHOLE SAMPLE HZN FF AL MN CEC BAR LL PI MOIST BAR DRY SOIL MOIST BAR BAR BAR SOIL NO. 6A1C 6B3A 6R3A 5C2B 6G7A 6D2A 8D1 8D1 4F1 4F 4A3A 4A1D 4A1H 4D1 4B4 4B1C 4B1C 4B2A 4C1 PCT <0.4MM <- - G/CC - - -> CM/CM <- - -PCT OF <2MM - -> CM/CM

PSDA IN ANDERTS SHOULD NOT BE USED FOR MAKING ENGINEERING INTERPERTATIONS BECAUSE OF INCOMPLETE DISPERSION OF AMORPHOUS MATERIAL. THEY ARE PRESENTED TO AID IN CLASSIFICATION AND ASSESSMENT OF MINERALOGY. REFER TO PEDON DESCRIPTION

1.40 1.40

1.46 1.47 0.002

1.41 1.43 0.005

1.40 1.43 0.005

6.33 4.56

6.70 4.70

9.22 5.89

7.75 4.75

	ECU	JADOR (002 -2	-3	-4	S 83FN- -5		02 -7	DA -8	TE 09/	/25/84 -10-	-11-	PEDON -12-	NO. -13-	83P 471 -14-	NA -15-	TIONAL -16-	SOIL -17-		2 OF 4 Y LABOR -19-	RATORY
SAMPLE NO.	HZN	5B5A	5B5A 602D	5B5A	5B5A 6Q2B	BASES	6H5A	AL 6G9A	SUM CATS 5A3A	NH4- OAC 5A8B) BASES + AL 5A3B >	AL SAT 5G1	SUM 5C3	NH4 OAC 5C1	<2MM			COND. MMHOS /CM 81	(6 NAF 8C1D		H20 801F 1:1
832191 832192 832193 832194 832195	1 2 3 4 5	4.6 4.9 6.1 23.4	1.9 2.5 3.3 3.8	0.2 0.3 0.4 0.4	0.3 0.8 1.0 1.2	7.0 8.5 10.8 28.8	1.0	 TR	8.0 8.5 10.8 28.8	5.7 6.7 8.3 9.3			87 100 100 100	100 100 100 100					9.8 9.8 10.0 10.3	7.0 7.4 7.6 8.5	7.4 7.8 8.2 8.7
													((- CLA	Y		1		•
SAMPLE NO.	HZN NO.												(7A21	- ~ X-F < 7A21 ELATIVE	20 7A21) 7A21	7A3	7A3	RES 1	WEATH 781A
832191 832192 832193 832194 832195	1 2 3 4 5													NX 6 NX 6 NX 6				-	•	0,	

FAMILY CONTROL SECTION: DEPTH 25-100 PCT CLAY 1 PCT .1-75MM 23 ANALYSES: S= ALL ON SIEVED <2MM BASIS V=75-20MM FROM VOLUME ESTIMATES MINERALOGY: KIND OF MINERAL NX AMORPHOUS

RELATIVE AMOUNT 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

		EXT P BRAY1	15 BAR MOIST	P RETENT	(ACII AL	O OXAL	ATE) FE	EXTR AL (KOH)	pH KCl
SAMPLE NO.	HZN NO.	6S3 PPM	4B2b	*	* *	*	*		8Clg
					-PCT OF <	<2MM		>	
832191	1		4.2	34	0.3	0.2	0.3	0.1	
832192	2		5.1	17	0.2	0.1	0.7	<0.1	
832193	3		5.9	22	0.2	0.2	0.7	<0.1	
832194	4		6.6	14	0.2	0.2	0.8	<0.1	
832195	5			-					

^{*} NEW ZEALAND PROCEDURE

CP83FN128

33		(((X-RAY) (DTA) (TOT ANAL)
SAMPLE NO.	HZN NO.	FA 7BlA	RE	7821 7821 7821 7821 782 782 782 782 782 782 782 782 782 782
83P2191 83P2192 83P2193 83P2194 83P2195	2 3 4	VFS VFS VFS	GA68 OT17 GC14 GS 1 GA64 GC18 OT17 GS 1 GA63 OT18 GC17 GS 2	NX 6 0.5 1.2 NX 6 0.5 0.5 NX 6 0.4 0.7

ANALYSES: S=ALL ON SIEVED < 2mm BASIS

MINERALOGY: FA = FRACTION ANALYZED RE = RESISTANT

KIND OF MINERAL: GA = GLASS AGGREGATES GS = GLASS OT = OTHER NX = NON-CRYSTALLINE GC = GLASS COATED GRAIN

RELATIVE AMOUNT: 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

MINERALOGY BASED ON SAND/SILT:

MINERALOGY BASED ON CLAY:

FAMILY PLACEMENT:

COMMENTS:

2

Series: Ecuador No. 3 127 Pate: 6/83

Pedon No: S83FN-325-003

Taxonomy: Coarse-loamy, mixed, isomesic Typic Humitropept

Latitude: 1°25'00" Longitude: 78°37'15"

Location: About 7 km south of Mocha. Urbina-Paramo area

Physiography: in uplands

Geomorphic Position: On a slope

Slope and Aspect: 10% W convex-convex Elevation: 3400 m M.S.L

Microrelief: hummucky

Air Temp. Summer: Winter:

Precipitation: 1000 mm Udic moisture regime.

Water Table: 5 to 10 meters

Drainage: Well drained Permeability: moderate

Stoniness: nonstony

Land Use: Grassland or grazing land

Erosion or Deposition: none

Parent Material: ash

Described by: T. Cook, G. Del Posso, and Soil Survey Staff.

Remarks: Date described: January 29, 1983. Field Moisture, 0 to 175 cm wet. Worm casts occupy 30-40% of the soil mass; 15-20 mm holes made by insects and grubs occupy 1% of the pedon.

Vegetation: cushion plant.

- 0 10 0 cm decomposed material; weak medium to coarse subangular blocky structure; loose; clear smooth boundary. This layer varies in thickness from 5 to 35 cm within a horizontal distance of 10 to 100 cm.
- Al 0 15 cm Very dark brown (10YR 2/2) loam; weak medium and coarse subangular block structure; friable, non-sticky and non-plastic; weakly smeary, very moist or wet; common very fine, fine, and medium roots; common very fine and fine tubular pores; gradual smooth boundary.
- 15 54 cm Black (10YR 2/1) loam; common fine faint dark brown (10YR 3/3) mottles; weak very coarse subangular blocky structure parting to weak medium subangular blocky; friable, non-sticky and non-plastic; weakly smeary, very moist or wet; common very fine, fine, and medium roots; common very fine and fine tubular pores; gradual smooth boundary.
- A3 54 98 cm Black (10YR 2/1) sandy loam; common fine faint very dark grayish brown (10YR 3/2) mottles; weak coarse prismatic structure parting to weak coarse angular blocky; friable, non-sticky and non-plastic; weakly smeary, very moist or wet; few very fine, fine, and medium roots; common very fine and fine tubular and common very fine interstitial pores; diffuse smooth boundary.
- 98 137 cm Black (10YR 2/1) sandy loam; common fine faint very dark gray (10YR 3/1) mottles; weak coarse prismatic structure parting to weak coarse angular blocky; friable, non-sticky and non-plastic; weakly smeary, very moist or wet; few very fine, fine, and medium roots; common very fine and fine tubular and common very fine interstitial pores; gradual smooth boundary.
- 2Ab 137 175 cm Black (N 2/0) loam; massive; friable, slightly sticky and slightly plastic; weakly smeary, very moist or wet; many thin continuous organic coats in pores; few very fine, fine, and medium roots; common very fine, fine, and medium tubular and few very fine interstitial pores.

ECUADOR 003

LAB CLASSIF: COARSE-LOAMY, MIXED, ISOMESIC TYPIC HUMITROPEPT

S 83FN-325 -003

1.1

0.3

ECUADOR-SMSS

832204

5 3.54

DATE 09/25/84 SAMPLE NO. 83P2200-2204 PEDON NO. 83P 473

PROJECT NO. 83P 83

1.03 1.13 0.031

U. S. DEPARTMENT OF AGRICULTURE SOIL CONSERVATION SERVICE NATIONAL SOIL SURVEY LABORATORY LINCOLN. NEBRASKA 68508

44.8

12.1 0.34

PAGE 1 OF 4 PAGES

GENERAL METHODS 1B1A, 2A1, 2B

-6-- -7-- -8-- -9-- -10- -11- -12- -13- -14- -15- -16- -17- -18- -19- -20-(- - -TOTAL - - -)(- -CLAY- -)(- -SILT- -)(- - - - - - -SAND- - - - - -)(-COARSE FRACTIONS(MM)-)(>2MM CLAY SILT SAND FINE CO3 FINE COARSE VF F M C VC - - - - WEIGHT - - - WT LT .002 .05 LT LT .002 .02 .05 .10 .25 .002 -.05 -2 .0002 .002 -.02 -.05 -.10 -.25 -.50 SAMPLE DEPTH HORIZON .5 1 2 .1- PCI Q NO. NO. (CM) -2 -5 -20 -75 75 WHOLE **** SEE FOOTNOTE REGARDING PSDA BELOW **** 832200 1 S 0- 15 A1 1.3 36.5 62.2 15.1 21.4 22.7 24.6 8.1 3.0 3.8 TR TR 832201 28 15- 54 39 TR A2 1.5 3/.7 60.8 15.6 22.1 26.8 25.3 6.0 2.1 0.6 TR 34 --832202 3 S 54- 98 Α3 1.3 33.3 65.4 17.1 16.2 24.1 28.1 9.3 3.1 0.8 TR TR 4.1 TR 832203 4S 98-137 ΑU 0.8 30.0 69.2 16.2 13.8 21.5 26.8 14.7 5.5 0.7 TR 1.17 832204 48 TR 5S 137-175 1.8 47.9 50.3 2AB 27.0 20.9 26.8 1,.3 5.8 1.4 TR 23 --ORGN TOTAL EXTR TOTAL (- - DITH-CIT - -)(RATIO/CLAY)(ATTERBERG)(- BULK DENSITY -) COLE (- - -WATER CONTENT - -) WRD N P S 15 - LIMITS - FIELD 1/3 OVEN WHOLE FIELD 1/10 EXTRACTABLE 1/3 15 WHOLF SAMPLE HZN ΓE AL MN CEC BAR LL PI MOIST BAR DRY SOIL MOIST BAR BAR BAR SOIL NO. NO. 6A1C 6B3A 6R3A 6C2B 6G7A 6D2A 8D1 8D1 4F1 4 F 4A3A 4A1D 4A1H 4D1 4B4 <- - - - - - - - PCT OF <2MM - - - - - -> 4B1C 4B1C 482A 4C1 PCT <0.4MM <- - G/CC - - -> CM/CM <- - -PCT OF <2MM - -> CM/CM 832200 1 7.83 0.478 0.50.313.23 15.69 0.79 0.98 0.074 66.9 20.4 0.37 2 2.99 0.219 832201 0.8 0.3 5.93 5.87 1.13 1.22 0.026 38.2 8.8 0.33 832202 3 1.64 0.6 0.2 4.62 4.08 1.25 1.31 0.016 27.8 5.3 ± 0.28 832203 4 1.23 0.5 0.2 7.00 5.88 1.27 1.32 0.013 25.2 4.7 0.26

9.78 6.72

^{*} CAUTION SHOULD BE USED WITH PSDA DATA WHEN USED FOR MAKING ENGINEERING INTERPERTATIONS BECAUSE OF INCOMPLETE DISPERSION OF AMORPHOUS MATERIAL. THEY ARE PRESENTED TO AID IN CLASSIFICATION AND ASSESSMENT OF MINERALOGY. REFER TO PEDON DESCRIPTION FOR FIELD TEXTURE.

```
PAGE 2 OF 4 PAGES
      ECUADOR 003 S 83FN-325 -003 DATE 09/25/84 PEDON NO. 83P 473 NATIONAL SOIL SURVEY LABORATORY
         -1-- -2-- -3-- -4-- -5-- -6-- -7-- -8-- -9-- -10- -11- -12- -13- -14- -15- -16- -17- -18- -19- -20-
        (- NH4OAC EXTRACTABLE BASES -) ACID- EXTR (- - - - CEC - - -) AL -BASE SAT- CO3 AS RES.
                                                                                      COND.(- - - - PH - - -)
          CA MG NA K SUM ITY AL SUM NH4- BASES SAT
                                                              SUM NH4 CACO3 OHMS
                                                                                      MMHOS NAF CACL2 H20
SAMPLE HZN 5B5A 5B5A 5B5A 5B5A BASES
                                           CATS OAC + AL
                                                                   OAC <2MM /CM
                                                                                    /CM .01M
      NO. 6N2E 602D 6P2B 6Q2B
                                 6H5A 6G9A 5A3A 5A8B 5A3B 5G1
 NO.
                                                              5C3 5C1 6E1G 8E1
                                                                                      81 8C1D 8C1F 8C1F
          1:2 1:1
832200
                   0.3
          4.2
               2.3
                        0.2
                            7.0 23.4
                                       0.7 30.4 17.2
                                      0.7 30.4 17.2 7.7 9
0.3 16.9 8.9 4.4 7
0.2 12.7 6.0 3.6 6
                                                                                          9.8
                                                                                               4.8
                                                                                                     5.4
832201
          2.1
               1.6
                   0.3
                        0.1
                             4.1 12.8
                                                               24
                                                                    46
                                                                                          10.2 5.3 6.0
832202
         1.6
              1.6
                   0.2
                        TR
                            3.4 9.3
                                                               27
                                                                   57
                                                                                          10.3 5.6
                                                                                                   6.1
832203
              1.7
                        TR
                                       0.1 10.7 5.6
         1.6
                   0.2
                            3.5
                                7.2
                                                    3.6
                                                        3 33
                                                                    62
                                                                                          10.1
                                                                                                5.7
                                                                                                   6.2
832204
          6.9
               5.5
                   0.3
                        0.2 12.9 16.0
                                      -- 28.9 17.6
                                                               45
                                                                    73
                                                                                           9.8
                                                                                                   6.3
         (- - - - SPODIC HORIZON CRITERIA - - - - )
                                                                  (- -NA PYROPHOSPHATE EXTRACTABLE- -) INDEX
                                                                  (-----)
          C FE AL FE+AL FE+AL AL+C OF
                                                                  (---- X-RAY - - - -)(- -DTA - -) TOTAL DOM
                                                                  (- - - - - <2U - - - -) RES WEATH
SAMPLE HZN
                      (- -DIVIDED BY- -) ACCUM
      NO. 6A4A 6C8A 6G10 DI-CI PCT PCT
NO.
                                                                  7A21 7A21 7A21 7A21 7A3 7A3 7B1A 7B1A
         <- PCT OF <2MM-> FE+AL CLAY CLAY
                                                                  <- RELATIVE AMOUNTS -> <- - - - - PCT - - - ->
832200
               0.4 0.3
                        0.9
                             0.5
                                       446
832201
               0.3
                   0.2
                        0.5
                             0.3
                                       630
                                                                  NX 6
832202
               0.2
                   0.2
                        0.5
                             0.3
                                      530
                                                                  NX 6
832203
                   0.1
                        0.3
               0.1
                             0.3
                                      402
832204
               0.3 0.3
                        0.4
                             0.3
                                      1064
                                                                  VR 1 NX 6
```

FAMILY CONTROL SECTION: DEF(H 25-100 PCT CLAY 1 PCT .1-75MM 39. SPODIC HORIZON: INDEX OF ACCUMUL 1234 ANALYSES: S= ALL ON SIEVED <2MM BASIS MINERALOGY: KIND OF MINERAL NX AMORPHOUS VR VERMICULITE

RELATIVE AMOUNT 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

Page 3 of 4

ECNYDOK 003 283EM-352-003

* NEW ZEALAND PROCEDURE

T S S N

SUPPLEMENTAL DATA SHEET

	4.0 4.0 4.0 4.0	2.0 3.0 2.0 4.0 7.0	E.0 E.0 E.0 E.0	ε.0 ε.0 ε.0 γ.0 γ.0	25 25 25 25 25	7.28 9.6 8.51 2.8		2 3 7 7	832204 832203 832203 832200
8СГД КСЈ БН	EXTR AL (KOH)	ATE) FE *	IS S OXAL,	AL AL *	P RETENT *	15 BAR MOIST 4B2b 	EXT P BRAY1 653	ON	SPMPLE NO.

CP83FN128

		(·	MINERA	LOGY)
						-	PTICAL ND/SIL)	()	(DTA)	(TOT A	
SAMPLE NO.	HZN NO.	PA 7Bla	RE 7Bla	7BlA	7BlA	7BlA	7B1A	7BlA	7BlA	7BlA	7BlA	7B1A		7A3 7A3	K20 6Q3A	FE 6C7A
83P2200 83P2201 83P2202 83P2203	2	VFS		GA59	GC19	OT14	GS 8						NX 6 NX 6		0.4 0.2	2.5
83P2204	5												VR 1 NX 6		0.2	1.7

ANALYSES: S=ALL ON SIEVED < 2mm BASIS

MINERALOGY: FA = FRACTION ANALYZED RE = RESISTANT

KIND OF MINERAL: GA = GLASS AGGREGATES GS = GLASS OT = OTHER NX = NON-CRYSTALLINE GC = GLASS COATED GRAIN

VR = VERMICULITE

RELATIVE AMOUNT: 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

MINERALOGY BASED ON SAND/SILT:

MINERALOGY BASED ON CLAY:

FAMILY PLACEMENT:

COMMENTS:

5

Series: Ecuador No. 7 Date: 6/83

Pedon No: S83FN-325-007

Taxonomy: Ashy over loamy, mixed, isothermic Typic Ustifluvent

Latitude: 0°05'25" Longitude: 78°25'00"

Location: about 15 km NE of Quito, in the Calderon area

Physiography: Terrace Geomorphic Position: On a slope

Slope and Aspect: 1% planar-planar Elevation: 2620 m M.S.L

Microrelief: smooth

Air Temp. Summer: Winter:

Precipitation: 600mm Ustic moisture regime.

Water Table: > 10 m

Drainage: Well drained Permeability: moderate
Stoniness: nonstony
Land Use: Idle land, adjacent land is cropped to corn

Erosion or Deposition: none

Parent Material: multiple ash deposits

Described by: T. Cook, G. Del Posso, and Soil Survey Staff.
Remarks: Date Described: January 24, 1983. Field Moisture, 0-10 cm dry, 10-67 cm slightly moist, 67-205 moist. All rock fragments are pumice. From 0 to 35 cm appears to be distrubed by artificial digging or scraping, therefore did not seperate a thin surface horizon.

Vegetation: Planted eucalyptus grove, chilca, herbs and kikuyu grass.

- A 0 35 cm Very dark grayish brown (10YR 3/2) loamy sand, light brownish gray (10YR 6/2, dry); massive; very friable, non-sticky and non-plastic; few very fine and common medium and coarse roots; many very fine interstitial and few very fine tubular pores; 10% 2 to 70 mm fragments; gradual wavy boundary. Included are fragments of charcoal, brick, rags and other artifact pieces.
- · 35 67 cm Dark grayish brown (2.5Y 4/2) loamy sand, light grayish brown (10YR 6/2, dry) and grayish brown (10YR 5/2, dry); common fine distinct dark grayish brown (10YR 6/2, moist) mottles; massive; very friable, non-sticky and non-plastic; few very fine and common medium and coarse roots; many very fine interstitial and few very fine tubular pores; 10% 2 to 70 mm fragments; clear wavy boundary. Included are fragments of charcoal, brick, rags, and other artifact pieces.
- 67 90 cm Grayish brown (2.5Y 5/2) loamy coarse sand, light gray (5Y 7/1, dry); structureless; very friable, non-sticky and non-plastic; few very fine, fine, and medium roots; many very fine interstitial, common very fine and fine tubular pores, 15% fragments; abrupt wavy boundary. At bottom of this horizon there is a layer 5 to 20 mm thick with 40% pumice fragments 2 to 30 mm in size.
- Black (10YR 2/1) and very dark grayish brown (10YR3/2) 90 - 107 cm fine sandy loam; structureless; very friable, non-sticky and non-plastic; few, medium and coarse roots; many very fine interstitial and a few very fine tubular pores; clear wavy boundary.
- 107 143 cin Very dark grayish brown (10YR 3/2) and black (10YR 2/1) loam; structureless; friable, slightly sticky and slightly plastic; a few thin patchy organic coats on peds few fine, medium and coarse roots; many very fine interstitial and common very fine and fine tubular pores; gradual wavy boundary.
- 143 185 cm Very dark grayish brown (10YR 3/2) and black (10YR 2/1) loamy sand; structureless; friable, non-sticky and non-plastic; weakly smeary, very moist or wet; few very fine and fine roots; many very fine interstitial, few very fine tubular pores; abrupt smooth boundary.
- 185 205 cm Dark brown (10YR 3/3) and very dark grayish brown (10YR 3/2) loamy sand; structureless; firm, non-sticky and non-plastic; weakly cemented; few very fine and fine roots; common very fine interstitial pores. A black and very dark grayish brown (10YR 2/1, 3/2) layer 5 to 10 mm thick occures at top of horizon.

ECUADOR 007

LAB CLASSIF: ASHY OVER LOAMY, MIXED, ISOTHERMIC TYPIC USTIFLUVENT

S 83FN-325 -007

ECUADOR-SMSS

DATE 09/25/84

SAMPLE NO. 83P2077-2083 PEDON NO. 83P 455

PROJECT NO. 83P 83

U. S. DEPARTMENT OF AGRICULTURE SOIL CONSERVATION SERVICE NATIONAL SOIL SURVEY LABORATORY LINCOLN, NEBRASKA 68508

PAGE 1 OF 4 PAGES

GENERAL METHODS 181A, 2A1, 2B

		-1	-23-	4	-5	-6	-7	3-	-9	-10-	-11-	-12-	-13-	-14-	-15-	-16-	-17-	-18-	-19-	-20 -
SAMPLE NO.	HZN NO.		HORIZO	N LT .002	-TOTAL SILT .002 05	.05 -2	LT .0002	LT .002	.002	.02 05	.05 10	.10 25	.25 50	.5 -1	VC 1	· 2 -5	WE 5 -20	ACTIONS IGHT - 20 -75 <75MM(3	 .1- 75	WT PCT O WHOLE
**** S	EE F	OOTNOTE	REGARDIN	G PSDA	BELOW 1	****														
832077 832078 832079 832080 832081 832082 832083	1S 2S 3S 4S 5S 6S	0- 3 35- 6 67- 9	5 A 67 AC 90 2C1 97 3C2 93 3AB 95 3ACB		25.2 24.7 25.8 36.7 35.7 25.9	72.4 72.9 71.5 55.7 53.0 69.1 73.4			13.4 13.7 18.9 20.4 20.2 13.6 9.8	11.8 11.0 6.9 16.3 15.5 12.3	11.9 10.5 5.4 16.1 15.3 12.7 16.2	20.3 19.5 20.2 21.1 21.5 24.3 21.3	24.1 23.9 23.5 12.0 12.6 22.0 15.4	12.8 14.3 15.6 4.9 2.9 7.9 12.3	3.3 4.7 6.8 1.6 0.7 2.2 8.2	2 3 3 1 TR 1 5	1 1 1 TR TR TR	19V 19V 29V 	69 71 77 40 38 57	22 23 33 1 TR 1 5
SAMPLE NO.	HZN NO.	ORGN T C 6A1C	" '	6R3A	FE 6C2B	AL 6G7A	MN 6D2A	(RAT LO CEC 8D1	D/CLAY) 15 BAR 8D1	- LIM LL 4F1	RBERG) HITS - PI 4F	MOIST 4A3A	1/3 BAR	OVEN DRY 4A1H	WHOLE SOIL 4D1	FIELD MOIST 4B4	1/10 BAR 4B1C	CONTEN 1/3 BAR 4B1C OF <2MM	15 BAR 482A	WHOLE SOIL 4C1
832077 832078 832079 832080 832081 832082 832083	2 3 4 5 6	0.61 0 0.49 0 0.15 0 0.45 0.66 0.20	.044		0.3 0.3 0.2 0.5 0.4 0.4	TR TR TR TR O.1 TR TR		2.00 1.83 1.19 0.97 0.88 1.00 1.87	1.67 1.46 1.00 0.82 0.68 0.94 2.10				1.24 1.23 1.14 1.28 1.38 1.57	1.26 1.27 1.38 1.43 1.66	0.002 0.007 0.030 0.025 0.012 0.019 0.020			19.5 18.4 28.5 27.5 19.3 17.4 20.1	4.0 3.5 2.7 6.2 7.7 4.7	0.17 0.16 0.24 0.27 0.16 0.20

^{*} CAUTION SHOULD BE USED WITH PSDA DATA WHEN USED FOR MAKING ENGINEERING INTERPERTATIONS BECAUSE OF INCOMPLETE DISPERSION OF AMORPHOUS MATERIAL. THEY ARE PRESENTED TO AID IN CLASSIFICATION AND ASSESSMENT OF MINERALOGY. REFER TO PEDON DESCRIPTION FOR FIELD TEXTURE.

	ECUADOR	007		:	S 83FN-	325 -0	07	DA	JE 097	/25/8h		PEDON	NO	83P 455	N A	TIONAL			2 OF 4	
	-1	2	-3	-4	-5	-6	-7	-8	-9	-10-	-11-	-12-	-13-	-14-	-15-	-16-	-17-	-18-	r LABOI -19-	-20-
																				
SAMPLE NO.	HZN 5B54 NO. 6N2E	5B5A 602D	5B5A 6P2B	5B5A 6Q2B	BASES	6H5A	AL 6G9A	CATS 5A3A	NH4- OAC 5ABB	BASES + AL 5A3B	SAT 5G1	SUM 5C3	NH4 OAC 5C1	CO3 AS CACO3 <2MM 6E1G	OHMS /CM		MMHOS /CM	(NAF 8C1D	-PH - CACL2 .01M 8C1F 1:2	H20 8C1F
832077 832078 832079 832080 832081 832082 832083	1 7.6 2 3.9 3 2.4 4 5.8 5 7.8 6 3.4 7 3.8	0.6 0.7 1.3 1.7 0.9	0.1 TR 0.1 0.2 0.9 0.8 0.8	0.3 0.3 0.2 0.2 0.2 0.2	8.3 4.8 3.4 7.5 10.6 5.3 6.0	1.6 2.2 1.2 1.7 2.0 0.5 1.3	TR	9.9 7.0 4.6 9.2 12.6 5.8 7.3	4.8 4.4 3.2 7.4 9.9 5.0 5.8			84 69 74 82 84 91 82	100 100 100 100 100 100					9.6 9.4 8.4 8.8 9.1 8.6 8.1	7.6 7.0 6.8 6.9 7.3 7.4 7.8	7.7 7.3 7.8 7.6 8.1 8.5 8.4
	(SPO	OLC HOR	RIZON C	RITERL															
SAMPLE NO.	(N C HZN NO. 6A4A	A PYROF FE	PHOSPHA AL 6G10	TE EXT FE+AL Ol-Cl	RACTABI FE+AL VIDED I PCT	LE) AL+C BY) PCT	INDEX OF	,				(7A21	7A21	- CLAY RAY RAY RAZI	Y)() 7A21	7A3) A) 7A3	TOTAL RES 7B1A	DOM WEATH
832077 832078 832079	1 2 3	0.1	TR TR TR	0.3	TR								FD 1	NX 6						
832080 832081	5		TR TR										FD 1	NX 6						
832082 832083	6 7	0.1	0.1 TR	0.5	TR		139						NX 6							

FAMILY CONTROL SECTION: DEPTH 25-100 PCT CLAY 3 PCT .1-75MM 69. SPODIC HORIZON: INDEX OF ACCUMUL 0 ANALYSES: S= ALL ON SIEVED <2MM BASIS V= 75-20MM FROM VOLUME ESTIMATES NX AMORPHOUS

RELATIVE AMOUNT 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

N S S L
S U P P L E M E N T A L D A T A S H E E T

		EXT	15	Р			ATE)	EXTR	рН
		P BRAY1	BAR MOIST	RETENT	AL	SI	FE	AL (KOH)	KCl
SAMPLE	HZN	6 S 3	4B2b	*	*	*	*		8Clg
NO.	NO.	PPM	<		-PCT OF	<2MM-		>	
832077	1		3.3	25	0.2	0.1	0.2	<0.1	
832078	2		4.0	7	0.2	0.1	0.2	<0.1	
832079	3		1.7	17	0.1	<0.1	0.1	<0.1	
832080	4		6.8	10	0.1	<0.1	0.1	<0.1	
832081	5		9.3	30	0.2	0.1	0.1	<0.1	
832082	6		4.8	17	0.1	<0.1	0.1	<0.1	
832083	7		4.6	3	0.1	0.1	0.2	<0.1	

^{*} NEW ZEALAND PROCEDURE

K2O FE <-----> <-----> <-----> <-----> <-----> <----->

0.5 1.4

Page 4 of 4

83P2078 83P2079 3 83P2080 4 VFS 83P2081 5

HZN

NO.

FA

RE

GA45 OT33 GS14 GC 8

FD 1 NX 6

FD 1 NX 6

0.6 2.8

83P2082 6 83P2083 7

83P2077 1

CP83FN128

SAMPLE

NO.

(------) (------) (---DTA--) (TOT ANAL)

> NX 6 0.3 1.7

ANALYSES: S=ALL ON SIEVED < 2mm BASIS

MINERALOGY: FA = FRACTION ANALYZED RE = RESISTANT

KIND OF MINERAL: FD = FELDSPAR NX = NON-CRYSTALLINE GA = GLASS AGGREGATES GS = GLASS OT = OTHER

GC = GLASS COATED GRAIN

RELATIVE AMOUNT: 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

MINERALOGY BASED ON SAND/SILT:

MINERALOGY BASED ON CLAY:

FAMILY PLACEMENT:

COMMENTS:

Series: Ecuador No. 8 137 Date: 6/83

Pedon No: S83FN-325-008

Taxonomy: Coarse-loamy over cindery, mixed, isomesic Andic Humitropept

Latitude: 0°32'10" Longitude: 78°38'40"

Location: This pedon is located on the NE side of Corazon Volcano about 10 km W

of Machachi.

Physiography: Mountains Geomorphic Position: Hillslope

Slope and Aspect: 45% NE planar-convex Elevation: 3840 m M.S.L

Microrelief: smooth

Air Temp. Summer: Winter:

Precipitation: 1200 mm (there is not data) Udic moisture regime.

Water Table: > 10 m

Drainage: Well drained Permeability: moderate

Stoniness: nonstony

Land Use: Grassland or grazing land

Erosion or Deposition: none

Parent Material: multiple ash and pumice deposits

Described by: T. Cook, G. Del Posso, and Soil Survey Staff.

Remarks: Date described: January 31, 1983. Field Moisture, 0-184 cm wet.

All rock fragments are pumice.

Vegetation: stypa ichu (paramo pasture)

- Al 0 26 cm Black (10YR 2/1) loam; weak medium and coarse subangular blocky structure; very friable, slightly sticky and slightly plastic; weakly smeary, very moist or wet; many very fine and fine, few medium roots; many very fine and fine few medium tubular pores; 1% 2 to 5 mm fragments; diffuse wavy boundary. Worm casts make up about 35% of the horizon.
- A2 26 63 cm Black (10YR 2/1) loam; weak fine and medium subangular blocky structure; friable, slightly sticky and slightly plastic; moderately smeary, very moist or wet; common very fine and fine rocts; many very fine and fine tubular, pores; 1% 2 to 5 mm fragments; diffuse wavy boundary. Worm casts make up about 35% of the horizon.
- A3 63 92 cm Black (N 2/0) gravelly loam; weak medium and coarse subangular blocky structure parting to weak coarse and very coarse granular; very friable, slightly sticky and slightly plastic; weakly smeary, very moist or wet, common very fine and fine roots; many very fine and fine tubular, pores; coarse fragments, 20% 2 to 40 mm; abrupt wavy boundary. Worm casts make up about 35% of the horizon. Pumcie fragments and roots increase in amount in the lower part of the horizon.
- 92 134 cm Yellowish brown (10YR 5/4) and brownish yellow (10YR 6/6) very gravelly coarse sand; single grain; loose, non-sticky and non-plastic; few very fine and fine roots; many very fine, fine and medium interstital pores; coarse fragments, 45% 2 to 50 mm; abrupt wavy boundary. Dominate colors of pumice fragments, many other colors present.
- 3Ab 134 184 cm Black (10YR 2/1) loam; many fine and medium dark brown (7.5YR 3/2) mottles; weak coarse and very coarse angular blocky structure; very friable, slightly sticky and slightly plastic; weakly smeary, very moist or wet; few thin patchy clay skins or organic coatings on peds and common thin clay skins or organic coats in pores many very fine and fine common medium tubular pores. Worm casts make up about 35% of the horizon.

ECUADOR 008

LAB CLASSIF: COARSE-LOAMY OVER CINDERY, MIXED, ISOMESIC ANDIC HUMITROPEPT

S 83FN-325 -008 DATE 09/25/84 SAMPLE NO. 83P2210-2214

ECUADOR-SMSS

GENERAL METHODS 1B1A, 2A1, 2B

PEDON NO. 83P 475 PROJECT NO. 83P 83

U. S. DEPARTMENT OF AGRICULTURE
SOFT SOIL CONSERVATION SERVICE
NATIONAL SOIL SURVEY LABORATORY
LINCOLN, NEBRASKA 68508

PAGE 1 OF 4 PAGES

-1-- -2-- -3-- -4-- -5-- -6-- -7-- -8-- -9-- -10- -11- -12- -13- -14- -15- -16- -17- -18- -19- -20-

								-				• •	.,			.0	- 19-	-20-
SAMPLE NO.	HZN NO.		HORIZON		.05 LT	LT .002	.002	.02 05	. VI .05 10	F . 10 25	М .25 50	C .5 - 1	VC 1 -2	2 -5	WE 5 -20	1GHT - 20	.1-	PCT 0
****	EE F	OOTNOTE R	REGARDING	PSDA BELOW	***													
832210 832211 832212 832213 832214	1S 2S 3S 4S		A1 A2 A3 2C 3AB	2.3 31.7 1.7 35.5 1.1 28.6 0.2 4.4 2.0 49.5	66.0 62.8 70.3 95.4		18.0 20.8 16.5 2.7 32.2	14.7 12.1	14.5 13.5 14.4 1.4 16.0	15.6 12.1	15.9 16.7 37.8	8.8 10.4 15.2 28.2 4.9	4.0 8.4	TR 2 5 9 TR	TR TR 8 14	3V 3V 40V 54V	53 52 79 99 32	3 5 53 77
SAMPLE NO.	HZN NO.	6A1C 6B	AL EXTR P 3A PCT	FE 6R3A 6C2B	AL MN 6G7A 6D2	CEC A 8D1	リラ	- LIM LL 4F1	P1 4F	MOIST 4A3A	1/3 BAR 4A1D	OVEN DRY 4A1H	WHOLE SOIL 4D1	FIELD MOIST 4B4	1/10 BAR 4B1C	1/3 BAR	15 BAR 482A	WHOLE SOIL 4C1
832210 832211 832212 832213 832214	2 3 4	7.15 0.4 6.29 0.3 5.58 0.47 1.53		0.8 1.0 0.9 0.3 1.0	0.6 0.7 0.7 0.1 0.5	10.71 16.55 14.00	15.55				0.88 0.97 0.84	1.03 1.11 0.93	0.053 0.045 0.025			49.2 40.7 53.7	20.4 19.4	0.25 0.20 0.23

^{*} CAUTION SHOULD BE USED WITH PSDA DATA WHEN USED FOR MAKING ENGINEERING INTERPERTATIONS BECAUSE OF INCOMPLETE DISPERSION OF AMORPHOUS MATERIAL. THEY ARE PRESENTED TO AID IN CALSSIFICATION AND ASSESSMENT OF MINERALOGY. REFER TO PEDON DESCRIPTION FOR FIELD TEXTURE.

	ECU	ADOR 0	08 -2	-3	-4	S 83FN- -5	325 -0 -6	08 -7	DA -8	TE 09/ -9	'25/84 ~10~	-11-	PEDON -12-	NO. 83	3P 475 -14-	NA -15-	TIONAL	SOLL	SURVEY	2 OF 4 / LABOR -19-	ATORY
SAMPLE NO.	HZN	CA I 5B5A 6N2E	MG 585A	NA 5B5A 6P2B	K 5B5A 6Q2B	ASES -) SUM BASES -MEQ /	1 TY 6H5A	AL 6G9A	SUM CATS 5A3A	NH4- OAC 5A8B	BASES + AL 5A3B	SAT 5G1	SUM 5C3	NH4 OAC 5C1	CACO3 <2MM 6E1G	OHMS /CM 8E1		MMHOS /CM 81	NAF 8C1D		H20
832210 832211 832212 832213 832214	1 2 3 4 5	2.4 2.1 3.0 0.6 3.0	0.8 0.7 0.8 0.1 0.8	0.2		3.3 4.2 0.8		0.6 0.4	35.3 35.0 4.7	18.2	5.3 3.9 4.6 0.9	28 15 9 11 7	11 9 12 17 16	20 18 23 29 28					10.9 11.2 11.2 10.7 11.0	4.7 5.1 5.0 5.3 5.2	5.6 5.9 6.0 6.1 6.1
SAMPLE NO.	HZN	(NA C 6A4A	FE 6C8A	PHOSPH4 AL 6G10	TE EXT FE+AL - DI DI-CI	CRITERI FRACTAB FE+AL IVIDED PCT CLAY	LE) AL+C BY) PCT	INDEX OF)					(((7A21	X-I - X-I 7A21	- CLA RAY - 2U - 7A21	MINERAI Y)) 7A21 TS ->	7A3) A) 7A3	(+ TOTAL RES 7B1A) -) DOM WEATH 7B1A
832210 832211 832212 832213 832214	1 2 3 4 5		0.4 0.4 0.4 0.1 0.2	0.6 0.6 0.1	0.7 0.6 0.6 0.5 0.4	0.4 0.6 0.9 1.0 0.3		888 1275 999 193 1230						NX 6 NX 6						1	GA42
																					~

FAMILY CONTROL SECTION: DEPTH 25-100 PCT CLAY 1 PCT .1-75MM 67. SPODIC HORIZON: INDEX OF ACCUMUL 2928 ANALYSES: S= ALL ON SIEVED <2MM BASIS V= 75-20MM FROM VOLUME ESTIMATES MINERALOGY: KIND OF MINERAL NX AMORPHOUS GA GLASS AGG

RELATIVE AMOUNT 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

Page 3 of 4

ECUADOR 008 S83FN-325-008

N S S L

		EXT	15	P	(ACID	OXAL	ATE)	EXTR	рН
		P	BAR	RETENT	\mathtt{AL}	SI	FE	AL	KCl
		BRAYL	MOIST					(KOH)	
SAMPLE	HZN	6 S 3	4B2b	*	*	*	*		8Clg
NO.	NO.	PPM	<		-PCT OF <	2MM-		>	
832210	1		26.6	80	0.8	0.3	0.5	0.9	
832211	2		28.7	90	1.2	0.6	0.7	1.4	
832212	3		24.1	87	1.2	0.6	0.6	1.3	
832213	4		4.5	32	0.3	0.2	0.3	0.3	
032213	-								

SUPPLEMENTAL DATA SHEET

^{*} NEW ZEALAND PROCEDURE

Page 4 of 4

CP83FN128

						-							
		()	
SAMPLE NO.	HZN NO.	FA 7Bla	RE 7Bla <		781A	7B1A	7B1A	7Bla	7BlA	7BlA	7BlA	7BlA >	K2C PE A 7A2I 7A2I 7A2I 7A2I 7A3 7A3 6Q3A 6C7A > <relative amounts=""> <pct></pct></relative>
83P2210 83P2211 83P2212 83P2213	2	VFS	1	GA4'2	GS20	FP19	FG 7	PR 4	AR 4	FE 1	OG 1	н д 1	1 NX 6 0.3 2.1 NX 6 0.2 2.1
83P2214	5												NX 6 0.2 1.9

ANALYSES: S=ALL ON SIEVED < 2mm BASIS

MINERALOGY: FA = FRACTION ANALYZED RE = RESISTANT

KIND OF MINERAL: FE = IRON OXIDES AR = WEATHERED AGGREGATES FP = PLAGIOCLASE FELDSPAR GA = GLASS AGGREGATES GS = GLASS

PR = PYROXENE NX = NON-CRYSTALLINE HG = GLASS COATED HORNBLENDE FG = GLASS COATED FELDSPAR

OG = GLASS COATED OPAQUE

RELATIVE AMOUNT: 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

MINERALOGY BASED ON SAND/SILT:

MINERALOGY BASED ON CLAY:

FAMILY PLACEMENT:

COMMENTS:

14

Series: Ecuador No. 10 Pedon No: S83FN-325-010

Taxonomy: Sandy, mixed, isomesic Fluventic Hapludolls

Latitude: 0°33'00" Longitude: 78°37'00"

Location: This pedca is located about 2 km N of Aloasi or 5 km W of Machachi.

Physiography: Terrace

Geomorphic Position: Summit

Slope and Aspect: 3% E planar-planar

Elevation: 3000 m M.S.L

Date:

8/82

Microrelief: Smooth

Air Temp. Summer: Winter:

Precipitation: 1100 mm Udic moisture regime.

Water Table: > 10 m
Drainage: Well drained
Stoniness: none Permeability: moderate

Land Use: pasture mostly kikuyo Erosion or Deposition: none Parent Material: mixed ash deposits

Described by: T. Cook, G. Del Posso, and Soil Survey Staff.

Date Described: January 25, 1983. Field Moisture, 0-180 cm slightly moist. The upper 117 cm seems to be mixed by many animals or uprooted trees. Fifty meters from pit in another pasture the surface horizon was very dark brown (10YR 2/2) to a depth of 25 cm. Field estimates of the weighted average particle size control section places it in the sandy family. Vegetation: Pasture grasses and planted Ecucalyptus grove.

- 0 10 cm Very dark brown (10YR 2/2) and very dark grayish brown (10YR 3/2) sandy loam; weak fine subangular blocky structure; very friable, non-sticky and non-plastic; common very fine and fine and few medium and coarse roots; many very fine and fine interstital few very fine tubular, pores; 2% 2 to 5 mm fragments clear wavy boundary.
- C1 10 28 cm Dark grayish brown (10YR 4/2) and dark brown (10YR 3/3) sandy loam; massive; very friable, non-sticky and non-plastic; very weakly smeary, very moist or wet; few very fine, fine medium, and coarse roots; many very fine interstital, common very fine and fine tubular pores; 2% 2 to 5 mm fragments gradual wavy boundary.
- 28 40 cm Dark grayish brown (10YR 4/2) sandy loam; massive; very friable, non-sticky and non-plastic; very weakly smeary, very moist or wet; few very fine, medium, fine, and coarse roots; many very fine interstital common very fine and fine tubular pores; 2% 2 to 5 mm fragments; clear irregular boundary.
- 40 53 cm Very dark gray (10YR 3/1) and dark grayish brown (10YR 4/2) coarse sand; single grain; loose, non-sticky and non-plastic; few very fine, medium, fine and coarse roots, many very fine interstital, few very fine tubular pores; 2% 2 to 5 mm fragments; clear irregular boundary.
- 53 85 cm Black (10YR 2/1, rubbed) and very dark gray (10YR 3/1, rubbed); loamy sand; weak fine and medium subangular blocky structure; very friable, non-sticky and non-plastic; few very fine, fine, medium, and coarse roots; many very fine interstital, few very fine tubular pores; 7% 2 to 10 mm fragments; diffuse wavy boundary. 15 to 20% of horizon is very dark gray (10YR) 3/1, moist) pockets 10 to 15 cm in size.
- 85-117 cm Black (10YR 2/1, rubbed) and very dark gray (10YR 3/1rubbed) sandy loam; weak fine and medium subangular blocky structure; friable, non-sticky and non-plastic; few very fine, fine, medium, and coarse roots; many very fine interstital, few very fine tubular pores; coarse fragments, 10% 2 to 20 nmm; gradual wavy boundary. 15 to 20% of horizon is very dark gray (10YR 3/1, moist) pockets 10 to 15 cm in size.
- 2Ab3 117 180 cm Dark reddish brown (5YR 2/2) loam; weak rine and medium subangular blocky structure; very friable, slightly sticky and slightly plastic; weakly smeary, very moist or wet; few very fine, fine and coarse roots; few very fine interstital, common very fine and fine tubular pores.

ECUADOR 010

ECUADOR-SMSS

LAB CLASSIF: SANDY, MIXED, ISOMESIC FLUVENTIC HAPLUDOLL

S 83FN-325 -010

DATE 09/25/84

SAMPLE NO. 83P2220-2226 PEDON NO. 83P 47?

PROJECT NO. 83P 83

11.00201 110.

U. S. DEPARIMENT OF AGRICULTURE SOIL CONSERVATION SERVICE NATIONAL SOIL SURVEY LABORATORY LINCOLN, NEBRASKA 68508

PAGE 1 OF 4 PAGES

GENERAL METHODS 181A, 2A1, 2B

		-1	-2	-3	-4	-5~-	-6	~7	-8	-9	-10-	-11-	- 12-	-13-	-14-	-15-	-16-	-17-	-18-	-19-	-50-
SAMPLE NO.	HZN NO.	DEPTI (CM)		ORIZON	LT	.002	.05	LT .0002	LT .002	.002	.02 05	. VF .05 10	. 10 25	м .25 50	.5 -1	VC 1	2	WE 5	ACTIONS IGHT - 20 -75 <75MM(3	.1-	PC1 O
**** S	EE FO	оотиот	E REGA	ARDING	PSDA 8	ELOW *	****														
832220 832221 832222 832223 832224 832225 832226	18 28 38 48 58 68	0- 1 10- 2 28- 4 40- 5	10 28 40 53 35 2	A C1 C2 C3 2AB1 2AB2 3AB3	2.7 2.3 2.5 1.0 1.3 2.1 3.5	25.5 24.7 27.1 15.1 20.5 27.0 42.5				14.1 14.0 17.1 8.2 10.7 14.5 26.5	11.5 10.7 10.1 6.9 9.8 12.5 16.0	15.2 16.8 15.1 11.6 15.9 17.5 16.0	24.3 21.7 22.4 22.5 18.3	17.2 25.5 18.7 13.5	11.9 10.3 11.3 16.3 14.9 13.4 6.1	5.5 4.2 5.1 8.1 6.2 8.2 2.9	4 3 4 5 6 7 3	1 1 3 4 1	4V 4V 4V 4V 14V 20V 7V	61 60 59 75 71 68 45	9 8 9 10 23 31
SAMPLE NO.	HZN NO.	ORGN 1 C 6A1C <	N 6B3A	•	6R3A	FE 6C2B	AL 6G7A	MN 6D2A	CEC	/CLAY) 15 BAR 8D1	- LIM LL 4F1	BERG) ITS - PI 4F 0.4MM	MOIST 4A3A	7 1/3 F BAR 4A1D	OVEN DRY 4A1H	WHOLE SOIL 4D1	FIELD MOIST 484	1/10 BAR 431C	CONTEN 1/3 BAR 4B1C OF <2MM	15 BAR 482A	WHOLE SOIL UC1
832220 832221 832222 832223	2 3 4	2.87 0 0.85 0 0.47 0 0.42	0.072			0.3 0.3 0.3	0.1 0.1 0.1 0.1		4.22 2.13 1.68 3.70	3.26 1.91 1.52 3.50				1.18					28.7	8.8 4.4 3.8	0.28
832224 832225 832226	6	0.77 1.04 0.93				0.3 0.4 0.4	0.1 0.1 0.1		4.31 3.62 2.43	3.08 2.19 1.49				1.10	1.12	0.007 0.005 0.009			25.9 32.6 28.8	3.5 4.0 4.6 5.2	0.23 0.26 0.25

^{*} CAUTION SHOULD BE USED WITH PSDA DATA WHEN USED FOR MAKING ENGINEERING INTERPERTATIONS BECAUSE OF INCOMPLETE DISPERSION OF AMORPHOUS MATERIAL. THEY ARE PRESENTED TO AID IN CLASSIFICATION AND ASSESSMENT OF MINERALOGY. REFER TO PEDON DESCRIPTION FOR FIELD TEXTURE.

	ECU	ADOR C	10		:	S 83FN-	325 -0	10	DA	TE 09	/25/84		PEDON	NO A	33P_477	N:A	TIONAL			OF 4 1	
		-1	-2	-3	-4	-5	-6	-7	-8	-9	-10-	-11-	-12-	-13-	-14-	-15-	-16-	-17-	-18-	-19-	-80-
											· 								~		
SAMPLE NO.	HZN	5B5A 6N2E	5B5A 602D	5B5A 6P2B	5B5A 6Q2B	BASES	6H5A	AL 6G9A	SUM CATS 5A3A	NH4- OAC 5A8B	BASES + AL 5A3R	AL SAT 5G1	SUM	NH4 OAC	CO3 AS CACO3 <2MM	OHMS /CM		MMHOS /CM	NAF	-PH - CACL2 .01M	H20
		<				-MEQ /	100 G				>		P	CT	6E1G	OLI		81	8010	8C1F 1:2	8G1F 1:1
832220 832221 832222 832223 832224 832225 832226	1 2 3 4 5 6 7	8.0 2.8 2.4 1.7 3.3 5.4 6.5	1.8 0.8 0.9 1.0 1.3 1.0	0.1 0.1 0.1 0.2 0.2 0.2	0.5 0.3 0.3 0.2 0.3 0.2	10.4 4.0 3.7 3.1 5.1 6.8 8.5	6.0 2.4 1.7 1.9 2.0 3.9 3.4	TR TR 0.1 0.1 TR 0.1 0.1	16.4 6.4 5.4 5.0 7.1 10.7	11.4 4.9 4.2 3.7 5.6 7.6 8.5	3.8 3.2 6.9 8.6	3 3	63 62 69 62 72 64 71	91 82 88 84 91 89					9.6 9.9 9.7 9.8 9.9	5.9 6.0 6.1 5.9 6.0 6.1	6.0 6.2 6.6 6.6 6.6
								0.1	,	0.7	0.0	•	,,	100					9.7	6.5	6.9
SAMPLE NO.	HZN	(NA C 6A4A	PYROF FE 6C8A	PHOSPHA AL 6G10	TE EXT FE+AL (Di Di-Ci	RITERI. RACTAB FE+AL VIDED PCT CLAY	LE) AL+C BY) PCT	INDEX OF)				(7A21	 - X-F X 7A21 LATIVE	- CLA RAY - RAY - RAZI	Y)() 7A21	DT) A) 7A3	TOTAL RES 7B1A	DOM WEATH 7B1A
832220 832221 832222 832223 832224 832225 832226	1 2 3 4 5 6 7		0.1	0.1 0.1 0.1 0.1 0.1 0.1	0.5 0.3 0.3 0.3 0.2 0.2	0.1 TR TR 0.1 0.1 TR TR		150						NX 6 NX 6 NX 6							
																			.~		

FAMILY CONTROL SECTION: DEPTH 25-100 PCT CLAY 2 PCT .1-75MM 68. SPODIC HORIZON: INDEX OF ACCUMUL O ANALYSES: S= ALL ON SIEVED <2MM BASIS V= 75-20MM FROM VOLUME ESTIMATES MINERALOGY: KIND OF MINERAL NX AMORPHOUS

RELATIVE AMOUNT 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

		EXT P	15 BAR	P RETENT	(ACII AL	OXAL	ATE) FE	EXTR AL	pH KCl
		BRAYl	MOIST			~ -		(KOH)	NCI
SAMPLE	HZN	6S3	4B2b	*	*	*	*	(KOn)	901~
NO.	NO.	PPM	(-PCT OF	<2MM			8Clg
						\ZIMI		>	
832220	1		7.5	20	0.2	0.1	0.2	0.1	
832221	2		3.8	14	0.2	0.1	0.2	0.1	
832222	3		3.8	14	0.1	0.1	0.2	0.1	
832223	4		3.3	14	0.1	0.2	0.2	= :	
832224	5		4.5	17	0.3			0.1	
832225	6					0.2	0.2	0.2	
	0		5.9	20	0.3	0.3	0.2	0.2	
832226	/		7.0	17	0.4	0.3	0.2	0.1	
									_•

^{*} NEW ZEALAND PROCEDURE

CP83FN128

																	
		((·			(OPTICAL ND/SIL	, ,T			MINERA)	((DT	 .A)	(TOT) ANAL)
SAMPLE NO.	HZN NO.	FA 781A	RE 7BlA <	7Bla	7Bla	7BlA	7Bla		7Bla	7Bla	7nla	7RIA	(CLAY- 7A2I 7A2I 7A2I 7A2I 7A2I <relative amounts=""></relative>	7A3	7A3	K20 6Q3A	PE 6C7A
83P2220	1																
83P2221 83F2222	2	VFS		GA40	ОТ29	GS17	GC15						NX 6			0.7	1.6
83P2223 83P2224	4 5												NX 6			0.4	0.0
83P2225	6															0.4	8.0
83P2226	7												NX 6			0.3	1.6

ANALYSES: S=ALL ON SIEVED < 2mm BASIS

MINERALOGY: FA = FRACTION ANALYZED RE = RESISTANT

KIND OF MINERAL: GA = GLASS AGGREGATES GS = GLASS OT = OTHER NX = NON-CRYSTALLINE GC = GLASS COATED GRAIN

RELATIVE AMOUNT: 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

MINERALOGY BASED ON SAND/SILT:

MINERALOGY BASED ON CLAY:

FAMILY PLACEMENT:

COMMENTS:

<u>_</u>

Series: Ecuador No. 11 Pedon No: S83FN-325-011

Date: 8/82

Taxonomy: Coarse-loamy over ashy (pumice), mixed, isohyperthermic Andic Humitropept

Latitude: 0°25'00" Longitude: 79°02'20"

Location: This pedon is located 20 km east of Alluriquin near Tandapi.

Physiography: mountains

Geomorphic Position: Hillslope

Slope and Aspect: 45% NE planar-convex Elevation: 1500 m M.S.L.

M.S.L.
Microrelief: Complex.
Summer: Winter:

Precipitation: 2100 mm perudic moisture regime.

Water Table: > 10 m

Drainage: Well drained Permeability: moderate rapid in upper part.

moderate in lower part. Stoniness: nonstony

Land Use: pasture.

Erosion or Deposition: none apparent.

Parent Material: multiple coarse and fine ash and pumice deposits. Described by: T. Cook, G. Del Posso, and Soil Survey Staff.

Date Described: January 27, 1983. Field Moisture, 0-202 cm wet. Vegetation: Native vegetation was subtropical mixed forest, now cleared and planted to pasture grasses.

- Al $0-25~{\rm cm}$ Black (10YR 2/1) loam; moderate medium and coarse subangular blocky structure parting to moderate medium and coarse granular; friable, non-sticky and non-plastic; many very fine, common fine, few coarse roots; many very fine and few fine tublar, common very fine interstitial pores; 1% 2 to 5 mm pumice fragments; gradual wavy boundary. About 1% fragments of brick, stones, artifacts, and charcoal.
- A2 25 57 cm Black (10YR 2/1) loam; weak medium and coarse subangular blocky structure parting to weak coarse granular; friable, non-sticky and non-plastic; weakly smeary, very moist or wet; common very fine and fine roots; many very fine and common fine tubular pores; 1% 2 to 5 cm pumice fragments; diffuse wavy boundary. About 1% fragments of brick, stone, artifacts, and charcoal.
- Black (N 2/0) sandy loam; weak coarse subangular A 3 57 - 84 cm blocky structure; friable, non-sticky and non-plastic; a few thin patchy organic coatings on peds; few very fine and fine roots; many very fine, and few fine tubular pores; 7% 2 to 15 mm pumice fragments; clear wavy boundary.
- 2Cl 84 120 cm Brown to dark brown (10YR 4/3) gravelly coarse sand; single grain; loose, non-sticky and non-plastic; few very fine roots; many very fine interstital pores; 25% 2 to 10 mm pumice fragments; clear wavy boundary. The upper few centimeters of the horizon has been stained black (10YR 2/1).
- 3Abl 120 133 cm Very dark grayish brown (10YR 3/2) loam; weak coarse angular block structure; friable, non-sticky and non-plastic; moderately smeary, very moist or wet; few very fine roots; many very fine, few fine tubular, and common very fine interstital pores; 1% 2 to 10 mm pumice fragments; diffuse wavy boundary.
- 133 156 cm Very dark grayish brown (2.5Y 3/2) loam; weak coarse angular blocky structure; very friable, non-sticky and non-plastic; moderately smeary, very moist or wet; a few very fine roots; many very fine, few fine tubular and many very fine interstital pores; 1% 2 to 5 mm pumice fragments; clear wavy boundary.
- Very dark brown (10YR 2/2) sandy loam; weak medium 3Ab3 156 - 178 cm subangular blocky structure; very friable, non-sticky and non-plastic; few very fine roots; many very fine, fine tubular and many very fine interstital pores; 7% 2 to 15 mm pumice fragments; clear wavy boundary.
- Yellowish brown (10YR 5/6) gravelly coarse sand; 178 - 202 cm single grain; loose, non-sticky and non-plastic; few very fine roots; many very fine and fine interstitial pores; 35% 2 to 30 mm pumice fragments.

PAGE 1 OF 4 PAGES

U. S. DEPARTMENT OF AGRICULTURE

NATIONAL SOIL SURVEY LABORATORY

SOIL CONSERVATION SERVICE

LINCOLN, NEBRASKA 68508

ECUADOR 011

LAB CLASSIF: COARSE-LOAMY OVER ASHY (PUMICE), MIXED, ISOHYPERTHERMIC ANDIC HUMITROPEPT

S 83FN-325 -011 DATE 09/25/84 SAMPLE NO. 83P2135-2142

ECUADOR-SMSS PROJECT NO. 83P 463
PROJECT NO. 83P 83

GENERAL METHODS 1B1A, 2A1, 2B

-1-- -2-- -3-- -4-- -5-- -6-- -7-- -8-- -9-- -10- -11- -12- -13- -14- -15- -16- -17- -18- -19- -20-

SAMPLE NO.		DEPTH (CM)	HORIZON	(CLAY LT .002 <	.002	.05	LT	LT .002	(SI FINE .002 02 OF <2M	.02 05	. Vf .05 10	. 10	.25 50	C 5	VC 1 -2	2 -5	WE 5 -20	ACTIONS IGHT - 20 -75 <75MM(3	 .1- 75	WT PCI O WHOLE
**** S	EE F001	TNOTE RE	GARDING	PSDA B	ELOW #	***														
832135 832136 832137 832138 832139 832140 832141 832142	15 25 2 35 5 45 8 55 12 65 13 75 15	0- 25 25- 57 57- 84 34-120 20-133 33-156 56-178 78-202	A1 A2 A3 2C1 3AB1 3AB2 3AB3 4C2	3.9 4.6 3.0 1.5 2.7 2.7 2.5 1.2		49.6 47.9 62.6 95.5 63.7 60.4 81.4 94.7			30.5 31.5 21.2 2.4 21.7 24.3 10.7 3.1	16.0 16.0 13.2 0.6 11.9 12.6 5.4	15.3 16.4 13.1 1.1 14.4 16.0 6.2 2.9	19.9 19.2 20.2 15.7 27.8 20.4 15.0	11.8 9.3 18.1 37.9 17.6 16.6 31.6 32.7	2.4 2.5 8.3 27.3 3.2 5.0 21.1 33.6	0.2 0.5 2.9 13.5 0.7 2.4 7.5 14.3	TR TR 2 6 TR 1 3 5	TR TR 1 2 TR TR 2 6	3V 3V 16V 36V 2V 2V 15V 47V	36 34 59 97 50 46 80 97	3 3 19 44 2 3 20 58
SAMPLE NO.	HZN NO. 6A	RGN TOTAL	١	6R3A	(D EX FE 6C2B MM	AL 6G7A	WN SCE	CEC 8D1	/CLAY) 15 BAR 8D1	- LIM LL 4F1	BERG) ITS - PI 4F 0.4MM	MOIST 4A3A	1/3 BAR	OVEN DRY 4A1H	WHOLE SOIL 4D1	FIELD MOIST 4B4	1/10 BAR 4B1C	CONTENT 1/3 BAR 4B1C DF <2MM	15 BAR 4B2A	WHOLE SOIL 4C1
832135 832136 832137 832138 832139 832140 832141 832142	2 3. 3 3. 4 0. 5 1. 6 0. 7 0.	08	7		0.7 0.8 1.0 0.3 0.5 0.5 0.6 0.4	0.6 0.5 0.6 0.1 0.2 0.2 0.2		6.97 3.30 4.90 1.40 2.37 2.33 2.56 1.67	5.95 2.52 4.07 2.33 2.56 2.33 2.16 2.17				1.02		0.016 0.021 0.012 0.006			72.8 42.5 45.8 31.1 29.0 26.4	23.2 11.6 12.2 3.5 6.9 6.3 5.4 2.6	0.31

^{*} CAUTION SHOULD BE USED WITH PSDA DATA WHEN USED FOR MAKING ENGINEERING INTERPERTATIONS BECAUSE OF INCOMPLETE DISPERSION OF AMORPHOUS MATERIAL. THEY ARE PRESENTED TO AID IN CLASSIFICATION AND ASSESSMENT OF MINERALOGY. REFER TO PEDON DESCRIPTION FOR FIELD TEXTURE.

	ECUAD	OOR 01	1 -2	-3	-4	83FN- -5	325 -0 -6	11 -7	DA -8	TE 09/ -9	/25/84 -10-	-11-	PEDON -12-	NO. 8	33P 463 -14-	NA -15-	T10HAL -16-	COLL	CHOVES	2 OF 4 7 LABOR -19-	A FANOL
SAMPLE NO	HZN 5	NH404 CA B5A 5 N2E 6	พน 5B5A 502 <i>ม</i>	5B5A 6P2B	к 5В5А 6Q2В	BASES	6H5A	AL 6G9A	SUM CAIS 5A3A	NH4+ OAC 5A8B	BASES + AL 5A3B	5G1		N114 OAC 5C1	CACO3 <2MM 6E1G	OHMS /CM		MMHOS /CM	NAF	- PH - CACL2 .01M 8C1F 1:2	H20'
832135 832136 832137 832138 832139 832140 832141 832142	2 3 4 5 6 7	0.6 1.7 1.6 2.0	1.9 0.3 0.2 0.1 0.1 0.2 0.2	0.3 0.2 0.3 0.3 0.3 0.3	0.6 0.2 0.1		27.7 24.5 25.7 2.6 11.5 10.5 9.9 2.6	0.1 0.4 0.1 0.1 0.1 0.1 TR	43.7 28.0 29.5 3.6 13.6 12.6 12.4 3.6	27.2 15.2 14.7 2.1 6.4 6.3 6.4 2.0	16.1 3.9 3.9 2.2 2.2 2.6	1 10 3 5 5 4	37 12 13 28 15 17 20 28	59 23 26 48 33 33 39 50					10.8 11.1 11.0 10.1 10.8 10.7 10.7	5.5 5.4 5.6 5.5 5.8 5.9 6.0 5.8	5.8 5.8 6.1 6.2 6.2 6.1 6.3
SAMPLE NO.	(- HZN NO. 6	-NA P C A4A 6	PYROP FE SC8A	HOSPHA AL 6G10	TE EXT FE+AL (D! DI-C!	RITERI. RACTAB FE+AL VIDED PCT CLAY	LE) AL+C BY) PCT	INDEX)				(7A21	- X-I - X-I 7A21 LATIVE	- CLAY RAY 2U 7A2I	()) 7A21	7A3) A) 7A3	TOTAL RES 7B1A	DOM WEATH 7B1A
832135 832136 832137 832138 832139 832140 832141 832142	1 2 3 4 5 6 7 8		0.4	0.6 0.6 0.5 0.1 0.2 0.2 0.2	0.8 0.6 0.4 0.3 0.3 0.3	0.3 0.2 0.2 0.1 0.1 0.1 0.1		1044 822 756 159 259 245						VR 1 NX 6 NX 6							

FAMILY CONTROL SECTION: DEPTH 25-100 PCT CLAY 3 PCT .1-75MM 56.
ANALYSES: S= ALL ON SIEVED <2MM BASIS V= 75-20MM FROM VOLUME ESTIMATES
MINERALOGY: KIND OF MINERAL VR VERMICULITE NX AMORPHOUS CL CHLORITE

RELATIVE AMOUNT 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

Page 3 of 4

NSSL SUPPLEMENTAL DATA SHEET

CAMPIE		EXT P BRAY1	15 BAR MOIST	P RETENT	AL	D OXAL.	ATE) FE	EXTR AL (KOH)	pH KCl
SAMPLE NO.	HZN NO.	6S3 PPM	4B2b <		-PCT OF	* <2MM-	* 	>	8Clg
832135	1		31.0	78	0.8	0.3	0.5	 0.8	
832136	2		21.5	83	1.1	0.5	0.4	1.2	
832137	3		22.9	89	1.7	0.8	0.6	1.5	
832138	4		5.0	27	0.4	0.3	0.4	0.3	
832139	5		13.3	70	1.0	0.6	0.2	1.0	
832140	6		12.1	66	1.1	0.6	0.3	1.0	
832141	7		9.8	52	0.8	0.5	0.2	1.0	
832142	8		5.3	40	0.3	0.2	0.2	0.4	

^{*} NEW ZEALAND PROCEDURE

Page 4 of 4

CP83FN128

		(MINERA	LOGY)
		(-	PTICAL ND/SIL)	(-X-RA	Y)	(1	OTA)	(TOT	ANAL)
SAMPLE NO.	HZN NO.	FA 7Bla	RE 7Bla <	7BlA								7A2I 7					7A3 <	7A3 PCT	K20 6Q3A	
83P2135	1								 											
83P2136	_											VR 1 N	NX 6						0.2	1.7
83P2137	_	VFS		GA49	GC21	OT20	GS10					NX 6							0.2	2.0
83P2138	_																			
83P2139	_	VFS		GA48	OT31	GC17	GS 5					NX 6							0.0	1.0
83P2140 83P2141	_																			
83P2141	-	VFS		GS55	GC 28	OT10	GA 7	-				CL 1 N	NX 6						0.1	0.7

ANALYSES: S=ALL ON SIEVED < 2mm BASIS

MINERALOGY: FA = FRACTION ANALYZED RE = RESISTANT

KIND OF MINERAL: VR = VERMICULITE NX = NON-CRYSTALLINE GA = GLASS AGGREGATES GS = GLASS OT = OTHER

GC = GLASS COATED GRAIN CL = CHLORITE

RELATIVE AMOUNT: 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 MODERATE 2 SMALL 1 TRACE

MINERALOGY BASED ON SAND/SILT:

MINERALOGY BASED ON CLAY:

FAMILY PLACEMENT:

COMMENTS:

15

Appendix II

Summary of micromorphological descriptions of soils for the Sixth International Soil Classification Workshop

R. Honorato

Facultad de Agronomía
Pontificia Universidad Católica de Chile

PERM	SAMPLE	RELATED DIST	RIBUTION PATTERN	SOIL PABRIC	1	PEDOLOGICAL P	EATURES		ORGANIC COMPOUND	zś.		
		Normal	Specific		Cutans	Sesq.Glab.	Pedotubules	Org.skeleton		Coloidal mull		N GRAINS
98 7	2081	rosphysic	Agglutinic(E) Porphyroakelic(B)	Isotic	-	××	-	-	23		plagioclases (A)	Degree of weathering
											volcanic rock frag.(8) amphpyroxenes(5)	nearly frash/complex alteratio some pyroxenas moderately altered pullicular
649	2093	Porphyric	Agglcongelic(E) Porphyroskelic(B)	Isotic	amorphous coatings	XX	××	x	x	**	punics withic frag.(C-) (coated and uncoated) plagicclases(A) wolcanic rock frag.(S) pyroxenes(S) quarts (T)	 nearly fresh some plagicclases sligthly sltered
909	2095	Porphyric	Agglutinic(E) Porphyroskelic(B)	Isotic	-	**	¥	-	***	-	plagioclases(A) volcanic rock frag.(S) pyroxenes(S) pumics vitric frag.(S-C quartz (T)	nearly fresh
001	2096	Porphyric Porphyri-	Agglutinic(E) . Porphyroskelic(B)	Isotic	-	**	-	-	×	-	plagioclases(A) volcanic rock frag.(S) pumics vitric frag.(S-C amphpyroxenss(S) micas (T)	nearly fresh
001	2108	granic Porphyric	Agglcongelic Porphyroakelic(B)	Isotic	amorphous coatings	X.F	ж	-	XXX	XX	pusice vitric frag.(A) (coated) amphpyroxenes(C) plagioclases(C) wolcanic rock frag.(S) micas (S)	neerly fresh
004	2120	Porphyri-	Agglcongelic Porphyroskelic(8) very loose congelic	Isotic	amorphous coatings	ж	×	-	XX	*	plagioclassa(A) pumics vitric frag.(C) amphpyroxenes(S) volcanic rock frag.(S) micas (T)	nearly fresh to sligthly alteration complex of plagioclases
204	7124	granic Porphyric	to dermatic		coatings	XX	•	-	*	**	pumice vitric frag.(A) (coated and uncoated) plagicclasss(A) volcanic rock frag.(S) amphpyroxnes(S) micas (S)	nearly fresh some amphpyroxenes strongly sltered
011	2117		Agglutinic(E) Porphyroskalic(B)	Isatic	-	XX	•	-	x	•	plegioclases(C) pumice vitric frag.(S) amphpyroxenes(S)	nearly fresh
		Phyri-granic	Agglcongelic Porphyroskelic(B)	Isotic	some amorphous coatings	•	•	•		XX.	plagioclases (A) punice vitric frag.(C) amphpyroxenes(S) micas [T]	nearly fresh
005	2152	Porphyric	Agglcongelic Porphyroskelic(B)	Isotic	few amorphous coatings	XX	ж	××	KX	xx	plagioclases (C-A)	nearly fresh
005	2154	Parphyric	Int. congelic-aggl, Porphyroskelic(B)	Isotic	xx amorphous	××	==	=		XX.	plagioclassa(C-A)	some amphpyroxenes strongly sitered nearly fresh to sligthly fresh some amphpyroxenes
00 \$	2155	Porphyric	Int. agglcongelic	Isotic	xx	××		_			pumice vitric frag.(S) (Coated) Lices (T)	strongly altered
			Porphyroskelic(B)					*	EX.		plagioclases(C) smphpyroxenes(T) pumice vitric frag.(S) [coated]	nearly frush

PZCCŃ	SAMPLE	RELATED DIST	RIBUTION PATTERN	SOIL PABRIC	P	EDOLOGICAL F	EATURES	·	ORGANIC COMPOUND			
		Normal	Specific		Cutana	Seaq.Gleb.	Pedotubules	Org.skeleton	Coloidal humus	Coloidal mull	EXELETAN Hineralogy	Degree of weathering
005	2157	Porphyric	Int. congelic-aggl. Porphyroskelic(8)	. Isotic	x	23	×	-	11	K.X	plagioclases(S-C) micas (T)	eligthly altered (complex alteration)
012	2168	Porphyric	Int. agglcomg.(z) Porphyroskalic(B)	Isotic	m (amorphous)	XX	жx	-	X.X	-	plsgioclases(C) pumice vitric frag.(8) amphpyroxenes(S)	sligthly sitered (complex siteration)
012	2170	Porphyric	Agglutinic(E) Porphyroskelic(B)	Isotic	*	XX	-	-	**	-	plagicclasss(C) volcanic rock frag.(S)	sligthly altered placioclases fractented
012	2172	Plasmi- porphyric	Agglutinic(E) Porphyroskelic(B)	Isotic	XX	KX	-	-	×	-	plagioclases (A)	sligthly altered plagioclasss fragmented
013	2103	Porphyric	Agglutinic(E) Porphyroskelic(B)	Isotic	×	**	*	-	×	-	pumice vitric frag.(S) plagioclases (A) quartz (T)	sligthly altered coins of planicclases
-	2184	Pleamic	Agglutinic(E) Porphyroekelic(B)	Isotic	XX (ferrans)	XX	-	-	×	-	scarce skeleton quartz (T)	
013	2185	Plasmic	Agglutinic(E) Porphyroakalic(B)	Isotic	xx (ferrene)	EXX	-	-	-	-	no grains	
002	2192	Phyri-granic	Dermatic '	Isotic	-	-	-	-	×	-	to thick in some coarse grains plagioclases are coated (A)	nearly fresh
002	2193	Phyri-granic	Dermatic	Imotic	7	-	- ,	-	×	-	volcanic rock frag.(\$) (glassy) plagioclases(C)	nearly fresh
002	2194	Mainly volcan	ic material with a fi	ne volcanic p	paste as mat:	rim including	phenocryste o	f feldesphats a	ind pyroxenes prin	cipaly.		
003	2201	Porphyric	Int.congaggl.(E) Porphyroskelic(B)	Isotic	-	K.S	×		жж	x	plagioclases(C) pumice vitric frag.(C)	plagioclease and pumice slightly fresh to atrongly altered
003	2202	Porphyric	Int.agglcong.(E) Porphyroskalic(B)	Isotic	-	xx .	×	-	X.F	*	plagioclases(C) pumice vitric freg.(C)	plagioclases and pumice aligthly fresh to strongly altered
003	2204	Porphyric	Agglutinic(E) Porphyroskelic(B)	fectio	-	7	-	-	*	-	pleqioclases (C) punics vitric frag. (C) amphpyroxenes (S)	plagioclases and pumice miigthly fresh to strongly altered
008	2211	Porphyric	<pre>Int.agglcong. Porphyroskelic(B)</pre>	Isotic	amorphous costings	EE	-	xx	xx	*	plagioclases (C-A) pumice vitric frag. (C-A) (coated and uncoated)	plagioclases and pumice sligthly fresh to strongly altered
000	2212	Parphyric	Int.agglcong. Porphyroskelic(B)	Isctic	amorphous coatings new purice	**	*	*	xx	*	plagicclases(C-A) purics vitric frag.(A) (coated and uncoated)	plagiocisses and pumice slightly fresh to strongly altered (scmething less)
000	2214	Porphyric	Int.egglcong. Porphyroskelic(B)	Isotic	amorphous coatings new pumice	XX.	×	-	ż	-	plagioclases (C-A) pumice witric frag.(A) (costed and uncosted)	plegiocisses and purite slightly fresh to errongly altered (something less)
010	2221	Granic	Dermetic(E)	Isotic	some smorphous costings	×	×	x	××	-	plegioclases/A- purice vitric fra: A)	alightly altered

Sey:

Nomenclature and criteria:

- Abondant xxx A
- Common xx C
- Scarce x S
- Traces T

Weathering Degree

G. Stoops, E. Risdom, J.Delvigne, E. Fitzpatrick and G. Paneque. 1978. Sub-group on weathering phenomena and neofromations a progress report.

Soil Fabric

Soil Fabric and Fedological Features

E = Eswaran B = Brewer

H. Eswaran and C. Baños. 1976. Related distribution patterns in soils and their significance, An. Edaf. y Agrb.

R. Brewer.1964. Fabric and mineral analysis of soils. John Wiley Inc.


Appendix III

Soil Moisture and Temperature Regimes

The following five pages show the determination of soil moisture and temperature regimes according to Franklin Newhall system of computation for selected stations in the Ecuador (Newhall, 1972). This information was taken from SMSS Technical Monograph No. 2, 1981 (Van Wambeke, 1981).

158

ECUADOR



SYMBOL TABLE:

USTIC --- S
UDIC --- D
APIDIC --- A
XERIC --- X
PERUDIC -- P

	DETERMIN	IATION	OF SOIL MOIS	THEE	a e c	INE .	.ccnp	O TAKE	Tn s	EDA	MELTN I	NEW	PALL	SVET	EM 0E	COMBI	1 T A T	LON		
FOR ECUA	DETERMEN	1211011	or sore acra	IONE	K L G	T 14 C 1	4CCUR	OTING	10 1	- N M	HVETH I	14C 71	ואננ	31311	cm ur	COMPL	JIAI	LUN		PAGE 1
- 李安宗李宗宗李宗宗李宗 (********	*****	*******	** = *:	****	***	****	**	* * * * 1	. * *	****	**	***	****	****	*****	***	****	***	*******
NAME OF	* MEAN SO) [+	TEMPERATURE	* C	UMUL	IV I TA	E DAY	S MS	C	*	HAX.	.co	NSEC	JTIVE	DAYS	THAT	MOS	15	*	MGISTURE *
STATION	* TEMPERAT	URE +	REGIME	*	ΙN		*	NHEN		2 1	HCIST					OR Y	*	MCIST		
*		FINT \$		#ONE	YEAR	2 I S	*50I	L TE	MP >5	5.*	IN ONE		* 51	EN SO	{ L * *	AFTER	*	AFTER	*	REGIME +
*	‡ \$	* *		*DRY	H/D	MGI	* DRY	H/D	IOM	*	YEAR		*TE	MP > 1	ອ **	SUMMER	₹ \$	WINTER	*	
4	* * *			*			*			*			*					SOLST.		*
- 中华农林州市中华海外市市。	****	****	*******	****	***	***	***	****	***	**	*****		* * *	****	* * * *	****	***			******
*AMBATO	*16.4 16.7	15.6#	ISOTHERMIC	* 245	115	0	\$245	:15	0	*	86		*	86	**	75	*	0	*	ARIDIC +
≠ANCON	*26.1 27.0	24.8*	ISOHYPERTH.	*328	32	0	+328	32	0	\$	18		ŧ	18	* *	45	*	-	*	ARIDIC *
*ANGEL (EL)	*14.2 13.5	14.3*	ISOMESIC	* 0	0	360	* 0	0	360	*	360		*	360	**	0	*	120		upic +
*ARGELIA (LA	1*17.5 17.e	17.0#	ISOTHERMIC	⇒ 0	0	360	* C	ō	360		360		*	360	¢ ‡	ō	*	120		UDIC +
*ASCAZUBI			ISOTHERMIC			360		0	360	*	360		*	360	**	ŏ	*	120		UDIC +
*BABAHOYO			ISOHYPERTH.		5.5	178	#127		176		219		*	219	**	13	ø	13		# USTIC
*BAHIA DE CAR	R*27.2 27.1	26.5*	I SOHYFER TH.	\$23G	40	81	¥239		81		98		*	98	**	15	*		ŧ	USTIC *
⇒BALZAR			ISOHYPERTH.				+119				229		*	229	*	14	€.	32		USTIC *
*BANOS			ISOTHERMIC				~ O	O	300		360		3	360	**	Ô	*	120		UDIC *
≎CALDERON			ISOTHERM!C		-		*141	219	٥		157		*	167	± e	38	×		÷	USTIC *
 ≉CANAR	*13.4 13.4			*192			±192		_	\$	158		*	168	* *	45	*	-		USTIC *
*CAR LAMANGA			ISOTHERMIC			315		45	315		360		*	360	2.0	0	*	96		UDIC *
*COLTMES			ISCHYPERTH.	_	-		* 65				295		*	295	0.2	0	*			USTIC *
*COTOCOLLAG			ISOTHERMIC			360					250 250		£	360	**	0	÷	62		UDIC +
*COTOPAXI	*10.1 10.3			* 0		360		0	560		360		*	33	**	0	_	120		
*CUENCA			ISOTHERMIC	-	-	360	-	0					*			_	*	1 20		UDIC #
*DAULE			I SOHYPERTH.		_		-	-	360		350		# :h	360	* #	. 0	-	120		"DIC #
*ESMERALDAS							*155		lól		197		-,	197	ध्य में •	15		0		USTIC *
*GUARANDA						_	*204		103		146		*	146	**	120	*	75		USTIC #
			ISOTHERM!C			269	-		259		360		*	360	**	0	4	7.4		USTIC *
#GUAYAQUIL EC							=163		161		197		*	197	**	15	**	5		USTIC *
*GUAYAQUI E			ISOHYPERIH.					35	144		179		*	179	**	15	.9	0		USTIC *
*GUAYLLABAMBA							*137		C		137		*	137	**	73	*	0		USTIC #
IBARRA	415.5 18.5	18.4	ISOTHERMIC	* 32							328		*	328	*	32	*	15		USTIC *
#ISABEL MARIA							*154	48	160		206		*	206	**	14	*	3		# JITZU
*LATACUNGA			ISOTHERMIC				* 223	137	O		110		*	110	**	75	*	2	*	USTIC +
*LIBERTAD (LA					34		* 326	34	0		19		*	19	**	45	*	0	*	ARIDIC *
*LIMONES			!SOHYPERTH.				÷ 0		329		360		¥	360	**	0	•	1.20	*	+ DIGU
*LOJA			ISOTHERMIC		9	360	* 0	0	360	*	360		*	360	**	0	*	120	*	UDIC *
*MACARA			ISOHYPERTH.				* 243	48	69		117		*	117	**	45	*	0	*	USTIC #
*MACHALA			ISOHYPERTH.			106	* 201	53	1.06	*	159		*	159	**	15	*	0	*	# DITZU
⇔MANTA	_		ISOHYPERTH.		27	0	*333	27	0	*	15		*	15	**	45	*	0	*	ARIDIC *
*MENDEZ			ISOHYPERTH.	-	G	360	* G	0	360	•	360		#	360	**	0	*	120	*	PERUDIC *
*MILAGRO	*27.1 27.5	26.3*	ISOHYPERTH.	*145	40	175	*145	40	175	÷	215		#	215	#7	15	*	10	*	USTIC #
≢OTAVAL O	*16.5 lt.1	16.8*	ISCIHERMIC	9 0	9	360	* O	0	360	*	360		*	360	**	0	*	120	*	UDIC ★
*PALMIRA	*13.4 13.4	13.0=	ISOMESIC	*245	115	3	* 245	115	C	ţ	115		*	115	**	75	*	0	*	USTIC *
⇒PAPALLACTA	#11.9 12.1	11.3=	ISOMESIC	* 0	0	360	* 0	0	360	*	360		*	360	**	0	*	120	*	PERUDIC #
*PATATE	*19.1 19.3	18.3*	ISCIHERMIC	* 70	290	0	* 70	290	0	*	290			290	**	70	*	û	*	* SITZU
*PICHILINGUE	*27.0 27.3	26.0*	ISOHYPERTH.	* 97	66	197	* 97	á6	197	*	263		*	263	**	Ō	*	28		USTIC #
*FILLARO	*14.2 14.2			* 0	0	360	* 0	0	360	*	360		*	360	7 *	Č	*	120		* 010U
*PORTOVELO	*25.7 25.6	25	ISCHYPERTH.	*103			£103	_	191		257		*	257	**	Č	*	26		USTIC #
*PORTOV1EJO			ISCHYPERTH.		51		#221	51	88		139		*	139	* *	15	*	0		USTIC #
*PUERTO BOLIV							*261	54	45		86		#	86	**	17	#	ŏ		ARIDIC *
*QUITO ECUADO						360		-	360		360		•	360	**	ō	*	120		UDIC #
*QU:TG			ISOTHERMIC			360			360		360		*	360	方庫	Č	13		\$	UDIG #
*RIUBAMBA			ISOTHERMIC	_	46		P314	46	o		46		*	46	#**	105	•	- 0		ARIDIC #
**********	****	****	*******	****	***	***	***	****	***	***	*****	**	***	****	6 40 4 1	****	***			******
COMPUTED BY	FERTRAL PRO	GRAM V	E08, APR 198	1														6	4 Y F	04/09/#Z

HAME OF STATION	* ME.	AN SO	IL *	TEMPERA		⇒ Cl	140F	VITA	≠≠≠ ≠% E D∧y	**** S MS	* * * * C	# #	********* 0.XAX	P##### CONSEC	F##### UTIVE	*** VAQ	******	******* MCS IS	***	PAGE *****
3741103	* TEM			REGIM	_	*	ΙN			WHEN		*	MCIST IN	SOME	PART		DRY			MOISTUR
	PANN+:	ろした! 四半! 立	412217			*ONE	YEAR	? IS	* SU1	L TĒ	r.P >	5*	IN ONE					* MOIS		
	b a	-				*DRY	M/D	HOI	*DRY	H/D	HOI		YEAR	*TE	MP > 8	**	SHMMED	* WINTE	K *	REGIME
*******	*****		# 			*			*			*		*		**	SOLICE	* SOLST	K *	
LINAS	±25 5	24 4	74 / A	*****	****	****	****	****	****	***	***	***	*****	****	***	* * * *	******	* 30631		
N GABRIEL	#14 2	12 2	24.4	ISOHYP	ERTH.	-	9	0	4351	9	0	*	9		9	**	75	*	~ ~ ~	****
N LORENZO	#29 A	13.5	14.04	ISOMES	IC :	* 0	0	360	. 0	0	360	*	360	*	360	**	õ	-	* (ARIDIC
NTA ISABEL	#20.7	20.2	22.44	ISUHYP	ERTH.	* 0	0	360	* 0	0	360	*	360	*	360	**	Õ) *) *	2100
NTO DOMING	*24 5	24 6	23.0*	1 SUHYP	ER TH.	* 256	94	0	* 266	94	0	*	94	*	94	**	75) #	UDIC
BACUNDO	*15.5	15 5	24.54	ISOHYP	ERTH.	* 0		360	•	0	360	*	360	*	360	**	.0	* 120	-	USTIC
	#27.C	70.3	12.24	ISOTHE ISOHYP	RMIC	* O		360	_	0	360	*	36 0	*	360	**	Õ	* 120	-	UDIC
	*13.0	14 2	12 10	ISOMES	ER TH.		_	360	_	O	360	*	360	*	360	**	č	* 120	-	UDIC
	13.5	12.7	13 0	ISOMES	16 1	• 0		360	_	0	360	*	360	*	360	**	ŏ	* 120	-	UDIC
	*15.5	15 5	15 5+	ISOTHE	10 1	• 0		360	_	0	360	*	36 0	*	360	**	ŏ	* 120		2100
*******	*****	エフャン	10.07	13UIHE	KMIC	* O	0	360	* 0	0	360	*	360	*	360	**	Ô	* 120		DICU

	DETERMIN	O NOITA	F SOIL MOISTL	JR E	REGIME AC	CORDING TO F	PAN	NKLIN	NEWHA	LL SY	STEM OF	F COM	PUTATI	ON			
FOR ECUA																PAGE	1
******	********	****	*****	* * *	****	********	* * * 1	* ** * * *	****	***	*****	****	****	*****	****	****	**
* NAME			TEMPERATURE					CONS			DAYS	*		.DAYS			*
* OF	REGIME	4/D	REGIME	*	OF MOISTL	IRE REGIME					T(2+3)	*		.MOIST	COMP	MCI.	*
■ STATION		T>8		*				SUM.	HINT	SUM.	TNIW	*	SUM.	WINT	SUM.	WINT	*
*********	*******		******	*			*					*					*
*AMBATO	ARICIC					******		*****	****	*****	***		****	*****	****	****	* *
*ANCON	ARICIC	86	ISOTHERMIC		WEAK	ARIDIC	*	75	11	104	11	*					*
*ANGEL (EL)		18	ISOHYPERTH.	*	TYPIC	ARIDIC		18	0	32	0	*					*
*ARGELIA (LA)	010u 010u	360 360	ISOMESIC		TYPIC	UDIC	*					*	180	180	180	180	*
*ASCAZUBI	0100		ISOTHERMIC	*	TYPIC	UDIC	*					*	180	180	180	180	*
*BABAHOYO	USTIC	360 219	ISOTHERMIC	*	TYPIC	UDIC	*					#	180	180	180	180	*
*BAHIA DE CAR	USTIC	98	ISOHYPERTH. ISOHYPERTH.		TYPIC	TROPUST.	*	167	52	167	66	*					¢
*BALZAR	USTIC	229			ARIDIC	TROPUST.	*	98	0	121	0	*					*
*BANOS	UDIC	360	ISOHYPERTH. ISOTHERMIC	*	TYPIC	TROPUST.	*	166	63	166	75	*					*
*CALDERON	USTIC	167	ISCIHERMIC	*	ARIDIC	UDIC	*	125			~ ~	*	180	180	180	180	*
*CANAR	LSTIC	168	ISOMESIC	*		TROPUST.		135	45	142	77	*					*
*CARIAMANGA	UDIC	360		*	ARIDIC DRY	TROPUST.	*	135	33	135	33	*					*
*COLIMES	USTIC	295	ISOHYPERTH.		UDIC	TROPUDIC	#	100	100	100			180	96	180	135	*
*COTOCCLLAO	0110	360		*	TYPIC	TROPUST. JDIC	*	180	100	180	115	*					*
*COTOPAXI	0110	33		∓	TYPIC	-	÷					*	180	180	180	180	*
*CUENCA	0110	360		-	TYPIC	UDIC UDIC						*	180	180	180	180	*
*DAULE	USTIC	197	ISCHYPERIH.		TYPIC		*	1.6	2.2	1/5	, ,		180	180	180	180	*
*ESMERALDAS	USTIC	146	ISOHYPERTH.		ARIDIC	TROPUST.	*	165	32	165	40						*
*GUARANDA	USTIC	360		*	UDIC	TROPUST.	*	0	146	0	156	*					*
*GUAYAQUIL EC	LSTIC	197	1SOHYPERTH.		17910	TRUPUST.	*	145	2.2	1.5	2.2	*	189	74	180	89	*
*GUAYAQUIL	LSTIC	179	ISUHYPERTH.		ARIDIC	TROPUST. TROPUST.		165 165	32	165	32	*					*
*GUAYLLABAMBA	LSTIC	137	ISCTHERMIC		ARIDIC	TROPUST.	*	75	14 105	165 107	14	*					*
*IBARRA	USTIC	328		*	UDIC	TROPUST.	*	75	180		116	•					*
*ISABEL MARIA	LSTIC	206	ISOHYPERTH.		TYPIC	TROPUST.	*	166	40	148 166	180 50	*					*
*LATACUNGA	LSTIC	110	ISOTHERMIC		ARIDIC	TROPUST.	*	105	27	105		_ <u></u>					*
*LIBERTAD (LA	ARICIC	19	ISOHYPERTH.		TYPIC	ARIDIC	-	19	0	34	32 0	.					*
*LIMONES	DIGU	300	I SOHYPERTH.		DRY	TROPUDIC	*	19	U	24	U		127	100	1.0		
*LOJA	LDIC	360	ISCTHERMIC		IYPIC	JDIC	*					-	134 180	18C	149	180	*
*MACARA	LSTIC	117	ISOHYPERTH.		ARIDIC	TROPUST.	*	117	0	117	0		100	18 0	18C	180	*
≯MACHALA	USTIC	159	ISOHYPERTH.		ARIDIC	TROPUST.	*	159	Ö	159	0	*					-
#MANTA	ARICIC	15	ISOHYPERTH.		TYPIC	ARIDIC		15	ő	27	Ö	-					-
∞MEND E Z	PERUDIC	360	ISOHYPERTH.		,	PERUDIC	*		U	21	U	*	180	180	1 00	100	-
*MILAGRO	USTIC	215	ISOHYPERTH.		TYPIC	TROPUST.		165	50	165	50	7	100	100	180	180	-
*OTAVALO	UDIC	300	ISCTHERMIC		TYPIC	UDIC	*	10)	,,	100	70	-	180	18C	100	100	+
₽PALMIRA	LSTIC	115		*	ARIDIC	TROPUST.		105	10	105	10	*	100	100	180	180	-
*PAPALLACTA	PERLDIC	360	· · · · ·	*		PERUDIC	*	100	10	105	10	*	180	180	180	180	7
⇒PATATE	USTIC	290		*	OIGU	TROPUST.	*	75	180	110	180	*	100	100	100	160	Ī
*PICHILINGUE	USTIC	263	ISOHYPERTH.		TYPIC	TROPUST.	•	180	68	180	83						-
*PILLARG	UDIC	360		*	TYPIC	UDIC		100	00	100	65	*	1 60	100	100	100	Ξ
*PORT CYELO	LSTIC	257	ISOHYPEPTH.		TYPIC	TROPUST.	*	180	62	180	77	→	180	180	180	180	*
≠PORTOVIEJO	USTIC	139	ISUFYPERTH.		ARILIC	TROPUST.	*	139	0	139	0	*					*
*PUERTO BCLIV	ARIEIC	86	ISOHYPERTH.		WEAK	ARIDIC	•	86	Ö	99	0	- *					<i>+</i>
*QUITO ECUADO	UCIC	360		#	TYPIC	UDIC	ė.		•	,,	U	*	180	180	180	120	-
*QUITO	010	350		*	TYPIC	UDIC	*					<i>∓</i>	180	180 180	180	180	*
*RIGBAMBA	ARICIC	46	ISOTHERMIC		WEAK	ARIDIC	*	45	0	46	0	*	100	100	1 90	180	-
********	********		******			*******	* * *	****	* * * * * :		*****	****	****	****	***	***	e #
COMPUTED BY FCF	RTRAN PROG	RAM VW	DB, APR 1981													4/29/	

NAME OF STATION	MOISTURE REGIME	CON.D. M/D T>8	TEMPERATURE REGIME	* * *		SUBDIVISION DRE REGIME	* CON * MCI: * SUM	ST (2	+31		****** DAYS F(2+3) WINT	****	COMP	DAYS MOIST	COMP	DAYS DAYS MOI	
SALINAS	ARICIC	9	ISOHYPERTH.	* * *	****	*********	*****	* * * *	* * * *	****	****	****	****	*****	****	****	* *
SAN GABRIEL	UEIC	36ú	ISOMESIC		TYPIC	ARIDIC	* (9	0	9	0	*					
SAN LORENZO	UCIC	36U	ISOHYPERTH.	-	TYPIC	01C	*					*	180	180	180	180	
SANTA ISABEL	USTIC	94	ISOHYPERTH.			UDIC	*					*	180	180	180	183	
DAINOC OTHAZ	UEIC	360	ISOHYPERTH.		ARIDIC	TROPUST.	* 94	+	0	94	0	*					
TABACUNDO	LDIC	360	ISOTHERMIC	*	TYPIC	J100	*					*	180	180	180	180	
IPUTINI	UDIC	360	ISOHYPERTH.		TYPIC	JDIC	*					*	180	180	180	180	
TI SALEO	OIGU	360	ISCMESIC		TYPIC	JDIC	*					*	180	180	180	180	
TULCAN	2100	360	ISOMESIC	*	TYPIC	ODIC	*					*	180	180	180	189	
JYUMB ICHO	UDIC	360	ISGTHERMIC	*	TYPIC	UDIC	*					*	180	180	180	180	
*********	*****	****	TOUINERMIL	*	TYPIC	UDIC	*					*	180	18C	180	180	

Appendix IV

In this appendix, eleven pages called a data sheet packet—an explanation is included. If you have any questions concerning the data sheet, these pages should be reviewed.

DATA SHEET PACKET - AN EXPLANATION

PARRICE

SAMPLED AS: LOAMY-SEZIETAL, MINED, MESIC TYPIC DISTROCHREPT

\$ 78VT-025 -002 SAMPLE NOS. 78P1791 - 1796 DATE 6/ 5/79

SPODOSOL STUDY WINDHAM COUNTY

U. S. DEPARTMENT OF AGRICULTURE SOIL CONSERVATION SERVICE NATIONAL SOIL SURVEY LABORATORY LINCOLN, NESRASKA CREERAL METHODS 1811, 241, 28

		-1	-2-	3	-4	-5	-6	-7												- 19-	-20-
SAMPLE NO.		(CH)			.002	.002	.05	.0002	CO3 LT .002 - PCT	7 T WB - 002 - 02 OF <21	COARS: .02 05	.05 10	.10 25	-SAND- 8 25 50	c .5 -1	7c 1 -2	2 -5	15E PR VR 5 	ACTIONS		
781791 781792 781793 781794 781795 781796	25 35 45 55	0- 5- 23- 35- 50- 61-1	23 35 50 61	11 821I1 822 823 83 20	11.9 10.4 11.3 10.4 7.4 5.7	38.5 39.0 42.6 38.9 25.8 7.8	49.6 50.6 46.1 50.7 68.8 86.5			22.4 21.2 23.4 20.3 12.7 4.5	16.1 17.8 19.2 18.6 11.1 3.3	12.8 12.1 12.4 11.4 6.4 3.4	11.7 10.4 11.4 9.8 8.7 10.0	5.4 5.5 4.6 5.5 7.3 12.4	8.1 7.8 6.9 9.0 15.9 27.4	11.6 14.8 10.8 15.0 30.5 33.3	7 5 6 7 10	13 27 11 20 29 30	5 48 19 37 35 37	88 50 78 90 96	25 83 56 74 79 85
SAMPLE NO.	82N 80.	6 A 1C	6815	E E E E E E E E E E E E E E E E E E E	TOTAL S 6RJA CP <	(D BX 2B 6C2B	TRACTA AL 6G7A	T) BLE BH 6D2A >	CEC 8D1	O/CLAY) 15 BAR BD1	(ATTER - LIE LL 4F1 PCT <	BERG) ITS - PI 4P	(- DOL PIELD HOIST 4AJL	K DEUS 1/3 BAB 4A10 G/CC -	OFEN OFEN OFEN OFEN	COLE SOIL AD1 COLE	(FIELD BOIST 484	HATER 1/10 BAR 481C -PCT (CONTEN 1/3 BAR 4B1C OF <248	15 15 848 482 (>	SOIL 4C1 CE/CE
781791 781792 781793 781794 781795 781796	1 2 3 9 5 6	5.59 2.41 1.33 0.91 0.58 0.41	0.208 0.113 0.064 0.053			1.8 1.9 1.8 1.7 1.1	0.3 0.7 0.5 0.4 0.3		2.81 1.26 0.61 0.55 0.62 0.63	1.32 0.92 0.60 0.53 0.54	31	1		0.90 0.98 1.17 1.20 1.50	1.01	0.010			34.7 30.0	15.7 9.6 6.8 5.5 4.0 5.3	0.25 0.27
SAMPLE SO.	HZN NO.	- #84 CA 5841 6#28	01C I 86 5841 602D	ITBACTA WA 584A 692B	BLE BI E 5B44 6Q2B	ASES -) SOM BASES -EBQ /	ACID- ITY 6#2B 100 G	BITA AL 6G1E	(SUB CATS SAJA	-CBC #H4- OAC 5A8A) B15E3 + AL 513B	AL SAT 5G1	-BASE SUM 5C3	SAT- HH4 OAC SC1	CO3 AS CACO3 <288 6818 >	S RES. OBBS /CB 811			(-PH -CACL2 .018 8C1E 1:2	820 8C1A 1:1
781791 781792 781793 781794 781795 781796	2 3 4 5 6					0.1 28 72 72		7.5 3.5 2.2 2.0 1.1 0.6	39.4 22.1 11.7 10.2 7.6 5.2	33.4 13.1 6.9 5.7 4.6 3.6	8.0 3.6 2.2 2.0 1.1 0.6	94 97 100 100 100	TR TR TR TR	1 12 72 72 72 1		23000				3.2 4.0 4.2 4.2 4.4 4.4	3.5 4.1 4.3 4.3 4.4
SASPLE BO.	HZN HO.	HA C 6A4A <- PC	-SPO PIRO PZ 6C5A	DIC HOR PHOSPHA AL 6G5A <288->	IZON C TE BIT PE+AL (DI DI-CI FE+AL	BITEBI BACTABI FE+AL VIDED PCT CLAY	A LE) AL+C BI) PCT CLAT	VCCOR OL INDEX)			****		(((7A2B	- 1- 7128	- CLA BAY - 7128 ABOUR	TINERA T) 7128	LOGY - (DT 713	7A3	TOTAL RES 781A	DOE TEATE 781A
781791 781792 781793 781794 781795 781796	3	1.8	0.6	0.3	0.4	TE	0.2	167 304 73 75 43 132													

PARILY CONTROL SECTION: DEPTH 25-100 PCT CLAY 7.6 PCT .1-758H 86.4. SPODIC HORIZON: INDEX OF ACCORDE 0 ESTIMATED BULK DESSITY FOR LAYER 1, 4, 5, 6, ABALYSES: S- ALL, ON SIRVED <2MB BASIS

DATA SHEET HEADING

WARWICK

SAMPLED AS: LOAMY-SKELETAL, MINED, MESIC TYPIC DYSTROCHAEPT

S 7887-025 -002

SAMPLE NOS. 78P1791 - 1796

DATE NOVEMBER . 1979

O. S. DEPARTMENT OF AGRICULTURE SOIL CONSERVATION SERVICE NATIONAL SOIL SURVEY LABORATION LINCOLN, HEBRASKA

SPODOSOL STUDY WINDHAM COUNTY

GREERAL BETHODS 1814, 241, 28

<u>Warwick</u> Name of the soil series, pedon sampled to represent.

Sampled As Classification (field) of pedon at time of sampling.

Pedon number \$78 VT-025-002

| S = Special sample | S = Special sample |
| 78 = calendar year sampled |
| VT = state where sampled - 2-letter FIPS code for state where sampled |
| 025 = FIPS county code |
| 002 = consecutive pedon number for calendar year for county |

Sample numbers--NSSL assigned numbers.

78 = year (PY)

P = permanent storage of data on NSSL computer disk (T = temporary storage)

1791 = last four digits is consecutive sample number (NSSL)

Date-date of printout

Project name and/or county sampled.

General methods for soil preparation -- see SSIR No. 1.

Gameral conventions. Unless otherwise specified analyses are calculated to an oven dry weight base. Therefore, if analyses are reported on a < 2 mm bese, it means that analyses are on all particles < 2 mm in size and results are calculated using an oven dry sample weight. Conversely, if the analyses are reported on a whole soil base, it means that the analyses are for all particles with coarse fragments included, and are calculated using an oven dry sample weight.

Footnotes

PARILY CONTROL SECTION: DEPTH 25- 71 PCT CLAY 51 PCT .1-75##

ABALISES: S= ALL ON SIEVED COME BASIS

MINERALOGI: KIND OF MINERAL HI HICA AT HONTHORILL RE RAOLIBITE CL CHLORITE QZ QUARTZ RELATIVE AMOUNT 6 INDETERMINATE 5 DOMINANT 4 ABUNDANT 3 SODERATE 2 SHALL 1 TRACE


Family control section:

Depth limits of control section; calculated weighted average % clay of control section; and calculated weighted average % 0.1-75 mm fraction for use in determining loamy vs. silty particle size class.

Estimated bulk densities for layer: . . . self explanatory.

Analyses: Tier 1, column 1 provides explanation of soil preparation code.

Mineralogy: Identifies clay mineralogy codes and relative amount codes.



General: First two columns (unnumbered columns) are the same for each tier. The first column on the left is the MSSL sample number, the second number is the horizon number. Except for columns 1-3, tier 1 is entirely particle size and coarse fragment data. Particle size distribution (columns 4-15) is reported on the < 2 mm fraction and is determined after destruction of organic matter, washing free of soluble saits, drying the sample, and obtaining a base weight, and dispersion with sodium The first column on the left is the NSSL sample

- Column 1. Space after horizon number for two letter preparation code.

 Space after horizon number for two letter preparation code.

 Space after horizon number for two letter preparation code.

 Space after horizon number for two letter preparation code.

 Space after horizon number for two letter preparation code.

 Space after horizon number for two letter preparation code.

 Space after horizon number for two letter preparation code.

 Space after horizon number for two letter preparation code.

 Space after horizon number for two letter preparation code.

 Space after horizon number for two letter preparation code.

 Space after horizon with space column preparation code.

 Space after horizon number for two letter preparation code.

 Space as "S" but clod parameters (letter 2, columns 12-14 and 16-18) reported on a whole soil base.

 Space after horizon number for two letter preparation preparation code.

 Space after horizon number for two letter preparation code.

 Space as "S" but clod parameters (letter 2, columns 12-14 and 16-18) reported on a whole soil base.

 Space as "S" but clod parameters (letter 2, columns 12-14 and 16-18) reported on a whole soil base.

 Space after horizon number for two letter preparation code.

 Space after horizon number soil base.

 Space after horizon are shipped on a whole soil base.

 - < 2 mm material. Results reported on whole soil base.
- Column 2. Denth in cm
- Horizon notation. Arabic numbers are used instead of Roman Numerals. Column 3. Printed horizon designation may be truncated since only six spaces available for notation. The entire notation is stored in computer
- Column 4. % total clay. Particles < 0.002 mm. Sedimentation and pipette analysis.
- Column 5. % total silt. Perticles 0.002 mm to 0.05 mm. Sedimentation and pipette analysis.
- Column 6. % total sand. Particles 0.05 mm to 2.0 mm. Sieve analysis.
- Column 7. % fine clay. Particles < 0.0002 mm. Centrifugation and pipette analysis.
- Carbonate < 0.002 mm. Sedimentation and pipette analysis, HCI treatment, and CO evolution % carbonate clay. Carbonate < 0.002 mm. Sedimentation and pipette analysis, HCI treatment, and CO Subtracted from col. 4 to obtain % noncarbonate clay when determining taxonomic particle size class. Taxonomy, p. 384, footnote 1.</p>
- Column 9. % fine silt. Particles 0.002 mm to 0.02 mm. Sedimentation and pipette analysis.
- Column 10. % coarse silt. Particles 0.02 mm to 0.05 mm. By difference--coarse silt = [100-(% clay + % fine silt + % sand)]
- Column 11. % very fine sand (VFS). Particles 0.05 mm to 0.1 mm. Sieve analysis. Treated as silt to distinguish loamy and silty particle size classes. See Soil Taxonomy, p. 383.
- Column 12. % fine sand (FS). Particles 0.1 mm to 0.25 mm. Sieve analysis.
- Column: 13. % medium sand (MS). Particles 0.25 mm to 0.5 mm. Sieve analysis.
- Column 14. % coarse sand (CS). Particles 0.5 to 1.0 mm. Sieve analysis.
- Column 15. % very coarse sand (VCS). Particles 1 mm to 2 mm. Sleve analysis.
- Column 16. Weight % coarse fragments 2 mm to 5 mm. Reported on < 75 mm base. Sieve analysis.
- Column 17. Weight % coarse fragments 5 mm to 20 mm. Reported on < 75 mm base. Sleve analysis.
- Column 18. Weight % coarse fragments 20 mm to 75 mm. Reported on < 75 mm base. Sieve analysis in field (at least 20 kg sampled needed) or calculated from volume estimates.
- Column 19. Weight % particles and fragments 0.1 mm to 75 mm. Reported on < 75 mm base. Sieve analysis and/or estimates of > 20 mm fraction. Used to distinguish loamy and silty particle size classes (see p. 384, Soil Taxonomy).
- Weight % fragments > 2 mm. Reported on whole soil base. Includes fragments from 2 mm to 250 mm in size. Calcul from sieve analysis and volume estimates. Used to convert analysis on < 2 mm base to whole soil base by formula: datum whole soil = datum < 2 mm x [1-(col. 20/100)].

 Coarse fragments assumed diluents in this equation. Column 20. Weight % fragments > 2 mm. Calculated

-1-- -2-- -3-- -4-- -5-- -6-- -7-- -8-- -9-- -10- -11- -12- -13- -14- -15- -16- -17- -18- -19- -20-

EXTR TOTAL (- - DITH-CIT - -) (RATIO/CLAY) (ATTERBERG) (- BULE DEBSITY -) COLE (- - - WATER CONTENT - -)
P S EXTRACTABLE 15 - LIBITS - FIELD 1/3 OVER WHOLE FIELD 1/10 1/3 15
PE AL RB CEC BAR LL PI HOIST BLE DRY SOIL BUIST BAR BAR
6S1A 6H3A 6C2B 6G7A 6D2A 8D1 8D1 4F1 4F 4A3A 4A1D 4A1E 4D1 4B4 4B1C 4B1C 4B2 ORGM TOTAL WEB THOLE JAKPLE HZM SOIL NO. DATE BETE AC I MO. -PCT OF <388 -PCT <0.488 <- - G/CC - - -> CB/CB <- - - PCT UP <288 --> CM/CM 78 781 1 1.57 0.143 0.71 0.33 1.29 1.55 0.063

- General. Unless otherwise noted, analyses are reported on < 2 mm base.
- Column 1. % organic carbon. Wet combustion analysis (Walkley-Black). Multiply % organic carbon by 1.7 to estimate organic matter.
- Column 2. % total nitrogen. Kjeldahl digestion analysis. Carbon/nitrogen ratios 3 10-12 in mollic epipedons. Higher ratios may indicate a nitrogen immobilization problem.
- Column 3 % total extractable phosphorous. Perchloric acid digestion. Presently not determined at NSSL.
- Column 4 % total sulfur. Leco ignition, KIO₃ titration. Used to assess potential acidity in sulfidic soils or mine spoils.
- Column 5 % iron. Dithionite-citrate extractable (Fe Di-Cl). Commonly called "free iron." Used in spodic horizon criteria
- Column 6 % aluminum. Dithionite-citrate extractable (Al DI-C!). Used in spodic horizon criteria.
- Column 7 % manganese. Dithionite-citrate extractable.
- Column 8 CEC-7/clay ratio. Ratio of CEC-7 (Tier 3, col. 9) to total clay (Tier 1, col. 4) Related to activity of clay fraction.
- Column 9. 15-bar/clay ratio. Ratio of 15-bar water (Tier 2, col. 19) to total clay (Tier 1, col. 4). If clay > 10% and if organic carbon/clay (Tier 2, col. 1/Tier 1, col. 4) is < 0.1, ratio should be between 0.3 and 0.5. Ratios > 0.6 may indicate a dispersion problem. See p. 384, footnote 1, Soil Taxonomy.
- Column 10. Liquid limit reported on < 0.4 mm base. Used in Unified and AASHO engineering classifications.
- Column 11 Plasticity index reported on < 0.4 mm base. Calculated by liquid limit (col. 10) minus plastic limit. Used in Unified and AASHO engineering classifications.
- Column 12. Bulk density at field moisture content q/cc (dry wt./moist vol.). Heasured on natural fabric clod.
- Column 13. Bulk density at 1/3-bar or 1/10-bar moisture content g/cc (dry wt./moist vol.) Measured on natural fabric clod. Used on SCS-SOILS-5. 1/10 bar run on sandy soils that have 15-bar moisture (tier 2, col. 19) < 5% throughout.
- Column 14 Ovendry bulk density d/cc (dry wt/dry vol.). Heasured on natural fabric clod.
- Column 15. COLE. Coefficient Of Linear Extensibility whole soil base. Calculated from increase in bulk density between moist (tier 2, col. 13) and ovendry conditions (tier 2, col. 14). Used in determining shrink-swell potential for SCS-SOILS-5 or total potential linear extensibility for vertic subgroups. Total potential extensibility equals horizon thickness in cm x COLE: summed for each horizon to the depth in question (assumes no large cracks between air-dryness and 1/3-bar moisture and that coarse fragments are diluents).
- Column 16. Weight % field moisture content (wt. H_2O/dry wt. soil x 100).
- Column 17 Weight % moisture content at 1/10-bar tension. Calculated from water retained in natural fabric clods for loamy and clayey soils and from water in < 2 mm sample for sandy soils. Used to represent field capacity for sandy soils.
- Column 18. Weight % moisture content at 1/3-bar tension. Calculated from water retained in natural fabric clods. Used to represent field capacity for loamy and clayey soils.
- Column 19. Weight % moisture content at 15-bar tension. Calculated from water retained in < 2 mm soil sample. Represents the wilting point.
- Column 20. Water Retention Difference (WRD) Reported as cm of water per cm depth of soil. Volume fraction for whole soil of water between 15 ber and an upper limit, usually 1/3 bar but 1/10 bar for sandy soils. After adjustment for salts (tier 4, coi. 11 salt sheet) and root exfoliation, WRD provides estimate of Available Mater Capacity (AWC).

-1	-2	-)	-4	-5	-6	-7	-8	-9	-10-	-11-	-12-	-13-	-14-	-15-	- 16-	-17-	-18-	- 19-	-20-
								ACID)										

	-																	
SIMPLE IO.	EZN BO.	584A 682B	5841 6020	584A 6P2B	584A 602B	2585 -) 258 8/ 385 -HEQ /	6.82B	AL 6G1F	CATS	OAC	* AL	SAT	SUB	OAC	CACO3	ORMS /CB		820 8C 1A
781793						TR									-	13000	1:2	A. 3

SAMPLE VO.	CA HZM 5B4A NO. 6M2M	40AC EXTRACTABLE BASES -; BG MA K 50H 5844 5844 5844 BASES 602D 6P2B 6Q2B 	TTT SUM WB4- CATS DAC 6H2B 5A3A 5A6A	NA 5D2 5 E	8ASE CARBOMATE SATURATION AS CACO3 SUB NH9OAC (2Em (20Hm 5C3 5C1 6E18 (PCT> (PCT>	CASO4 AS (PB) GTPSUM SAT CACL2 <288 <2088 PASTE ONE 6FIC 6FIC 3C18 8C18
78 781		5.3 0.1 1.1		TR TR	1	(* -PCT -> 1:2

- General: The NSSL has acid and salt data sheet formats. The first two tiers are the same for each format, but tiers 3 and 4 differ. All analysis on tier 3 are reported on the < 2 mm base. Columns 1 through 4 are bases extractable in normal ammonium acetate (N NH₄OAc) at pH 7 (CEC-7).
- NH₁OAc extractable calcium (Ca) meq/100 g. Note that Ca is not reported for horizon 1 salt sheet because the sample is calcareous. NH₁OAc dissolves and extracts Ca from carbonates, gypsum, and other soluble salts. Commonly used as soil trit for Ca. "Convert to Kg/Ha = Ca meg/100/g x 20 x thickness in cm x bulk density (tier 2, col | 13) x {1-(Vol. > 2 mm/100)}. Column 1.
- NH₁OAc extractable magnesium (Mg) meg/100 g. Commonly used as soil test for Mg. Convert to kg/ha = Mg meq/100g x 12 x thickness (cm) x bulk density (tier 2, col. 13) x $\{1-(Vol. > 2 \text{ mm/}100)\}$ Column 2.
- Column 3. NH_uOAc extractable sodium (Na) meg/100 g.
- $NH_{Q}OAc$ extractable potassium (K) meg/100 g. Common soil test for K. Convert to kg/ha = K meg/100 g x 39 x thickness (cm) x bulk density (tier 2, col. 13) x {1-(Vol. > 2 mm/100)} Column 4
- Column 5 Sum of bases (Ca + Mg + Na + K) extractable by N $\mathrm{NH_{L}OAc}$ (Cols. 1+2+3+4)
- Column 6. Extractable acidity measured at pH = 8.2 meg/100 g.
- Normal potassium chloride (KCl) extractable aluminum (Al) meq/100 g (acid sheet only; blank on salt sheet) Normally little present if pH > 5 (col. 20).
- Cation Exchange Capacity at pH 8.2 (CEC-8.2). Calculated by summing NH, OAc extractable bases (col. 5) a extractable acidity (col. 6). Not reported on salt sheet for horizons with carbonates or soluble salts. Column 8.
- Column 9. CEC at pH 7 (CEC-7). Measured by retention of ammonium by soil at pH 7
- Column 10. Acid sheet. Bases plus Al (CEC-unbuffered). Calculated by summing NH_{ij}OAc extractable bases (col. 5) and KCl extractable Al (col. 7). Commonly called effective CEC of acid soils.
- Salt sheet. Exchangeable Sodium Percentage (ESP). Calculated by dividing extractable Na (col. 3) by CEC-7 (col. 9) and multiplying by 100. Extractable Na is adjusted for soluble Na (fier 4, col. 3) if present, NH₄OAc will extract both exchangeable and soluble Na. Calculated, Exchangeable Na = Extractable Na meq/100g [(soluble Na meq/1)(SP-Tier 4, col. 10 salt sheet)+1000]. Used is natric horizon criteria. Column 10.
- Acid sheet. % aluminum saturation. Calculated by dividing KC1 extractable A1 (col. 7) by bases plus A1 (col. 10) and multiplying by 100. Considered root limiting if $\geq 60\%$. Column 11
- Column 11. Salt sheet. Sodium Absorption Ratio (SAR). SAR = Na (Tier 4, col. 3)/v Car Mg/2 (tier 4, cols [+2) Criteria for natric horizon and used in assessing satinity problems. Set up to assess quality of irrigation water
- % base saturation, sum of cations. Calculated by dividing sum of bases (col. 5) by CEC-8.2 (col. 8) and multiplying by 100. Used in most affic-ultic separations in Soil Taxonomy. Column 12.
- \S base saturation, NH_uOAc. Calculated by dividing sum of bases (col. 5) by CEC-7 (col. 9) and multiplying by 100 Used in mollic, umbrid, and eutro-dystro breaks in Soil Taxonomy. Column 13.
- Column 14. % calcium carbonate equivalent. Carbonates of Ca + My analyzed. Expressed as CaCO₃ equivalent. Used to assess calcic horizons and mineralogy class.
- Column 15. Resistivity onms/cm. Determined on saturated paste. Used to determine potential hazard for corrosion of uncoated
- Salt sheet. % CaSO, as gypsum < 2 mm base. Used to determine potential hazard for corrosion of concrete, also to assess gypsic horizons and mineralogy class. Column 16. Sait sheet.
- Column 17. Sait sheet. % CASO₄ as gypsum < 20 mm base. Used in mineralogy class--see Taxonomy p. 387.
- Acid sheet. pH 1:1 N KCl or NaF pH. KCl pH used as indicator of net negative charge. Commonly 1 unit less than pH in 1:1 soil-water suspension. NaF pH > 10 indicates presence of structually weak Al associated with allophane, organic matter, or halloysite. Acid sheet.
- Column 18. Sait sheet. pH of saturated pasta.
- pH in 1:2 soil-0.01 M calcium chlorida ($CaCl_2$) suspension. Used to eliminate possible seasonable variations in pH. Normally about 0.5 units below the 1:1 soil-water pH (col. 20) but may be greater for acid soils. Relates somewhat with base saturation when BS > 60%; ie, if CaCl_pH is 6.0, NH, OAc base saturation is commonly > 75%; if pH is > 7.5, the soil is normally saturated with bases. Used in reaction classes—see Soil Taxonomy pp. 388, 390. Column 19.
- Column 20. pH in 1:1 soil-water suspension. Normally varies according to season of the year.

-1-- -2-- -3-- -4-- -5-- -6-- -7-- -8-- -9-- -10- -11- -12- -13- -14- -15- -16- -17- -18- -19- -20-

ACID

(----) TOTAL DOB PR+AL PR+AL AL+C OF (- -DIVIDED ST- -) ACCUM SABPLE HZW NO. NO. 6141 6C51 6G51 DI-CI PCT 7811 7811 712B 712B 7128 713 <- PCT OF (288-) FE+AL CLAY CLAY PELITIVE ABOUNTS -> 781791 781792 0.2 304 SALT

(- - - - - - - WATER ZYTHACTED PROM SATURATED PASTE- - - - - -TOTAL BLEC. ĸ CO3 ECO3 CL 304 820 SALTS COMD. SAMPLE HZE EST. MU. 6818 6010 6P18 6Q18 6I18 6J18 6E1C 6E1C 6E1C 805 20888 <- -PCT -> 78 784 0.7 58.0 0.1 2.80 07 3 82 3 ER 3 CL 1 EE18

General. The first 12 columns of the acid and salt data sheets differ and are discussed separately. All analyses in tier 4 except mineralogy are reported on the < 2 mm base. Clay mineralogy is reported on total clay base unless otherwise noted. Grain counts are on base of fraction(s) specified in heading.

ACID

- Column 1. \S sodium pyrophosphate (0.1 M Na $_4$ P $_2$ 0 $_7$) extractable carbon (C-pyro). Used in spodic horizon critaria with < 0.1 \S Fe-Pyro (col. 2).
- Column 2. % sodium pyrophosphate extractable iron (Fe-pyro). Used in spodic horizon criteria. Usually considered to be ron associated with organic matter
- Column 3 Sodium pyrophosphate extractable aluminum (Al-pyro) Used in spodic horizon criteria.
- Column 4. Pyrophosphate Fe+Al/dithionite citrata Fe+Al ratio. Calculated by dividing Fa-Pyro + Al-pyro (cols 2+3) by Fe DI-CI + Al DI-CI (tiar 2, cols. 5+6). Used in specific horizon criteria.
- Column 5. Pyrophosphate Fe+Al/clay ratio. Calculated by dividing Fe-pyro + Al-pyro (cols. 2+3) by % clay (tier 1, col 4) Used in spodic horizon criteria.
- Column 6. Pyrophosphate Al plus carbon/clay ratio. Calculated by dividing C-pyro + Al-pyro (cols. 1+3) by % clay (tier 1, col. 4). Used when column 2 < 0.1.
- Column 7. Index of accumulation of spodic horizon. Calculated by subtracting 1/2 clay % (tier 1, col. 4) from CEC-8.2 (tier 3, col. 8) and multiplying by thickness (cm) of subhorizon. Total index is sum of all subhorizons of spodic horizon. See Soil Taxonomy p. 32 for limits of spodic horizon.

SALT

- Column 1-9 are soluble cations and anions (salts) in a water extract from a saturated soil paste.
- Column 1. Water soluble Ca meg/1.
- Column 2. Water soluble Mg meg/1
- Column 3 Water soluble Na meq/1
- Column 4 water soluble K meg/1.
- Column 5. Water soluble carbonate (CO₃) meg/1. pH (Tier 5, col. 20) must be \geq 9 in order to have carbonate anion.
- Column 6. Water soluble bicarbonate (HCO₁) meq/1.
- Column 7. Water soluble chloride (C1) meq/1.
- Column 8. Water soluble sulfate (SO_L) meq/l. Used along with gypsum (Tier 3, cols. 16 & 17 salt sheet) in estimating potential corrosivity of concrete.
- Column 9. Water soluble nitrate (MO_{χ}) mag/1.
- Column 10. Weight % water content of saturated pasts (Saturation Percentage--SP). Used to convert from meq/l (cols, 1-9 sait sheet) to meq/100g. Datum meq/100g = (Datum meq/I)(SP)/1000.
- Column 11. % total estimated soluble salts in soil. Calculated from conductivity (EC, col. 12 sait sheet) of saturation extract. Total sait meq/1 = -4.2333 + 12.2347 (EC, col. 12 sait sheet) + 0.0580 (EC) = 0.0003 (EC) Total sait % = total sait meq/1 x 0.00064 SP (col. 10 sait sheet).
- Column 12. Electrical conductivity (EC) of saturation extract metho/cm. Used as a measure of salinity.

BOTH ACID AND SALT DATA SHEETS

- Columns 13-16. Relative amounts of clay minerals. 1--trace, 2--small, 3--moderate, 4--abundant, 5--dominant, 6--indeterminate
- Columns 17-18. Quantitative amounts of clay minerals in percent (Differential Thermal Analysis--DTA).
- Column 19. % total resistant minerals. Optical count of minerals in dominant fraction (silt or sand). Is used to assess family mineralogy class.
- Column 20. % dominant weatherable mineral in fraction(s) indicated.

SOIL INTERPRETATIONS RECORD

	A	D) 	F	F	_G	H	. 1	J	K
	OEPTB (M)	USDA TE	ATUEL	1	winter		IL PROPERTIES	FAACT.	ŕ	DIAM SIN.	M FERML LI	131 E	LIGHT	PLAS. HOTT
#1	<u> </u>			 				(PCT)		- "-	•	, ma		INDEX
				-										-
II.				ļ —				 					-	
						_	_	_	_					
"	MPTH	CLAY	MORST BULL Divisity	MRWA- BALITY	AVAILABLE WATER CAPACITY	SOL SUCTION	R	Deleta - Dali				EXC	CORRESELY	Y 117
<u>ar</u>	(100)	:PCT OF <340	לוכד פן	(D. MI)	IN THE	(pH)	iMeOS/CM	POTENTIAL		i cao				OCHIE
<u> </u>	DUFTY AS											\exists		
$\dot{\mathbb{H}}$	ABOVE								#=					
	Column				r zone. See ti	er 1, col	. l. Conver	t ca to	n-in	cm/2.5	14			
	column (10 cm = 0+4 in. 1 cm = 6+6 an	1 11-15 A	nd val esti	mates of	coarse	frages	ınti in	descri	gtion.	
					l, cols. 4-6 and m, silty clay.									
(columen (See cal	enginearir s. F, G, H, pedon 2,	. 1, 3,	sification. Sei K. = * CH	e *Gu+de	for Interpre	ting Eng	inmeri	ig Uses	21 201	15" (61	.037	
(Calumn (gineering o		ication. See G	EUS and G	Cals. F, G,	н, т, э,	•					
(column i	. Fraction	> 3 inches	(75 m	a) calculated for [(cols. 16 + 17	rom (lata (in tier 1 1 - cal 70)	/100 + (cois. 1	6 + 17	+ 18) -			
		Evamoles	-nedon 1 1	0-21								0-55)	•	
	Ölümn /				(material < - 5									
		% passin Example-	g 94 = (100 -padon 1, 0) - (co)-iu cas	(17 + 18)									
(column d	. A passin	g 410 sieve	(mate	rial < 2 am), (Calculate		n tier 1	,					
		EX 8PPD 18 *	•pedon :, :	3-10 cm	16 + 17 + - (1 + 7 + 35)		38) + <u>625</u> .							
(Column I	. % passin	g \$40 sieve	(mate	rial < 0 42 mm) ((cols. 14 * 15	estimate (col. G/)	using data	in tier	1.					
		Example.	-pagon I, c	i-in cour	(4.7 + 4 0)(62			1 # 62-5	• 275					
•	column 1	* passin	kg \$200 siev	re = (c	erial < 0.074 m ol. G/100) {(co	m) calcula 4 + 5)	te using da + (coi. 11/	ta in ti 2)	er 1.					
		% pa	-pedan 1, (issing \$200 : 8)} = 0.62	sieve .	(62/100) [.13	2 + 54 2) + (9.6/2)]	. (0.62) [(67.	4)				
,	olumn .	. Liquid I		See t	er 2 co1 10									
(column r	. Plastici		91), Şi	e tier 1, 101	11.								
,	olumn t	. Same as	col A.											
(Column 1				, col. 4 % clay = 45.8%									
c	olumn x				er 1, col 15. bulk density *	1 11 0/0								
c	olumn O	Permeabil				1.17 Y/C	•							
c	olumn P	and restr	*ictions an	d for s	AVC) Estimate alts. If no ro	ot restri	ctions or sa	11. 70). Ilts VRD	VRO m	st be d	adjuste eter of	d to ro	ating d	epth
c	oluman Q	Soil read	tion (pH).	See :	layer minimum A ier 3, col. 20 reaction = 5 1.	0.13	•							
c	olusan R	Salinity.	. Based on	EC (ti	e: 4, coi 12, EC = 2.8 ratio	salt shee	t). Saw GIE	.us for c						
c	olumn S	Shrink-sw	rell potent	ial. 8	ased on COLE (t	er 2, co	1. 15) See	GIEUS fo	r clas	148.				
c	oluman T				by series. Can		-	weier nom	ograph					
c	otu sa n U	T factor-	-values as:	igned	by series.									
			-		ee Technical Gu									
	olumn 4	Example	pedon 1, 0	-10 cm	by multiplying % organic matte	r + 4.55 .	4 1.7 + 7 49							
c	o iusen X	for class	43.		. Based on Res m zone acidity					-			540 GI	EUS
c	olumn Y	Example Gypsu Conta	pedon 2, 8) m is parti ins > I's g	n sp (1-140 ci llly so /psuss,	ed on soluble s and pH (tier 3 tier 4, col 10 m. 50, ppm = 9 luble in water it has a potent adjusted for pr	, salt shi 0 x 79 x i and contr ial source	eet) x 0.48, 0.48 = 3412 ibutes 30 me e of > 7,000	ppm or m	oderate	rating		tier 3,		

Form 5 will most likely be converted to metric system within 1980. For purpose of comparison to data sheet, depths are listed in cm.

Example Pedon 1

ACID SHEET

SAMPLED AS: LOAMT-SKELETAL, MIXED, FRIGID PRAGIOCHREPT

S 7788-031 -001

Sample NOS. 78P 552 - 556 DATE JULY 1979

U. S. DEPARTMENT OF AGRICULTURE SOIL CONSERVATION SERVICE MATIONAL SOIL SURVEY LABORATORY LINCOLN, MEBRASKA COOK AND ST. LOGIS COUNTIES

GENERAL METHODS 1814, 241, 28

				,	.,															
		-12	3	-4	- 5	-6	-7	-8	-9	-10-	-11-	-12-	-13-	-14-	-15-	-16-	-17-	-16-	- 19-	-20 -
SAMPLE ED.		DEPTH (CH)	RORIZON	CLAT	TOTAL SILT	51 ND -05	PINE LT	CO3	.002	.02	.05	. 10	-SAND- 6 .25	 . 5	VC 1 -2	(-C011	5 -20	IGBT - 20 -75	. 1-	PTT OF MHOLE
78 552 78 553 78 554 78 555 78 556	25 35 45	0- 10 10- 23 23- 38 38- 63 63-122	11 82181 82281 83 CX	13.2 19.5 12.2 8.8 12.7	54.2 42.6 23.2 40.5 41.4	32.6 37.5 64.6 50.7 45.9			36.4 28.2 15.5 35.9 31.1	17.8 14.4 7.7 14.6 10.3	9.6 7.1 6.0 12.0 10.3	9.5 8.8 7.2 11.9	4.8 4.9 5.3 6.3 5.8	4.7 7.3 11.3 8.9 8.6	4.0 9.8 34.8 11.6 11.2	1 6 8 5 8	2 16 8 9	35 31 29 30 27	52 67 77 66 66	50 61 58 52 61
SIMPLE NO.	HZN HO.	6 M 6 M	AL EXTR P 18 651A PC1	TOTAL S 6831	FE 6C2B	TBACTA AL 6G71	(T) IBLZ H# 6D21	CZC 8D1	CLAY) 15 BAR 8D1	(ATTER - LIR LL 4F1	BERG) ITS - PI 4F	(- BUL PIELD HOIST 4131	R DBMS 1/3 918 4110	ITY -) OVER DRY 4118	COLE NHOLE SOIL 4D1	(FIELD HOIST 484	1/10 818 481C	1/3 Blb 4B1C	15 15 BAR 482	WED WHOLE SOIL 401
78 552 78 553 78 554 78 555 78 556	1 2 3 6 5	4.35 0.2 2.27 0.1 1.21 0.0 1.24 0.0 0.26 0.0	60 50 77 69 17		1.3 0.9 2.0 2.0 1.6	0.2 0.7 0.7 0.7		1.76 1.05 0.98 1.16 0.86	1, 26 0, 72 0, 74 0, 95 0, 51				0.90 1.04 1.20 1.47 1.55	1.18 1.50 1.61	0.026 0.008 0.007			35.3 22.1 22.3	16.6 14.1 9.0 8.4 6.5	0.14 0.13 0.13
SIMPLE NO.	azn no.	(- #H40AC CA H 5B4A 5B 6#2B 60	EXTRACT: G Hi 41 5841 2D 6P2B	ABLE B K 5891 602E	ASZS -) SUM DASES -HEQ	1CID- 1TY 6828	EXTR AL 6G1E	SOB CATS 5131	-CEC #84- OAC 5484	BASTS + AL 5A38	AL SAT 5G1	-BASE SUM 5C3	S1T- #84 O1C 5C1	C1C03 <288 6218	RES. OHES /CH 881			(PH - CACL2 .01M .01M .021E .1:2	H2O BC1A 1; 1
78 552 78 553 78 554 78 555 78 556	1 2 3	8.3 2 2.9 0 0.6 0	.2 -	0.5	11.0 3.9 0.7 0.7 6.0	19.7	1.3	30.7	23.2	12.3	11	36	47						4.3	5.1 5.1 5.3 5.3
SIMPLE NO.	EZN NO.	(5 (HA PY C F 6A4A 60 <- PCT 0	PODIC HOI ROPHOSPHI F AL 53 6G51 P <2HH->														(0 (< 713	71) TOTAL) RES 781A) L DOM
78 552 78 553 78 554 78 555 78 556	1 2 3 4 5	0	0.1 0.9 0.4 0.3 0.4 0.1 0.1	0.8	0.1		24 1 255 166 342 440						_	KK 1			KK 15			

PARTLY CONTROL SECTION: DEPTS 25-100 PCT CLAY 11.3 PCT .1-7588 67.6. SPODIC HORIZON: INDEX OF ACCUROL 0 ESTIBATED BULK DEMSITY FOR LAYER 1, 3,

AWALYSES: S= ALL ON SIRVED <208 BASIS

SIMBBALOGY: KIMD OF SIMBBAL OR VERSICULITE KE KAOLIMITE BI BICA

RELATIVE ABOUNT 6 INDETERMINATE 5 DOMINANT 4 ANUMDANT 3 HODERATE 2 SHALL 1 TRACE

Example Pedon 2 SALT SHEET

WATDER

SAMPLED AS: CLAYET, MONTBORILLOWITIC, (CALCAREOUS), PRIGID, SHALLOW TYPIC USTORTHENT

5 77ED-001 -008

SAMPLE NOS. 782 775 - 780 DESATURATED HYDRAULIC COMPORTIVITY STORY

DATE 6/29/79 379

U. S. DEPARTMENT OF AGRICULTURE SOIL CONSENTATION SERVICE MATJOWAL SOIL SURVEY LABORATORY LINCOLM, MEDBASKA, AND KORTH DAKOTA AGRI. EXP. STA., YARGO,

GENERAL METHODS 1814, 211, 28

		-1	-2-	1						_									MI. EXP		
						-5		,	-8	-9	-10-	-11-	-12-	-13-					-18-	- 19-	- 20 -
					(-TOTAL SILT)	(c	LAY) (~ -5	ILT	1 (-SAND-		_) (~COL		ACTIO	5 (8H) -) (>2HH)
SABPLE	HID	DEPT	H	BORIZGE	LT	. 002	05	1.			COMMI	4 17			Ç	*C		42	IGHT -		3 T
NO.	FO.	(CH)			.002	05	-2	.0002	.002	÷. 02	.u2 ~.05	. 05	. 10	- 50	. 5	1	2.	5	20	. 1-	PCT OF
					<				- PCT	OP <2	MH (3	U		30	-1	2	-5	-20	~75	75	HOLE
78 775		•			_							,				,	(- P	CT DP	<td>791} -></td> <td>SOIL</td>	791} ->	SOIL
78 776		0- 10-		16	45.8	46.6 45.9 51.0 50.0 54.7 43.5	5.6			33.9	14.7	2.8	1.4	0.5	0.7	0.9	TR			,	73
78 777		23-		AC C1	52.4	45,9	1.7			35.2	10.7	0.7		0.3	0.3		TR				TR
78 778		46-		CRI	40.1	51.0	0.9			37.8	13. 2	0.2	0.2	••			TR				TR
78 779		56-		CR2	45.2	58.7	0.1			41.8	8. 2		0.1				TB		-	**	
78 780	65	81-1	40	CR3	56.3	43.5	0.7			12.6	72.0	0.1	0.1			••	TE			TR	TR
							***			12.0	0.9	0.1	0.1				~~	••		TR	
		OPGY	7051																		
		c		L ZITR	5	(D	TRACTA	T)	(RATIO	D/CLAT	(ATTE	BBEBG 1	(- BOL	E DEES	ITT -)	COLE	{ - - ·	-WATER	CONTE	ST	TRD
SAMPLE	EZW					y R	17						FIALD	1/3	OTER	# BOL 5	LIELD	1/10	1/3	15	BHOLE
FO.											B P 1	4.5	BOISE	NAU N	DET	SOIL	MOIST	BAR	BAB	BAR	SCIL
		<		PC1	07 (288		>			PCT	<0.0BB	ζ	G/CC -	>	CHICK	4 84	- B1C	481C 08 4384	482	47.1
78 775		1.42																- 10.	Ur \200	,	- 0/00
78 776		0.58							0.82	0.37	52	27		1. 17	1.46	0.077			27.9	17.1	9.13
78 777		0.22							0.73	0.34				1.48	1.83	0.073			26.9	17.7	0.14
78 778		0.16								0.36		36		1. 97	1.76	0.062			24.9	17.4	0.11
78 779		0.13								0.37				1.53	1. 95	0.065			29.3	18.6	0.09
78 78)	6	0.12	0.03	0						0.35		34		1.59	1.00	0.054			23.4	16.9	2.10
														1. 33	1. /3	0.037			25.6	19.8	0.09
	(- 1184	OVC	EXTRACTA EA A 584A D 692B	BIE BI	SES -1	ACID-		(C)	·	PICE										
		CY	H G	E A	K	5 0 H	ITT		SUN	NE 4-	MA	348	51	ATION	CARBO	STATE		24 45 2508	()
SAMPLE	821	5B41	5B4	1 5841	5 B4 A	B1585			CLTS	OIC									PASTE	CAC12	
wo.	DO.	6 H 2 H	602	0 6928	6Q2B		6 H 2 B		5434	5161	5D2	52	5C3	5C1	621P		6F1C	6910	8011	9017	
		ζ			BEC	/ 100	G			·>	PCT		<p< td=""><td>CT- ></td><td><p< td=""><td>CT -></td><td><f< td=""><td>CT -></td><td>0015</td><td>1:2</td><td></td></f<></td></p<></td></p<>	CT- >	<p< td=""><td>CT -></td><td><f< td=""><td>CT -></td><td>0015</td><td>1:2</td><td></td></f<></td></p<>	CT ->	<f< td=""><td>CT -></td><td>0015</td><td>1:2</td><td></td></f<>	CT ->	0015	1:2	
78 775	1		6.9	5 0.1	1.3					37.6										•••	
78 776	2		6.6		0.7					3/.3	IR	TB			2				7.7 7.5		
78 777		28.0		6 0.1	0.6	36.3				36.6	70	TH		0.0	2						
78 778	4	19.0	12.0		0.6	31.9				36.5	• • •	7.		97					7.5		
78 779	5	10.3	17.	0.4	0.6	28.3				33.5	•			RA.					7.7 7.8		
78 780	•	5.0	15.	3 0.5	0.5	36.3 31.9 28.3 21.3				27.5		5		77					7.5		
											_										
	(47	THE EX	TR ACTE	PROR	SATUR	ATED P	ASTR-			1	1)
		CA	MG	PA	ĸ								ELEC.								
SABPLE	HZF			· -	-				304	703	0.20	957	9111	(- 1-	HAT -)	(D7	A)	TOTAL A	BIE: JAN
¥0.	WO.	6 # 1 B	601	6 6 7 18	6Q1B	611B	6J1B	6 K 1 C	6L1C	6810											
		<			m	20 / L1	TER -			>	<f< td=""><td>PCT -></td><td>/CH</td><td><- B #1</td><td>ATIVE</td><td>1000</td><td>7840 75 -></td><td>'A3</td><td>/43</td><td></td><td>DC7A</td></f<>	PCT ->	/CH	<- B #1	ATIVE	1000	7840 75 ->	'A3	/43		DC7A
78 775																					- • •
78 776	ż	25.9	7.	5 0.4 3 0.5 3 0.7 7 1.5 9.7 3 23.6	0.4		J. 1	TH	J. 1		60.0	79	0.60	HT 4	ar 3	KK 3	CL 2			2.0	4.6
78 777	ž	34.4	9.	0.7	0.4		1.4	0.1	38.2	-	57.0	0.1	2.60		1		~ ·			_	
78 778	•	21.0	15.0	1.5	0.5		1.2	TB	67.6		68.0	0.1	2.80	aT 2	ni 3	KK Z	QZ 2			1.5	4.1
78 779	5	21.0	30.0	9.7	0.4		1.0	TR	61.2		71.0	0.2	4.20								
78 780	6	18.0	35.5	23.6	0.4		0.7	TE	90.0		79.0	0.3	5.50	5T 4	82 3	KK 2	CL 2	KK 13		2 7	4.6
														-							0

PABLLY CONTROL SECTION: DEPTH 25- 46 PCT CLAY 48.1 PCT .1-7588 0.7

AMALYSES: S= ALL ON SIZYED <288 BASIS

BIBBBALOGY: RIED OF BIFFRAL BY HOSTBORKLL HI HICA KE KAOLIEITE CL CELORITE QZ QUARTZ

RELATIVE ABOUNT 6 INDERFRANIPATE 5 DOMINANT 4 ABOUNDANT 3 HODERATE 2 SHALL 1 TRACE

Example Pedon 1

Pedon Classification: Fragiochrept; loamy-skeletal, mixed, frigid

Series Classification: Same Soil: Series not designated Soil Mo.: 377191-031-001

Location: Cook County, Minnesota SM4, NE4, NM4, Sec. 7, T. 62 N., R. 2 E. about 50 feet north of curve in road, 1,190 Feet south and 2,100 feet east of northwest section corner, about 9 miles north of Jrand Marais.

Climate: Some characteristics of temperatures are: annual normal 360 F (20 C), winter normal 110 F (- 120 C), summer normal 63° F (16° C); some characteristics of precipitation are: mean annual 29 inches (74 cm.), June to September 12 inches (30 cm.), mean snowfall 65 inches (165 cm.)

Parent Material: Grayish brown loamy-skeletal till of the Rainy Lobe of the Late disconsin glaciation.

Lancscape Setting: Sampling site is on the upper part of a 3 percent southwest facing convex slope that is on the side of a ridge about 1,000 feet long and 150 feet wide. The elevation is about 1,600 feet. The local relief

Vegetation: Mainly white spruce 3 to 14 inches d.b.h. with few aspen 2 to 3 inches d.b.h. Understory is mainly hazel

Prosion: None

Permeacility: Slow to moderately slow.
Ground water: Water table is probably perched above the fragipan for short periods.
Described by: R. Lewis October 13, 1977.

Sampled by: S. Holzhey, M. Mausbach, R. Turner, H. Finner, D. Grigal and J. Engen.

- Al 757572 O to 10 cm. (O to b inches). Elack (10YR 2/1) cobbly loam; strong fine granular structure; very friable, slightly sticky, and nonplastic; few worm casts; many fine and very fine continuous pores; many very fire to coarse roots; about 15 percent coarse fragments; about 5-10 percent larger than 5 cm.; clear smooth boundary. (3 to 13 cm. thick)
- 10 to 13 cm. (not sampled) (4 to 5 inches). Discontinuous horizon occupies about 10 percent of pedon; reddish gray (5TR 4/2) loam; moderate very fine and fine granular structure; friable, slightly sticky, and nonplastic; abrupt broken boundary. (2 to 5 cm. thick)
- 921hir 79P553 10 to 23 cm. (4 to 9 inches). Dark reddish brown (5YR 3/3) cobbly loam; brown (7.5YR 5/4) dry; moderate very fine and fine granular structure; very friable, slightly sticky, nonplastic, and smeary; many fine and very fine discontinuous pores; many very fine to coarse roots; about 15 percent coarse fragments; about 5 to 10 percent are larger than 8 cm.; gradual wavy boundary. (10 to 20 cm. thick)
- B22hir 78P55h 23 to 36 cm. (9 to 15 inches). Dark reddish brown (5YR 3/h) cobbly loam; moderate fine granular; very friable, slightly sticky, nonplastic, and smeary; many fine and very fine discontinuous pores; many very fine to coarse roots; about 10 percent coarse fragments; about 5 to 10 percent are larger than 3 cm.; gradual
- 737555 38 to 63 cm. (15 to 25 inches). Discontinuous horizon occupying about 15 percent of pedon; brown (7.5YR u/L) cobbly leam; moderate fine subangular blocky structure parting to moderate fine granular structure; friable, slightly sticky, nonplastic, and slightly smeary; common fine and very discontinuous pores; common fine and very fine vertical roots; about 15 percent coarse fragments; about 5 to 10 percent are larger than 8 cm.; clear wavy boundary. (0 to 25 cm. thick)
- 799556 63 to 122 cm. (25 to 48 inches). Dark grayish brown (10YR 4/2) cobbly loam; weak medium and coarse platy structure; firm, and brittle; few very fine horizontal pores; few very fine and fine rocts in upper part, dominantly between plates; about 45 percent coarse fragments; about 35 percent greater than 3 cm.

Remarks: Colors are for moist soil. Backhoe was used for the sampling pit. Lithology of rock fragments is mainly gabbro and basalt. Few "red rock" fragments.

Example Pedon 2

Soil Type: Wayden silty clay. Sample No.: S77:ID001-4 (FIPS) Date: September 12, 1977. Sampled by: M. Mausbach, F. Schroer, M. Meyer. Described by: F. Wilhelm, G. Muckel, S. Ekart, E. Gamble. Area: About 2 miles east and 213 miles north of Reeder, Adams County. Location: 300 feet east and 330 feet north of the southwest corner of the :Wig. Sec. 30, T131N, R95N. Classification: Clayey, montmorillonitic (calcareous), frigid, shallow Typic Ustorthent. Vegetation: Summerfallow (spring wheat in 76). Parent Material: Clayey shales of the Ft. Union (Tongue River) formation. Physiography: Hilltop. Relief: Convex, simple. Slope: 2 percent. Permeability: Slow (0.06 - 0.2 in./hr.). Moisture: Moist throughout. Drainage: Well drained. Ground Water: >5 feet. Erosion: Slight. Aspect: Northeast. Elevation: Approximately 2,900 feet. Pedon Description: Wayden silty clay. (Colors are for dry soil unless otherwise stated.)

- Grayish brown (2.57 5/2) silty clay, dark grayish brown (2.57 4/2) moist: strong very Ap 0 to 10 cm. fine granular structure; very hard, firm, very sticky and very plastic; common very fine roots; many very fine pores; slight effervescence; abrupt smooth boundary.
- Light brownish gray (2.5Y 6/2) silty clay, grayish brown (2.5Y 5/2) moist: moderate very fine subangular blocky structure; very hard, firm, very sticky and very plastic; common very fine toots; many very fine pores; tengues of Ap less than 5 mm thick; common irregularly snaped soft like messes; slight effervescence; clear wavy boundary.
- C1 23 to 46 cm. Pale yellow (5Y 7/3) silty clay with 30 percent scale fragments, clive (5Y 5/3) moist; moderate very line subangular blocky and rock structure; very hard, very tirm, very sticky and very plastic; common very fine rects; cormon very fine pores; common medium yellowish brown (10YR 5/8 mois*) from stains; common gypsum crystals; common pressure faces; tongues of Ap less than 5 km thick; clear wavy boundary.
- Crl 46 to 56 cm. Pale yellow (5Y 7/3) clay shales, olive (5Y 5/3) moist; rock (fissile) structure; extremely hard, very firm, very sticky and very plastic; common very fine roots with most following fissures and cleavage planes: few very fine pores; common medium yellowis; brown (194R 5/9 moist) iron stains; few tongues less than 5 cm wide; common pressure faces; common gypsum crystals; clear wavy boundary.
- Cr2 36 to S1 cm. Pale yellow (5Y 7/3) clay shales, olive (5Y 5/3) moist: rock triasite/ stronger, extremely hard, very firm, very sticky and very plastic; few very fine roots following fissures and cleavage course for some firm some firm for the form (7.3v 4/L moist) from stains; common gypsum crystals; planes; few very fine pores; common medium dark brown (7.5Y 4/4 moist) from stains; common gypsum crystals; common fine paleoroots; common pressure faces; gradual wavy boundary.
- Light gray (5Y 7/1) clay shales, gray (5Y 5/1) moist; rock (fissile) structure; extramely hard, very firm, very sticky and very plastic; few very fine roots following fissures and cleavage planes; very few fine pores; common gypsum crystals; common fine paleoroots; few pressure faces.

Additional Notes

Large horizontal and vertical cleavage planes extend through the shales. Some are as large as 19 inches by 24 inches. These are coated with brown (10YR 4/3 noist) and some have up to 2 inches of weathered material along these planes. The rock structure size increases with depth. Lignite was encountered at a depth of 55 inches. Also, an Unsatruated Hydraulic conductivity study is being conducted at this site.

Appendix V

This appendix contains a supplemental data sheet for all of the pedons sampled.

SCUADOR 001

SAMPLED AS:

S 330N-325 -001 ECUADOR-5NSS

SATULE NOS. #3P2103 - 2110 DATE 11/30/83

U. S. DEPARTMENT OF AGRICULTURE SOIL CONSERVATION SERVICE NATIONAL SOIL SURVEY LABORATOMY LINCOLN, WEDBASKA

GENERAL METHODS: ENGINEERING PRACTIONS JURY CALCULATED PROBUSDA FRACTION SIZES

	GEN	SHAL :	1 et h	1005;	eugi	aeda:	I DH	PRACT	I ost	ara	13 34	LCUI	ATEI	PRO	H US	DA	PHACT	C HOI	IZE	s			,	20020	-		
SA INGE	112 N NÚ.	(IN)	 !	 -201 1021	9 P 8 3 <	H C	G 1 E N 1/2 N C	T A 1 H E	E G € 3 /4 S —•	3/il	1 L 4 4 (-4	5 S 10 U (I	น เม 40 2 ย	G 200 8->	s s 20 <-d1	I E S CRO	A V R 2 NS->	COM USDA 1	ULA.	TIVE LESS .25 -HH-	CUBVE THAN .10 .05	PRACT DIAM 60 (P	IONS (< ETERS (50 ERCENT	ILP>	<-PCT	CD C	CC
d30103 d30104 g30135 d30130 g30117 g2010d g30119 g30110	45 55 75	a- 1a- 29- 33-	18 29 33 45 59 72	12 201 3431 3452 3405	100 100 104 100 100 100	100	98 37 73 97 98	100 97 79 57 95 97 97	73 40 44	96 70 11 14	96 67 36 36 75	49 13 16 13 14 15 12 13	42 32 31 41	1	10 1 11 29 27	16 16 16	5 1 1 k 5 7 3	90 52 19 91 93	44 11 95 86 34	35 3 75 75 71	51 37 26 17 1 60 45	7 0.16 7 2.42 127.12 5 0.10 9 0.12	0.097 0.829 21.034 0.063 0.074	0.004 0.004 0.020 0.468 0.003 0.003		42.0 > 100 57.9 33.5 43.5 58.9	1.6 1.6 0.5 0.7 1.2 1.4 1.9 2.0
\$4 10 LB	.12	>2 .:	10 5 2 0 5 1 0 H	1 6 50 7 77 - 67 3F 23 2	5 5 75 2 ~20	1 6 20 6-	-3 2	1.1) -	75 -2	' ·	1 1 15 • • • • 5	.2: I∂ 5 -2).a== 		3014 3014 1/1	50 0	AUN Hara Suops	ESGIL ESGIN ESICE	EER!	ING TUR	50 I I	-<2 M SURV 15 BAR	H PRAC EY E OVEN		RING I	T 1/3	105 BAR <2
83 193 33 134 93 (195 83 21 6 95 197 93 193 13 21 19 83 (115	1 2 3 4 5 6 7 6	75 7		1 7 	1 4 7 4 2 27 5 54 7 6 5 4 1 4 7 4	78 78 15 16 1	1 1 1 1 1 1 7	25	1 7 9 7 1 1 5	:7 :1 :0		3 9 11 1	43 24 24 34		1.4 1.2 1.4 1.3 1.3 1.2	9 7 5 4 5 9	1.47 1.34 1.38	1.71 1.00 1.67 1.50	1 1 1 1	.92 .93 .86	1.29 1.35 1.23	1.11 1.29 1.35 1.26	1.11 1.29 1.35 1.27	1.42 1.62 1.65		0.80 0.98 0.92 1.05	1.39 1.05 0.96 1.15
avadu t	112	/ ¥ >2 _5 -1	n (-(() 2 () -	L of	1 F 3 75 2 -20 ic	2 u 1 20 -5 4u 1	5 -2 4101	(53) (53) (2) (8) 50	 	1 7 1 7 22 22) () () ()	9 P) 2 D	5) 1 14 U.3 P	-10 - (CZ) - (CZ)	(1 K	1 0 5 C C SUM 	T PR	O ACT	c	BAR	LINR WHOL <-1/5	AR EXT	< U (IN/ U (IN/	LITY) 2 HH LW) ->	V B HOL SUIL <in< th=""><th>D) E <2 BB</th></in<>	D) E <2 BB
8 (11) 83 114 93 115 13 116 33 2107 93 2101 33 7109 93 2110	2 3	3 .	 	4 2	1 3 1 3 15 4 3 4 3 3 2 4 2 4 2	7 h 2 h 3 3 T7 1	1 1 5 4 1 1 1 1 1 1	77 37 96 97 96	23 20 24 28	17 15 1 20 11 14 16	1	2.1	26 25 12 23 31 45			1	7.13	4.56	1 - 2 -	.72	0.100 120 079		1.0	0.4	1.1	0.17 0.18 0.25 0.21 0.20 0.20	0.23 0.26 0.2 :
S CAPUS NO	42 40)2 75 4 1 >2 75 	? I ! 3 5 2 -2)F >	G H T L E -2 -0 -214+3 73 7	P S O I .05- 5 .00 AND+S 9 80	R A C L LT 2.002 LLT> 81	YC (0 N -<2 H 51	5 - N F N DS- 1 P CT 39	C L R A OP 3		٥			1-20	. 2011	651				3)) (CLAY LT C .002 - 74.1-)	011 1 4	- D1 1111 P	BICAL) CON- UCT. MHHOS MH- 37	(CUMULI 5ALT HG/ KG 15)	C. AMO IN. O	P H2O R TO IRDRY SOIL)
3 32 10 3 4 32 10 4 8 32 10 5 5 JZ 10 6 8 32 10 7 3 32 10 8 3 32 10 3 8 32 10 3 8 32 10 3	1 2 3 4 5 6 7 6	75	1 7 13 75 7 5 4	1 6 3 5 15 4 21 2 1 5 1 5 4 5 3 6	9 33 2 15 4 1 1 42 7 38 9 32	5 5 1 7 6 7 9	2 4 10 24 2 2 2 2	7 9 14 34 7 7 7	15 15 16 30 11 13 15	22 20 17 7 17 22 22 22	17 16 16 1 17 16 13	15 15 12 2 16 15 11	15	5 5 3 1 6 4 10 7		1	PSL PSL St. COS PSL PSL PSL PSL	51.7 62.7 71.4 95.5 51.0 53.3 53.1	34 25 42 31	1.3 5.1 3.7 7.3 5.9	4.9 5.0 2.5 0.8 6.1 7.3 9.2 6.3	5.2 6.0 5.3 5.7 6.4 6.1 6.4					
SAMPLE 332103 832104	0 P	LOW - 21	LO¥ 21	DIP 21 19	CLAY 4.9	5 A N	D 7	oc 2.65 2.33	7	5/C 12.2 12.1	¥ 1	3/C	ı								(F 3/ 1 12.2		s/c 2	와 OC 2.7		-	

ECUADOR 002

SAMPLED AS:

S 93PH-325 -002 ECUADOR-SHSS SAMPLE MOS. 83P2191 - 2195

DATE 11/30/83

U. S. DEPARTMENT OF AGRICULTURE SOIL CONSERVATION SERVICE NATIONAL SOIL SURVEY LABORATORY LINCOLM, MEBRASKA

GENERAL METHODS: ENGINEERING PRACTIONS WERE CALCULATED FROM USDA PRACTION SIZES

	GERI	511 4 5	13 1. ()	,,,,,		T II			inne i	, on.	***									-							
SA IPLE		43C		((O)		Р E 3 <	2 [2 N 3/2 N C	T,A	G 3 3/4 S	3/8	₽ A 4 <-N	S S 10 U H	1 8 40 8 E	G 200 R->	S I 20 <-MIC	P; V 2 5 2 RONS~>	USD 1.	.5	LESS .25 -8 K-	CURVE THAN .10 .0	DIA: 5 60 5 <:	eters 50 Percen'	(57) AT 10 Tile>	LL P LL P C-PCT	ONI- I PHTY > CO	VTUH CC
837191 832192 832193 832194 832195	25 35 45	0 - 21 - 35 - 43 - 47 -	. 35 . 43 47	3.2 3.3 20		100 100 100	100 99 33	100 99 79	98 66	100 98 57	100 98 55	10 0 9 8 5 2	100 97 49	99 94 46	57 29	4	3 1 4 1 2 1	100 96 48 D	100 94 46	98 91 45	76 3 d0 2' 73 2' 37 1	9 0.00 9 0.00 620.9	8 0.06 8 0.076 1 2.65	7 0.023 0 0.023 9 0.036	3 1 3	3.4 3.9 >100	1.4
SAME	H 2	<	50 UP	0 L : 250 -75 Pct (75 -2	5 75 -20 Hole	0 I 20 -5 50I	L (1 -2 L	18) - <2 >	-<: 75 -2 <	75 - 75 - 20 - 20	1 294 20 5 0 2 4	CTIC 5 -2	N <2 in->		SOIL 1/3	- D&& O A S W I S A S B S S S S S S S S S S S S S S S S	T R SOI R SIOR SIOR	P E I L HEER! I SA!	R ING TUR TKD	SOI 1/3 BAH	T <2 L SUR 15 BAR	VOL	U M E CTION ENGINE! BOIS!	G\CC) (VO BAT AT 1/3 	ID) IOS BAB <2
832191 832192 832193 832194	j	гя 3 51			3	2	TH	TR 1	100 100 97 49	3	 2 4 J	Th	CR 1	97		1.46 1.44	1.47	1.7	1 1	.91	1.40 1.46 1.41 1.40	1.47	1.47	1.71	1.91 1.88	J.62 0.34	0.82
SVILE	a •	>2 2	V Н 50 ОР	0 L 250 -75	2 75 -2	75 -20 -901	0 1 20 -5	L L 5	(4.1) (2 Le 50	2- 2- 01 L-	.05 .00	00 - rt	P L	> burz h	- (/! [4] -[1]	K)	(2) SJ t CA PS	4 4 4 5 5 5 6 6 6 6 6 6 6 6 6 6 6 6 6 6	RACT - C	10 H- 15 BAB H20	1/3	VIIO <- 1/ 15 Bar	LE 301	TO (IN, 15 BAR	C2 MM	SOIL	.E <2 HH <-WI\\
83,191 81,192 81,193 81,194								T9 1	100	19 30	16 15 15 11	1	20 13 9	27 25 43 22	12 14		7.35 3.50 12.00 24.00	6.7 6.7 9.2	10 4 12 5 15 4	.70 .89 .75	0.200 0.556 0.583	0.5 0.4	0.5	u.5 0.5	0.5 0.7	0.18 0.20 0.17	0.26
SAMPLE NO		>2 >2	4 () 75 −2	20 -2 >24H	.05	0 [-05- -002	L LT 2.00	 2 7C	-<2 5 5 C	1M N DS: N -P CT	P R S	TP	SIL C SIL) H TS F [CL AT	(13C) (11 (11)	941869 87 PSD: 2 84-1	2 - 3 - 05 1 - (.0 .0 .PCT	\(\)(\) 1LT 5- 002	CLAT LT .002 .2Md-)	CT5 CT5 O 14 ((-TLEC ATS- IST. OHMS	TRICAL COM- DUCT. MMHOS) (CUHUI SALT HG/ KG	T. AH. In. (In. (DUNTS) DF H2C AR TO AIRDRY
8 3219 1 8 3219 2 8 3219 3 9 3219 4 9 3219 5		, 51		1	72 68	29	1 1 1	1		3		52 45 43	22 21 22	9 10			LVP: VPS! VPS!		.3 2 .7 ! .4 3			7.0 7.4 7.6 8.5					
SAMPLE 8 32191	U P	LOW	Ln 4	W DT	7 (0	LAY 0.9	9 8	N D	oc		5/0	c 1	5/0	¢	0.6			UP LC 0- 5)¥ L	,0¥ [017 S	/C 7 76	V S/C	₩ OC 0.6	+		

ECHADOR 303

SAMPLED AS:

2CUADOR-SMSS

S 43FH-325 -003

SAAPLN # . 83P2200 - 2204

DATE 11/30/03

U. S. DEPARTMENT OF AGRICULTURE SOIL CONSERVATION SERVICE WATIONAL SOIL SURVEY LABORATORY LINCOLN, WEBBASKA

GENERAL METHODS: ENGINEERING PRACTIONS FORE CYLCULATED FROM USDA PRACTION SIZES

	GEN	CHAL	ETI	ods:		NGIN	THEF	NII S	PRACT	11015	261	E C	LCUI	LATE	o enc	OH US	SDA	Phace	ION 5	3128	S							11.31	
SAUPLE TO.	H2 Y			101	(-	PE	2 I	8 N 3/2	1 H E		3/8	P 1	3 5 10 U 1	H E	G 200 H->	3 20 <-41	1 E	1 NS->	1	.5	.25	. 10 .	N DI.	HETE 5 -PERC	BS (E	TA (HE	ATTERBERG LL P: <-PCT: 22 2:	UNI- FHTY	VTUR CC
332200 632201 832202 832253 632254		0-	54	A1 A2 13 14 2A5			100 100 100		100	100		100 100 100	100	91 96 44 10 17	51 55 44 43 66	16 17 13 17 29	7 8 9 7		99 99	93 97 96 94 99	85 91 87 79 93	66 59 52	39 0. 35 0. 31 0.	09 0. 10 0. 13 0.	066 078 093	0.008 0.007 0.006 0.007 0.004		16.2	1.7 1.9 2.2 2.3 1.7
SATELR		>2 2	# C 50 2 UP -	50 -75	75	75 -20 HOLE	20	-2 L	1:1) -	75 -2	75 -20 -00T	20 -5	1071 3 -2	<2	3.7	5011	50	RV EY VEN	T SOI!	EER SA	TNG		<2	RVEY-	RACT - EN	TELON SANIDA 12101	RING	-RAT	IOS
8322 0 9322 1 9314 2 8322 13 8322 14	1 2 3 4 5	IR IR	 		=======================================		=======================================	TR	100 100 100 100 100	11111			10	100 100 100 100 100		1.1	13 25 27	1.22 1.31 1.32 1.13	1.35	5 1	1.49 1.70 1.78 1.79	0.79 1.13 1.25 1.27 1.03	0.9 1.2 1.3 1.3	0 1. 0 1. 1 1.	98 22 31 32 13	1.32 1.56 1.60 1.59 1.49	1.49 1.70 1.78 1.79 1.64	2.35 1.35 1.12 1.09 1.57	2.35 1.35 1.12 1.09 1.57
SAMPLE		>2 2 <	50 : UP -	50 75	15 75 -2	75 75	-5	5 -2	(AII) (2 LE S	2- .05	.05.	L C . 00	a \ 12 D	K>	RAT	1)	i K k Y	1 0 : <2 C SUM CITS	113 PF R C MH4- OAC 67	ACT	15 TON-	L A Y LE 1/3 HAR 6)	- WIS	0LE S /5 HA ()V R -0	OIL R TO EN		LITY) 2 HH IN) -> OVEN -DRY 73	SOIL	E <2
132 0 d32 11 932 2 d3, 13 132, 4	1 2 3 4 5	TR TR TR	==		=======================================		=======================================	TS TA	100 100 100 100 100	19 26 31 33 20	11 16 16 14 19	1	17 16 18 20 15	35 32	10		1	1.27	5.4.	3 5	.87 .08 .03	5.692 1.733 1.231 1.625 1.722	5. 2. 1. 1. 2.	0 2 3 1 0 1	.4	5.6 2.0 1.J 1.0 2.5	7.4 2.6 1.6 1.3 3.1		0.37 0.33 0.28 0.26 0.34
SAMPLE	HZ NO	(V W >2 7 PCT 76	11 0 5 7 0 P 2	L E	S	n I 05- .002 D+SI	L LT .002	VC (-<2 : s: c	H I	POPS	VP	SIL C SIL	75 7	CL	IN PIEI	PRRA LU -<2	RE) BY BY PSDA HM-)	3A41	or.	11LT	LT .002	CL2	IST.	5 /2	UCT.	KG 1	IN. C 1/3 BA 5HAR A (WHL	R TO
8 32200 8 32201 9 32202 3 32203 8 32204	1 2 3 4 5				63 62 66 70 51	37 38 34 30 49	1 2 1 1 2	1 1 1	3 2 3 6 1	5 5 7 15 6	25 26 29 27 17	23 27 24 22 27	22 22 15 14 21	16 17 16	1 2 1 1 2			VPSL VPSL VFSL VFSL VFSL	62.2 65.0 65.0 50.2	1 1	16.5 17.7 13.3 10.0	1.3 1.5 1.3 0.3 1.8	4. 5. 5. 5.	3 6 7					
SAMPLE 932200 832201	0	LOW - 15 - 54	101 11	5 19	5	LAY 1.3 1.5	5AN 62. 60.	.2	OC 7.8 2.9		3/0 17.5 40.5	3	4 5/ 47.A 43.3		7.8 4.3			(JP LU 0- 15 5- 54		15 1 11	5 47	5/C .8 4	7.8	2 13	V 00 7.8 7.0			

SUPPLEMENTARY CHARACTER 12 ATTON DATA

RCUADOR 004

SAMPLED AS:

ECUADOR-SMSS

s 93FN-325 -004

SAMPLE NOS. 8392119 - 2120

DATE 11/30/83

U. S. DEPARTMENT OF AGRICULTURE SOIL CONSERVATION SERVICE NATIONAL SOIL SUNVEY LAUGRATORY LINCOLN, MEBRASKA

GENERAL METHODS: ENGINEERING FRACTIONS WERE CALCULATED PROMISED PRACTION SIZES

															, .	5													
SAIPLE		(I) 180			к- он	,	t		n z	3	,	(- n	וה ט	1 6	パーノ	V11.	~11U F	43-,	(11 17 -		·-, 、		EHCENI	(7,6HM) (HM) A: 10 FILE-	, (エン しし	DATION - CUR- Y YTUR CC 4 25
8 12119	48 58 68 78		- 18 - 25 - 33 - 59 - 66 - 75	203 203 346 304		100 100 100 100 100		81 72 100	69 55 100 100	100	55 53 36 100 130	45 23 100 44	n0 41 34 13 103	17 12 7 96 92 76	67 62 66	37 11	1 TR 17 17	TR TH TK 3	75 32 27 15 100 97 78	65 20 15 8 97 99 72	50 9 40 90 61	34 3 1 1 74 68 51	23 0 119 117 TR28 56 0 54 0 40 0	.39 .00 .41 .62 .06 .07	0.243 4.700 7.273 22.490 0.036 0.040	3 0.00° 3 0.00° 3 0.25° 2 0.36° 0 0.59° 5 0.00° 0 0.00°	9 6 6 9 1 1 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	45. 73. 47. 47. 18. 21. > 10	4 1.3 0 1.8 2 0.2 6 0.3 8 1.9 3 0.8 5 0.7
SAMELS	11.0	>2 3	1 H (0 L 25J -75 PCT	9 75 -2 OF 1	75 -20 HOLE	0 I 20 -5 : 50I	L (; -2 :L	1.1) -	75 -2	75 81 75 -20 -PCT	1 20 20 -5 0! 4	10110 5 -2 175	5) 04 (2 16-1		(4 1	SUN OV)84 (34 (454 (40 F)	102 S 1083 210r	P 8 L N E E B T S A	R ING TUR	0 N 	1 1 > S LL S 15 BA	V 2 Mi URV 1 R	O L L	MOIST	G/CC) (V	TIOS 3 BAR 2 <2 88
812119 832120 832121 332132 832123 832124 832125	3	20 59 66 81			9 1 2	3 11 40 39 57 	15	9 11 9	80 41 34 19	51	39 57	1 3 10 16 15 1	6 9 11 9 	94 80 41 34 19 100 98		1. 16 1.96 2.07 2.37 1.09	5 1 7 1 1 1	1.16	1.5 1.5	1 1 7 1 7 1	1.72	1.02	! 1. ! 1.	06 15 16	1.07	1.40	1.64	1.28	1.60 1.60 1.43 1.43
SAMPLE	H Z	>2 2	-1 H -50 -UP	() L 253 -75	F 75 -2	75 -20 -20	0 I 20 -5 UP	L 5 -2 ≆801	(da) - <2 -8 SC	2- -05 -01 L	.05-	1/1 - LT 2 .00	3 1 20	९)स एड ह >	・ (ノ4 RAT -10	ı	 !	< 2 C SUM	40 P	R ACT - - -	10#- 15 .Ait d20		 -> 1 0	HUL: 1/5 5	E SOIL	TRNSIB: 10 (18, 15, 18, 72	- nn 2) -> (ul) -> OVEN	- WIIO	LE <2 MM W/IN->
832120 932121 932122 932122 932123 932124 932125	1 2 3 4 5 6 7	5 2 7 1				1 5 30 31 50 7	13	1 7 3 	49 29 100	3.0			14 21 25 22 13 11 11									0.305 0.869 0.677 0.667 0.185		.5	1.9 1.4 2.1 2.4 2.1	1.6 1.3 1.4 2.1 1.5	2.4	0.27	0.29 0.29 0.47 0.47 0.27
SANPLE NO	17 80)2 7	E I H O S :	G H L D 20 -2	t 2- .05	PR 0 I 05-	4 C L LT .002	T 1	lon ≺2n SA	5 - 6 8 8 80 8 8	- C L	, A 7 C T	T I O	' E E	CL AY	(-fux (orts in Field	TUK TUK IPP I	150 150 17	(5 0 0 0	1 (1 15 T 5-	CLAY	CL2 CL2 CL3) (* - (* 1 (*	- DLECT RES- ST. C	*RICAL) *CON- *MCT. *MM405 **TM- **97	(COMMI SALT TG/ KG	T. Am IN. 1/3 B 1/3 B	DUNTS) OF HZO AR TO AINDRI
8 32120 9 32121 8 32122 9 32123 8 32123 8 32124 8 32125 8 32126	2 1 4 5 6 7	59 56 91	66 91 2	1 9 19 27 24 2	19 45 46	39 22 1 1 TR 55 52 33	4	2 7 22 20 21 TR 1	30 37 36 2 3	19 12 26 31 33 7 8	2 1 2 1 16 7 7 17 18 18	17 14 3 1 1 18 15	15 10 1 14 19 13	25 17 3 2 1 36 36 30	2 1 1 1 3 4		5 0 0 5 5 4	OL OS OS OS IL	\$1.7 73.7 96.6 97.6 97.6 93.5 45.6	1 3 1 1 1 1 1 5 5 3 3	9.1 7.1 3.0 1.7 1.3 1.4 1.4	2.1 1.8 0.6 0.9 0.9 3.1 3.6	\$ 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	.6 .3					
SAMPLE 832119 832120	UP 0- 24-	LOV - 24 - 45	LON 24	9 II	 Р С Ц	LAY 2.1 1.0	SAN 50. 71.	D 8 1	oc 4.26 2.51	 3 3				7									s/c .0	29.0	s/c	₩ 0¢ 4.3			

ECUADOR 005

SAMPLED AS:

S 83PH-325 -005 ECUADOR-SMSS

SAMPLE NOS. 33P2151 - 2153 DATE 11/30/83

U. S. DEPARTMENT OF AGRICULTURE SOIL CONSERVATION SERVICE NATIONAL SOIL SURVEY LASORATORY

		eral			: 1	ZNGIN	18581	NG I	RACT	'I ONS	421	is ca	LCOI	LATEI	o ∄RO	ព មន្ត	A PR	ACT	ION SI	ZKS					NBBK.		iver L	SURAL	561
SAULE	HZ.1	(I) 06F	*) *TH	110 12	K-	(I	N C	T A 1 H 2	5	>	<- 1	UĄ	nε	k->	<-m10	れのほら	->	<	3	M		> <	PRECE	RITTE	>	ATTER- BERG LL PI <-PCT2 22 2.	GRAD UNI- TATY CU	ATION CUR- TUR
331151 33_152 83_153 33_154 33_155 43_156 6_3157 33_158	25 35 45 55 65 75	0 - 3 - 11 - 21 - 39 - 43 - 56 -	- 11 - 21 - 31 - 39 - 43 - 56	34 24 201 206 348	2	100 100 100 100 100 100	100 100 100 100	100 39 100 100 100 100	100 93 103 100	100 97 100 100 100	100 47 100 100 100	100 47 103 138 198 100	100 17 100 100 100 100	100 46 99 99 99	58 51 56 59 52	J6 26 19 24	12 10 12 10	2 2 1 1 2 2	97 9 100 10 37 9 100 10 100 10 100 10	17 9	9 6 6	79 5 68 4 62 3 67 4	5 0.0 3 0.0 7 0.0 3 0.0	6 0.0 8 0.0 9 0.0 8 0.0	39 0.0 61 0.0 71 0.0 62 0.0	003 004 006		17.9 16.7 18.5 15.1 17.2 18.2	0.9 1.7 2.1 1.8 1.7
SAMPLE	HZ NO	>2 2	250 -UP	250 -75 PCT	75 -2 08 -	75 -20 1110 1. 1	20 -5 : 501	5 -2 L	<2 >	75 -2 <	75 20 - PCF	23 -5 07 (-2 :75	<2 111->		SOIL	SURV OVE	H H Y	ENGINE	ERIY Satu -ate	G H D	50 I 1/3 8A B	L SUK	VEY-	HOI	95 % H	G/CC I ING I ATUR ATED 48	LT 1/3	916
33:151 33:152 33:153 83:154 84:155 94:156 83:157 33:153	2 3 4 5 6 7	3				3		TR	97 100 100 100 100 100] 	3		cn 	100 97 100 100 100 100 100		0.73 0.86 0.06 1.05 0.66	0. 0. 0. 1. 0.	97 95 97 95 97 99	1.14 1.35 1.30 1.32 1.40 1.22	1.4 1.5 1.5 1.6 1.4	5 4 3 5 5 8 7	0.73 0.34 0.85 0.85 0.89 1.05 0.77	0.82 0.93 0.93 0.96 1.08 0.81	0.8 0.9 0.9 1.0 0.8	4 1. 5 1. 5 1. 7 1. 9 1. 9 1.	14 33 30 32 34 22	1.50 1.45 1.52 1.53 1.55 1.65 1.48 1.37	2.63 2.08 2.12 2.01 1.52 2.44	2.63 2.15 2.12 2.01 1.52 2.44
23 / LB	HZ NO) 2 2	0 2 H 50 UP	L U L 250 -75	0 3 P. 75 -2	75 -20	20 -5	2 - 2 - 2 1 - 11 1 - 11 - 11 - 11	- X (H.II) - <2 .≳ S∪	1 2- 11) L l 	3 A P: 2 D	S) R 95 ES >	(C/) (/W 6AT -EU	(il) PIHI CLAI	T L S CA	0 S < 2 - C U 11 T S	67	CT IO 15 BA H2 6H	C L H B O	LE 1/3 9Ak 69	(LIN WHO <-1/ 15 15 DAR 70	EAR P. LE SO 5 BAR OVE -DR 71	XTEHS. IL TO (. H 1! Y B.	1 0 LL <2 1 H/I 5 A B	.ITY) ! nu !nj-> oven -dry 73	W B WHOL SOIL CIN 74	/ LB-/
8 32151 8 32152 8 33153 8 33154 8 32154 8 32155 9 32156 9 32157 9 32158	3 4 5 6 7	1			1	1		TR	99 100 100 100 100 100	12 18 20 19	17 15 13 11 14 19 10	1 1 1 1	20 31 19 23 23 25 25 30	49 41 49 45 44 35 45	12 11 11		15. 13. 16. 20. 16. 9.	30 35 53 54 64 74	5.90 6.00 7.4. 1.25 6.72 1.33	5.2 6.5 3.2 10.1 10.0 5.4 9.5	8 1 0 2 4 2 7 3 0 3 2 3 0 2	.375 .400 .471 .167 .000 .634	3.4 3.4 3.0 2.4 0.1 3.3	5. 4. 3. 3. 1. 4.	4 4 4 4 1 3 d 3 2 2 3 U 9 J 10	.0 .5 .0 .9 .8	4.3 4.2 J.8	0.35 0.34 0.23 0.29	0.31 0.38 0.35 0.34 0.23
SAMPLE NO	4 Z 9 O	>2 7	11 O 15 -2	L 5. 20 -2	2- .05	0 I 05- .032	L LT .302		≺2 //1. SA C	# 5 # DS = #	R A	СТ V?	SILT C	אנ כז פ	CL AY	(UST" IN FISLE	41 67 70 24	P D P A	1840 2- .05	5 tL -05-	T 2	CLAY LT .002	CA- CL 2 - 0 1/1	RES- IST- ONNS	CON- DOCT	- . H	(CUMUL) SALT IG/ KG 19) 98	IN. U AC E/I A RAB	P H20 R TO IRDRY
832151 332152 932153 932154 932155 832156 832157 832158	1 2 3 4 5 6 7 8	3	3		46 55 64 58 55 64	57 54 42 36 42 45 36 16	4 2 2 1 1 2 2	TR	TR TR TR TR	1 4 4 4 3 4 15	19 21 26 34 30 29 34 44								39.4 45.0 55.3 63.2 57.2 51.3 62.3 93.3			2.0 1.7 1.2 1.1 1.9 2.2	5.7 5.6 5.4 5.3 5.4 5.5 5.5						
SAMPLE 832151 832152 832153	0- 7-	LOW 7 27 54	2	7 7 2	7 0	4.0 2.0	39.	D 4 0 8	0C 5.71 2.69 2.22	3 2 7	5/C 9.8 22.5 32.8	1		: 2					1 LOV - 7 - 27										

SCUADOR 006

SAMPLED AS:

S 83PN-325 -006 ECUADOR-SMSS SAMPLY HOS. 9302065 - 2070

DATE 11/30/83

U. S. DEPARTMENT OF AGRICULTURE SOIL CONSERVATION SERVICE HATIONAL SOIL SURVEY LABORATORY LINCOLM, NEBRASKA

GREERAL METHODS: ENGINEERING PRACTIONS FREE CALCULATED PROBUSDA PRACTION SIZES

			2100	٠, ,					20.13						., ., .,			1011 3	12.0	-								
SAMPLE	HZN NO.	(IN) Dept		 09- 404	9 q £	R C	ያ ዘ 3/2	T A	G P. 3 ∕4	3/1	P A 4	S S 10 U H	1 H	G 200 R->	5 1 20 <-#10	E V S Kons	2 ->	USDA 1	5	LESS .25 -ah-	.10 .	M DI 05 6 -> <-	amet O -per	ers (50 Icent	10 (nd	ATTEN BERG LL P <-PCT 22 2	UNI- I PHT! > CU	CUR- VIOR CC
8 32065 8 32066 8 32067 8 32068 8 34069 8 32070		4- 22-	59 2A	2 3 41 92	100 100 100 100 100 100	99 38 99 99	99 99 90 99 99	98 98 98 98 90	47 97 60 99 97 87	97 97 59 98 97 67	47 97 57 98 97 86	97 97 50 98 46 85	90 90 42 92 91 70	65 67	37 16 36 36	17 17 7 17 16 12	5 4 1 2 2	96 46 97 95	92 92 43 94 92 79	84 38 87 87	68 30 71 74	53 0. 2319. 56 0. 55 0.	07 0 00 2 06 0 06 0	.043 .000 .036	0.003 0.003 0.003 0.003	 	22.3 > 100 18.9 17.7	0.7
SA ALE	нZ	(¥ ¥: >2 25: -0' < 25 2'	H O L D 250 P75 PCT	75 -2	5 75 -20 MOLE	0 I 20 -5 SOI	-2	(1) - (2 >	-<7 75 -2 <	5 48 75 -20 FCT	-2 50 447	CTIC 5 -2 75 t	S) H <2		(2 E	I G — 4H SURV OVE — DH	H OLE EY	T P Soil Phuin	EER: SAT	R Ing	א ט	T T <2 IL SU 15	0 4 13 4 E 10 4 10 4	L U PRAC E EN	M E TIOH NGINEE MOIST	G/CC BING SATUB -ATED 48	(VO RAT AT 1/3 WHOLE	IOS BAR <2
931065 832066 042067 833968 932064 832070	1 2 3 4 5	3 3 50 2 4 15		3 50 2 4 15	3 40 2 3 13	3	TR TR 7	97 97 50 98 96 05	3 50 2 4 15	3 40 2 3 13	3	7	97 50 98 96 35		1.00 1.31 1.08 1.03	1. 1. 1.	09 41 17 15		1 3 1	.62 .82 .67	0.79 0.98 0.87 1.06 1.01 0.92	1.0 0.9 1.1 1.1	4 1 4 0 3 1 0 1	.07 .96 .15		1.54	1.65 1.02 1.45 1.57	1.70 2.05 1.50 1.62
3AHPLE NO	HZ NO	(V (원 0 : 9 250 9 -75	75 -2	75 -20 -PCT	0 I 20 -5 0ľ	110L	หส) - <2 ห 50:	A 2- .05 I L	T I T 1 .05-	.00) A P(2 D	S) R IRES P	(\Z) (\N TAH 01-	(8 A) PINE	T I	0 S <2 -C 0 M TS	T HM PR E C	O ACT	C IO N – 15 B A R H 2 O	LE 1/3 BAR	LI) HH - (-1,	HEAR OLE /5 B 0 C	SOIL A3 T	<	LITY) 2 Hn (N) -> UVEN	WHOL	/IN->
832065 837066 832067 832068 932069 332070	2	1 1 25 1 2 6		1 1 25 1 2 6	1 1 20 1 1 5	1	TR TD 3 1	99 99 75 99 98	17	15 19 11 21 21 16	1 1 1 1	16 16 10	45 46 35 49 50 50	15	0.03	13. 13.	97 33 75	5.55 4.03 6.56	4 . 4 .	. 18 . 4 2 . 3 4	1.106 0.789 1.375 0.875 1.462 2.231	2.12.2.2.4.1	0 0 1	5.1 2.9 2.5 2.7 3.7 5.6	4.1 2.0 2.0 2.2 2.3 4.5	3.0 3.3 2.8 3.8	0.32 0.30 0.26 0.33 u.35 u.35	0.30 0.34 0.34 0.36
SAMPLE	H Z N O	¥ II >2 75 -:	0 L ! 20 ? -2 ? >2n:	2 .05 1+3AN	0 [05- -002 D+SI	L LT .002 LT>	VC ((2 H) 5A) C	i P ins- ins-	R A P OP S	C [VP AVD+	1 0 SILT C SILT	я - s г	CL \Y)	(patr In Ethib	11 120 3 6 2 6 13 1 2	50 24 -)	3 \ 40 2 - .45	.,,	76.P 192 18	CLAY LT .002	03.2 01.3 -0.13	721 1721 140 140	S- (. DI HS : - 42	204- JCT - 16405	KG 1'	IN. U 1/3 BA 5BAR A	F H2) K TO INDF (SUIL)
832065 832066 832067 832069 832069 832070	2	3 5 51 5 2 4 1 15 15	1 10	27 43 42	52 51 23 55 54 44	5 4 1 3 3 2	1 1 7 1 1 2	4 6 3 5		17 17	2.0	19 17 16 19 21 21	30	5 4 2 3 3 3		51 54 51 51 51	L.	44.6 45.5 53.0 42.5 12.2 46.9	50 44 54	1.5	4.7 3.8 2.4 3.2 2.6 2.6	4.7 5. 5. 5.	1 1 1					
SAMPLE 832065 832066	0-	- 10	10 VO	10	LAY 4.7 3.8	44.	6 (oc 5.217	,	5/C 4.5 12.0	¥	3/C 9.5 1.4	 ч	0C 6.2 5.8			0-	2 507 - 10 - 51	1, 1; 1 () 1 ()	ט גינ ניגינ ניגינ	1P S 0 9.	5 .0 19	¥ 5 1.5	/c	₹ 0¢ 5.2 5.0			

ECUADOR 307

SAMPLED AS:

S 33PH-325 -007 ECUADOU-SHSS

SAIPLE NOS. 63P2077 - 2083

DATE 11/30/83

U. S. DEPARTMENT OF AGRICULTURE SOIL CONSERVATION SERVICE MATIONAL SOIL SDEVEY LABORATORY LINCOLN, MEDRASKA

	GEN;	EH 7 L	m eti	mas	: 8	UGIN	SERI	NG I	PHACT	I 0115	128	r ca	LCUI	.1 T P. C	· PRO	ក បទរ	DA P	FACT	IOH S	225				.022,	******	•		
SALPLE to:	HZA	(IH		!!O	H-	P E	R C 2 I	3/2	TA	G E 3/4 5	1/4	2 4	รร 10 ยส	G I H 40 2 E	P G 200 9->	5 1 20 <-#10	D C R S CROH	A √ B √ 2	CUMU USDA 1!	JLATI LX 5 _2	72 55 5.	THAN 10 -05	PRACT DIAM 60 60	ETERS 50	<76HB) (HH) AT 10 TILE> 21	BERG LL P	UWI- I PHTY > CU	ALOB- COB-
832077 832078 832079 832080 832081 932082 832083	45 55 65 75	14- 26- 35- 42- 56- 73-	26 35 42 56 73 91	10 201 302 3AB 3AC 40	J	100 100 100	94 92 100 100	10 <i>i</i>) 100	100 100	72 100 100 100	72 160	100 100 100	94 35	58 49 90 93 64 72	25 21 53 56 38 34	11 16 12	10 7	2 8 11 5 3	63 9 97 9 99 9	52 4 53 3 93 8 96 8	7 1 4 7	29 2° 23 19 60 44 62 4° 43 3°	0.46 0.80 4 0.10 7 0.09	0.31 0.44 0.06 0.05	3 0.012 4 0.012 3 0.009 5 0.003 7 0.002 0 0.005 3 0.011		38.6 92.9 38.0 58.7	1.8 2.1 3.6 1.9 2.1 2.5
SA4PL3		>2 2	E H C 50 3 UP -	: . 250 -75	3	75 -20 HOLS	20 -5 5 501	L ()	(2	75 -2 <	l I 75 18 75 -20 -PC f	28A 29 -5 OP C	TI: 5 -2 75	S) 1 (2 1->		SOIL	I : Guille OV C-	G H HOLE VEY	T P SOIL- SMGINI MOIST	ZEH IN	G R D		<2 M . SURV	M PRA EY— OVEN	U H E CTION ENGINEE HOIST	RING	-HAT	BAR <2
832077 332078 332079 332079 332061 332062 532033	2	23 32 1 TR	 		22 23 32 1 1 5	19 19 25 	1 1 1 TR 	1	79 77 68 49 100 49 95	ے 1	19	1 Ca 	3 1 TR 1	78 77 68 99 100 99		1.46 1.36 1.36 1.36 1.36 1.56) 1) 1) 1) 1	.43 .52	1.61 1.60 1.65 1.64 1.64 1.35	1.8 1.8 1.3 1.9	7 7 0 6 8	1.23 1.14 1.28	1.26 1.26 1.36	1.25 1.26 1.27 1.30 1.43 1.66	1.46 1.46 1.63 1.65	1.86 1.98	0.89 0.89 0.91 1.05 0.92 0.68 0.83	1.15 1.32 1.07 0.92 0.69
SAMPLE 90	11 2	>2 2	О И Н 50 UP	L 1 0 L 250 -75	75 -2	75 -20 -PCT	0 I 20 -5	F 1 5 -2 WHO	1 A (MH) <2 LE SO	2- .05	r I 7 1 .05-	0 /3 LT	N B A P(S) P PES P	(C/) - (/N RAT -IO	(R A	f I	. ∪ S <2 C	MM PR/ E C A44- UAC 67	O NCTIO 15 Ra	н — — Н	A Y) LE 7/3 BAR 69	¥HOI <-1/5 15	E SOI BAR		(2 Ha	WROL	Z <2 AH
6 12077 3 11279 6 12179 3 32080 6 3 32082 8 32082		12 17 1 1 k				10 10 13 	1 1 7R T9	1 2 1 1 1 1 3	48 84 33 99 100 99	30 35 27 23 41 38	10 10 3 13 19 15 12	1 1 4 0	27 22 10 ?2	21 20 26 35 26 27 28	11		1 1 1 1	.92 .70 .21 .12	2.00 1.03 1.19 0.97 0.88 1.3)	1.4 1.0 0.8 0.6	6 0 0 1 2 0 8 0 4 0	.333 .370 .329 .106	0.2 0.7 2.d 2.0 0.7 1.5	0.2 0.7 3.0 2.5 1.2 1.9	0.8 3.4 2.0 0.7 1.5	0.8 3.7 2.5 1.2 1.9	0.17 0.16 0.24 0.27 0.16 0.20 0.19	0.18 0.29 0.27 0.16 0.20
SAMPLE		>2 7	11 () 5 2 -2 0P 2	L I 20 : -2 244	S 2 . 0 S . • S A N	0 I 05- .002 D+5I	L LT .002	VC (SA C	и Р и ЭS- и РСТ	R A	1 D 4 P 4 D H A	SILT	'S 'Y	(1) (1)	1027	17.1 E 10 12 E 12 E	MUD 1 SDA	5A90 2 =	3 (L -)5-	2 2	CLAY LT (CA- :L2 t .01m	RES- ST. OHMS	TRICAL) CON- DUCT. MNHOS 2 MM	SALT HG/ KG 1	IN. O 1/3 BA 5BAR A (WHL	F H2O B TO IBDRT
832077 432078 832079 932090 832031 932082 832083	1 2 3 4 5 6 7	3.3	23		58 37 50 60 60 72 72	20 19 18 39 40 27 23	2 2 8 13 5	3 5 7 2 1 2 8	13 15 16 5 3 0	25 24 24 13 19 23 16	21 20 21 23 24 26 22	12 11 6 17 17 13		14 19 19 22 23 14	2 2 3 3 13 5		1. S F F	Ն 51. 5 L Ն	55.7	24. 25. 36. 35.	7 9 7 7 9	2.4 2.7 7.6 11.3 5.0 3.1	7.6 7.0 6.d 5.3 7.3 7.4 7.9					
SAMPLE 9 32077 9 32078	0	LOW - 35 - 67	35	(DI)	5	2.4	SAN 72. 72.	4	0.61 0.49		5/C 30.2 30.4	3	5/0 0.2 0.2		0.6 0.6			ij	5 FOX	14 14	n I. 18	P S,	/C N	5/C 2	0.6			

SUPPLET "HTAUY CHARACTERIZATION DATA

ECUADOR 008

SAMPLED AS:

S SSPH-325 -000 ECUADOR-SHSS SANFLY NOS. 53P2210 - 2214

DATE 11/30/83

U. S. DEPARTMENT OF AGRICULTURE SOIL COMSERVATION SERVICE MATIONAL SOIL SURVEY LABORATOMY LINCOLM, MEBRASKA

GENERAL METHODS: REGINEERING FRACTIONS WERE INCOLARD FOOR USDA FRACTION SIZES

	GEN	ERAL GE	ruops:	PNGI	N K K H 1	CHG P	RACT	IONS	VER	* :1	rcui	. 1															
SAMPLE O.	HZ H	(FI)	903 120	- 1 - 1	R C 2	3/2	T A (i	3/1	4 4 (-N	5 S 10 U H	1 N 40	G 200 H->	20 S .	D L P. S CROS	A V 2 (5->	USD!	10 L.J 10 L.J	LESS LESS -25	THA 10.	E PRA N DI 05 6	CTI AME 0	ONS (< TERS (50 RCBNT	76HM) HM) AT 10 ILE>	ATTER- BERG LL P: <-PCT: 22 2	GBAD UNI- UNI- CU	ATION CUB- CC
832210 832212 832212 832213 832214	25 35 45	0- 1 10- 2 25- 3 35- 5 53- 7	5 A2 6 A3 3 ZC	100 100 100	44	99 00 73	57	97 63 46 100	19 100	97 52 32 100	45 47 23 100	78 34 11 72	18 1 31	21 8 1 14	1'R 15	1 TR 2	43 19 99	81 36 13 94	66 28 4 05	48 21 1 67	35 0. 1419. 127. 51 0.	. 18 . 30 . 122 . 07	0.110 3.339 1.034 0.046	0.006 0.005 0.026 0.398 0.004		34.3 > 100 68.1 20.4	0.9
SAPPLE	H2	(W P. 4 H >2 250 	I G O L E 250 -75 -PCI O	n T 3 75 75 -2 -20 F whol	0 I 20 -3 E 501	P A 1. (H 5 -2	λ (H) − (2 .—>	€ -<7' 75 -2 -<	1 I 5 11 75 -20 PG2 (0 23 23 -3	CT16	;))4 (2)1->		JICE 1/3	1 :- 1U2 0\ 1-	G 40 4 Y 8 8	T T SOLI EXGL HOLS	P 8 L N 2 2 1 T S 1	A H ISG ATUR	0 M	I T < 2 IL SU 15	V S SS SS SS SS SS SS SS SS SS SS SS SS S	VEN C	N E TION HGIHEE HGIST	G/CC HING SATUR -ATED 48	\ \ \ 8AT AT 1/3 \\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	ID) IOS BAH <2 BB
331230 342231 331112 331113 331114	2 3 4 5	3 5 53 77		5 3 53 40 77 54	14	2 5 9	95 47 23 100	53 77	54 	T R 0 14	2 5 9	47 23 100		1.00 1.30 2.20 1.00	2 3 5	1.14 1.42 1.15	1.39 1.69	y . 5	1.62	0.97 0.84 1.05	1.0)5)1 3	1.11 0.93 1.15	1.36	1.55 1.60 1.52 1.65	1.65 1.01 1.52	1.73 2.15 1.52
SA 19	42	V 0 	L U B O L 250 -75	1 4 75 75 -2 -20 PC	5 u 1 20 -5	7 R L (5 -2 VIIO E	A (88) < 2 (8: 30	0 2- 05 11	1 1, 1 05- 102	0 1 17 10	4 d 10 d 10 d	S) R PUES P	(CZ) 13.6 14.6 14.6	(3) (3) (1) (1) (1)	T 1	L O : <2 C SUM :ATS	5 : : MA PI 2 C=: 384: 080	PO RAC: - -	TTON- 15 44 R H2O	L A Y) (1.1 - VH >	HEA IOLE 1/5 5 1 R	R EXT	ENSI51 (15	LITY) 2 MM TN)->	4IH WHOL WHOL WHOL WHOL WHOL WHOL WHOL WHO	. U) .E <2 .MM
8 fau 10 43 111 8 3 2 2 12 8 3 2 3 13 8 3 2 3 14	3	1 2 26 65		2 1 26 20 65 45	12	3	49 74 35 100	15 19 14												2.348 2.70b 3.182 1.550		. 6 . 7 . 5	3.1	3.3 2.7 2.7 2.5	4.6 3.5 3.1	0.25 0.20 0.23	0.21
S inple	u 7	(W % W H >2 75 -2 PCT OF 76 77	I G H O L R 20 2 -2 . >2Mn+	T F S O I 05- 05 .00 SAND+S	R A C L LT 2.002	7 T	0 H -(2 3) 	5 ~ 3 P H DS ~ H P CT (C L R A P DP S	A Y C T V P A S D +	I (SILI C SILI	H T	CL AY	(-TF) (DST) (N (HTL)	₹198 6971 5 6 42 8	181 (420 17 250 A 18-)	7 A M S 2 - - 05 (1	5 .	3 A(H S(LT);- .002 .0P .	CLAY	(P" CA- CL2 -015) (- R IS	RLECT RS- T. D HMS	RICAL) CON- OCT. MHROS NH	SALT HG/ KG 1	T. AMG IN. G 1/3 DA 5BAR A (BHL	OUNTS) F 1120 B 10 IRDRY SOIL)
8 322 10 8 322 11 8 322 12 9 322 13 8 322 14	3	3 3 5 5 53 53 77 77	2 13 23		2 1 TB	4 3 16	11 15 28		19 16 12	14 15 1 16	15 12 2 18	17 3 33	2 1 TH 2		5	iL iL ins is L	70 95.0 48.!) . ; ;	35.5 24.6 4.4 44.5	2.3 1.7 1.1 0.2 2.0	5 . 5 . 5 .	, 1 ,0 ,3 ,2					
5 NHP L Z 8 3 2 2 1 0 9 J 2 2 1 1	0	LOW L - 26 - 63	26 26	CLAT 2.3	5A)	• D	OC 7.146 6.29	6	3/C 28.7 36.7	'i 2	5/0	. 1								I P 18 28			s/C	W OC			

ECUADOR 009

SAMPLED AS:

5 83FH-325 -009 ZCUADOR-SHSS SAMPLE MOS. 33P2091 - 2016

DATE 11/30/03

U. S. DEPARTMENT OF AGRICULTURE SOIL CONSERVATION SERVICE NATIONAL SOIL SURVEY LABORATORY LINCOLN, NEURASKA

GENERAL METHODS: ENGINEERING PRACTICES ARMS CALCULATED PROX USDA PRACTICE SIZES

																	TON 2									
57.918 .0.		(IN) Dept	10H 221 H	R- 3	. H C	3/2 H C	T A	3 /4 5	3/8	1º A	10 U #	40 8 8	G 200 R->	P S S 20 <-:11	U A L P. V S CRONS	2 ->	CUH USDA 1	ULATI LE: 5 .2:	E CUAT	/B PRA IN DI .05 6	- PER O VWRI	NS (< ERS (50 ICENT	10 10 10	ATTER	ONI- PHTY CO	CUE-
832091 832092 832093 832094 832095	35 45 53	10 - 27 - 47 - 54 -	72 3AH1	100 100 P 5 1 100	100 1 99 1 94 1 1 C	98 90 1 T 1 T	96 0 H 98	ษ5 ถ0 S)5 77 H)5 74 J T	94 67	6 1 49 D &	49 26 1 E	2:J 13 8 I i	13 0 1 1 1	3 1 E	91 61 D	84 7: 51 4	2 55 1 29	42 0. 21 0.	.13 0 .95 0	.463	0.003 0.004 0.012		35.2 82.9	1.4
833096			86 JA32		99					47	94	d 1 	47	23	14	5 	91	a4 7:	52	40 0.	14 0	-087	0.003		44.1	1.1
SV.PLE	H 2	>2 25 -U	E I G H O L B O 250 P -75 PCT O 7 29	: 5 75 75 -2 -20 P VHOL	20 -5 E SOI	L (6 5 -2 [L	(2 >	√7 75 -4 √	5 ## -75 -20 PCT	3 F#3 20 -5 OF <	OTIO -2 75	5) 0N (2	ı	Soil 1/3	I G WHO SURVI OVE: - Day	и Б 1 О Г Б Н	T P SOIL BUGIN BUIST	ERRING SATUR	U N so	I t <2 IL SU	-D BAEA HH A O	L U PRAC E En	H E TION HGINZE MOIST	G/CC) RING A SATUB -ATED 48	(VO BAT T 1/3 WUOLE	IOS BAR <2
8 J2091 8 J2093 8 J2094 8 J2095 8 J2096	2 3 4	68 - 3 -		1 6 5 33 20 69 50 3 2 6 2	6	1	94 67 32	69 69 69 69	5 20 50	T a	1 7 4 1	99 94 67 32 97		1.0. 1.3; 2.1; 1.2;	2 1. 2 1.4 1 2 1.5	11 48 20	1.46	1.64	0.96 0.96 1.05	1.0	5 1 7 1 5 1	.07 .21	1.43	1.52 1.61 1.65 1.75 1.76	1.60	1.70 1.52
31/962	H Z	>2 25:	D L II H () L D 250 P -75	r: 75 75 -2 -20 PC	20 20 -5 108	(5 -2 VHOL	(2 (2 (8 SO)	2- .05 I L	r 1 .05- .012	/3 LT .00	1 A PC 2 D	S) R = L 3 R ('	(C/) (/N +AT -TO	R A) (I) PINE CLAY	T I () S <2 . -C III IS	TO 8.4 PRI 8 C 844-	0 (10 k 15 h 18 h 112 c	LAY) (LI - WH 1 15	NEAR OLE /5 B OR -	EXT. SOIL AR TO Ven	2 4 5 1 B C < 0 (I N / 15	2 NA IM)->	N NOT	E <2 BH /IN->
832091 832092 832093 832094 832095 811390	1 2 3 4 5 6			1 2 2 16 10 55 40 1 1 3 1	3 7	1 3 7 1		20 23 24 22	12 15 9 1 21 17	1 1 2	17 10 20 14	42 44 34 40 37	17 16		12.8 32.0	18 10 10	4.68 7.40 20.00	3.68 5.35	0.749 0.882 2.400 0.390	2.	2 9	5.6 2.9 3.9 1.6	4.2 2.3 3.7 1.4 1.3	1.0	0.29 0.32 0.25 0.31 0.27	0.33
SAMPLE 30	HZ NO	W H >2 75 -2 PCT OF	I G H O L P 20 2 -2 (>233+5	F F S O I 05- D5 .00: SAND+S:	R A C L LT 2.002 (LT>	7 I	0 H <2 H SAH C	S - I F I DS	C L R A F	A Y C T VP	I O SILT C SILT	R 7 N - S P	P.) CL 11	(-TEX (DETE IH FIFT D	TO NE- NB I HE BY PSD 2 HII-	-) D A	(P S SAND 2- .05 (PC	D 4 (SILT .05-	LT LT LT 0.02	(PII) CA- CL2 -01H ((-81 RE: IST. OH!	L#CTF 5- (- DC 15 8	ICAL) CON- ICT.	(CUMULT SALT IG/ 1 KG 15)	. AMO IN . O. /3 BA	UNTS) P H2O R TO IRDRY BOIL)
832091 832092 832093 932094 832095 832096	q	1 1 6 6 33 33 60 69 3 3 6 6	1 ! 13 ! 18 !	50 39 54 40 47 19 31 1 50 47 57 37	3 1 17 17 4 5	2 3 10 22 1 3	9 14 35 6 0	13 16 32 10	19 17 7 18		15 12 2 16	24 29 17 2 32 26	f 9 4		FSL TSL COS FSL PSL		56.4 55.1 51.4 46.6 40.4 57.2	40.3 32.6 3.3 46.5	3.4	5 5	3					
SAMPLE 832091 832092	0-	26	ON DIP 26 26 40 14	CLAY 7.1		D 2 6	OC - 168		S/C 7.9 6.4	7	5/C	7	0C 5.2			0 - 0 5	LOY 25	Say I	of P :	5/C .9	¥ S/ 7.9	/C 6	¥ 0C			

CHARACTERIZATION DATA

ECUADOR 010

SAMPLED AS:

S 83PN-325 -010 ECUADUR-SHSS

SAMPLE NOS. 43P2220 - 2225 DATE 11/30/83

U. S. DEPARTMENT OF AGRICULTURY SOIL CONSERVATION SERVICE HATIONAL SOIL SURVEY LABORATORY LINCOLN, MEBRASKA

GENERAL HETHODS: SHOTHERRING PRACTICHS WORR CALCULATED FROM USDA FRACTION SIZES

		- INAL			Dia.													NA.				50 100				Miles.		20.	
SAMPLE NO.	HZN NO.	(I)		HOR 120	1000	RS	C E	N 7	1 3	1 4	3/8	2 4	5 5	1 N	G 200	20	L E 5 Chon	2 Y	1.	.5	LPSS -25 -4H-	.10 .0)5 6 -> <-	AMET O -Pei	TERS (10 10	<-PCT	UNI- I PHTY > CO	VTUR CC
832220 832221 832222 932233 832224 8323.15 832216		11-	21 33 46	2112	10	00 5	19	98 98 98 94 90	97 97 97 97 97 90 84	96 96 96 90 87 40 43	16 96 16 16 16 75 73	95 95 95 95 95 84 76 42	91 42 31 10 78 69	71 75 72 62 58 52 79	33 33 35 20 24 27 49	15 15 13 a 9 11 27	8 7 3 4 4 5 13	2 2 1 1 1 3	86 36 33 73 63 86	75 79 76 68 62 54 81	60 63 60 45 47 45 72	40 41 25 29 32	25 0. 27 0. 14 0. 17 0. 20 0.	22 (25 (39 (46 (78 (78 (.149 .154 .290 .289	0.008 0.008 0.028 0.028 0.028		15.2 21.5 54.4	
SAATUR		>2 2	H O 50 2 UP -	50 75 C7 0		75 2 20 -	1 L 20 -5	-2	1- <2 ->	~77 75 -2 <	5 aa 75 -20 20T	20 20 -5	-2	<2 1.1->		SIIL 1/3	SUN OV	EH NCTR HOTE	T 1 SOII ENGI: NOIS:	E SA SA	R	50	T T-<2	W C	PRAC	MGINES LION	G/CC RING SATUR -ATED 48	HAT 1/3	BAR <2
632210 932221 932212 832273 832214 832216		9 10 22 31		=======================================		4 4 4 1.3 20 7	1 1 1 1 3 4 1	3 4 5 6 7	91 92 91 90 73 69	4 9 10 22 31 11	1 1 1 1 1 1 7	1 1 1 3 4 1	4 3 4 5 6 7 3	91 92 91 90 73 69		1.5 1.5 1.5 1.5 1.3	3 1 3 1 7 1 3 1	.23 .40 .35	1.64	4 1 4 1	.85	1.18 1.21 1.10 1.11	1.1	4 2		1.52 1.46 1.43	1.73 1.75 1.69 1.69	0.93 0.99 1.25	1.25 1.19 1.41 1.39
SATELE		West Control	0 V II 50 2 UP -	L U 50 75	4	# 3 1 75 3 20 -) I 20 -5	-2 -2 -4	(11) (2)	1- 11-	.09-	L1	1 D	14KS	141	PIJ PIJ SLA	2 -	0 8 -42 50.2 (A78	35. 71 8 C 910- 040	ract - :	10 N-	LE 1/3	- V:: (-1 15	OLE /5	SOIL SAR T OVEN	ENSIB! O (IN) 15 DAH 72	(2 AH- (1H)-> OVEN -DHY 73	SOIL CIN	D) E <2 HH /IN-> 75
332210 8321.1 832212 832225 832225 832226	1 2 3 4 5 6 7	5 0 12 16	<u>-</u>	 	5 5 6 12 16 5	2 2 2 7 10 3	1 1 1 1 2 2 1	1 2 3 3 4	95 96 95 94 83 84 75	11 11 4 1 25 22	13 11 14 1 9	1 1 1 1 1 1	43 22 13 13 21 19 24	32 27 31 31	11 12 11		3 4 2 4	.16	4.2. 2.1. 1.6. 3.76 4.3 3.6. 2.4	3 1	.19	0.615 0.286 0.257	0.	5	0.7 0.5 0.8	0.A 0.u 0.9	0.8 0.6 0.9	0.23	0.29 0.27 0.31 0.26
SAMPLE NO	42 NO	>2 7 PCT	11 0 5 2 -2 0P >	0 2 -2 . 2HH+	T S O01	I L. 5- L1 002.0	 002 r>	YC (SAN	I P I DS - M PCT	R A	V ?	SILI) - S F	CL AY	14 14 131 1	ERNI D P	Y SDA H-1	2-	.0	11.T 5- 00?	LT .002	CL2 .01%	15	D IHS - <2	COM- UCT. UMUS MM-	(CUMUL SALT MG/ KG 1	1/3 BA 50AB A (WHL	R TO
8 32220 8 32221 8 32222 0 32223 8 12224 8 32225 8 32226	1 2 3 4 5 6 7		9 8 9 10 22 31	5 6 9	69 66 76 62 50	24 23 25 14 16 19 39	3 2 2 1 1 1 3	6 4 5 9 6 8 3	12 11 12 16 15 14 6	17 10 19 26 19 14 10	23 25 22 23 23 19 20	16 17 15 12 16 18 17	12 11 10 7 10 13 17	14 14 18 0 11 15 27	3 2 3 1 1 2 4		S	L S S S L SL	71 73.0 70.4 13 78 70.6 54.0	1 2 1 1 2 2 2 2 2 2 2 2 2	5.5 4.7 7.1 5.1 0.5 7.0 2.5	2.7 2.3 2.5 1.0 1.3 2.1 3.5	5. 6. 6. 6.	0 1 9 0 1					
5 AMPLE 8 32220 8 32221 8 32222	10	LOW - 10 - 28 - 40	10 10 28 40	10 18	2	.7	71.8 73.0 70.4	0	OC 1.871 1.853	3	S/C 26.6 31.7 28.2	2 2	5/0.6		00 2.0 1.6 1.2			0	P LO: - 10 - 23			0 26	5/C -6 2 -7 2	6.6		w oc 2.9 2.0			

ECUADUR 011

SAMPLED AS:

S 83FW-325 -011 ECUADOR-SMSS Sample NOS. 83P2135 - 2142

DATE 11/30/83

U. S. DEPARTMENT OF AGRICULTURE SOIL CONSERVATION SERVICE NATIONAL SOIL SURVEY LABORATORY LINCOLN, NEDRASKA

GENERAL SETHOUS: ENGINEERING FRACTIONS 4286 CALCULATED FROM USDA PRACTION SIZES

				mica																		3.142.5	North Control			State Line		La Contra	Marie Line
SAURLS . (8	HZN NO.	(III		noa Izo	-	P 3	2 I	E N	T A	3/4	3,/9	PA	10	I N	G 200 R->	\$ \$ 20 <-MI	I E S CROI	V K	1	5	.25 -88-	. 10 .	DI 05 6	METE 5 -PERC	RS (1	10 10 LE>	LL P	- GRAD UNI- I PHTY > CU 3 24	CUR-
832135 832136 832147 832143 832140 932140 932141 832142	18 28 38 48 58 68 78 88	10- 22- 33- 47- 52- 61-	33 47 52 61 70	11 A2 A3 2C1 3A11 3A12 3A13 4C2		100 100 100 100 100 100 100	99 95 90 10 99 96 46	94 99 92 42 99 99 93 77	98 87 72 98 98 88 63	47 97 94 64 93 98 45	47 97 84 63 93 93 93	97 97 83 62 98 98 98 34 47	47 97 31 36 98 97 11 42	92 92 63 28 90 66 51	57 60 36 3 44 47 18	13 35 20 2 24 26 11	16 17 9 1 11 12 5	4 4 2 1 3 3 2 1	97 97 79 49 37 95 75 36	94 94 72 33 94 90 53 22	83 85 57 12 77 74 32 8	66 41 3 50 54	51 0. 30 0. 3 3. 36 0. 38 0.	08 0. 28 0. 54 1. 14 0. 13 0.	048 166 154 101 084 404	0.003 0.003 0.006 0.205 0.004 0.017			0.8 1.5 0.3 1.7 1.3
SALEUR	HZ NO	>2 2	H C 50 2 UP -	250 -73 PCT 0	75 -2	75 -20	0 [20 -5	-2 L	(2	₹7 75 -2 	5 14 75 -20 PCF	774 20 -5	-2 75	<2 1.1->			SU !	HOLE RVEY	T (SOII E.(GIA #0157	EER!	NG UR	50	<2	MM P	RACT - EI H I	TION IGINEE IOIST	RING) (VO 	IOS
332116 332116 832117 031118 832119 931110 131111 932112	1 2 3 4 5 6 7 8	3 19 44 2 3	=======================================		3 3 19 44 2 3 19 58	3 16 36 2 2 15 47	TR 1 1 6	TR 13 2 6 PR 1 3 5	97 97 91 56 98 97 81 42	3 17 44 2 3 19 53	3 15 30 2 2 15 47	73 1 2 74 1 0	TR TR 2 6 FR 1 3 5	97 97 81 56 93 97 31 42		1.0 1.1 1.0 1.1 1.0 1.1 1.2 1.2	4 3 4 1 7	1.20	1.33	}	.49 .65 .70 .71 .75	0.76 1.02 0.99 1.12 1.19 1.13	1.0 1.0 1.1 1.2 1.1	5 1. 5 1. 5 1.	07 06 16	1.31 1.45 1.44 1.47 1.54 1.43	1.47 1.64 1.62 1.70 1.74 1.70	2.40 1.55 1.35 1.32 1.19 1.09	1.60 1.68 1.37 1.23 1.35
SA total	H2 20	>2 2	50 2 UP	250 -75	75 -2	75 -20 -PC 1	23	1 (-2 1:101	(EP	2- .0,	7 1 .J5-	1.73	1 A	R	RAT)	Y	<4		ACT		LE 1/3	- 44	DLE S /5 BA OV R -D	OIL R TO PH	<	LITY) 2 Md IN) -> OVEN -OBY 73	MHOL	/IH->
33.116 33.116 33.116 832139 832140 832141 832142	1 2 3 4 5 6 7 8	1 5 30 1 1 9			1 1 1 1 9 43	1 1 7 25 1 1 7 35	1 1 TR 1 4	TR 1 1 1 1 4	97 92 74 99 97 91 57	14 11 34 27 27 31 29	11 12 1 14 16 6 1	1 2 1 1 1 1	16 13 13 13 23 20 25 26	55 41 42 34 39 27	12 14 15			1.21	6.47 4.30 4.30 1.40 2.37 2.33 2.56	2.	.52 .07 .56	1.872 0.343 0.767 0.444 0.222 0.480	5. 1. 1.	3 1 7 2 9 1 5 0	.5	5.4 1.3 2.0 0.9 0.6 0.9	7.3 1.6 2.3 1.2 0.6 1.2	0.38 0.3' 0.3\ 0.27 0.27 0.22	0.2/ 0.2/
SAMPL3	nz 110	>2 7 PCT	3 0 5 2	LR	5 05 SAN	0 I 05- .002	L LT .002 LT>	YC	c s	H DS- H PCT	R A	C T	STI.1) W -	CL	(DET IN IN FINL	ERMI	THED Y Y Y DE	2- .05	.05 .05	LT -	LT	CA- CL2 .014	RES IST. OHM	- DI	CON- UCT.	SALT MG/	T. AMO IN. O 1/3 BA 5BAH A (VIIL 99	P H20 R TO IRDRY SOIL)
832135 832136 832137 832131 832139 832140 832141 832142	1 2 3 4 5 6 7 8	2 3 19	3 19 44 2 3 19 58	3 8 1 3	50 49 52 54 64 60 67 40	47 48 29 2 34 37 13 2	4 5 2 1 3 3 2 1	TR 1 3 14 1 2 8 14	2 3 9 23 3 5 22 34	12 10 19 38 18 17 32 33	21 20 21 16 29 21 15	16 17 14 1 15 16 6 3	17 17 14 1 12 13 6	32 33 22 2 22 25 11	3 3 3 1			FSL FSL FSL FSL FSL FSL FSL FSL FSL FSL	47.5 47.9 62.6 45.3 63.7 60.4 41.4 94.7	31 36	.5	3.9 4.6 3.0 1.5 2.7 2.7 2.5 1.2	5.	4 5 5 9					
5AMPLE 832135 832136	0	LOW - 25 - 57		DIP 3 25 0 15	THE R	LAY 3.9 4.6	5AN 49. 47.	6	OC 7.24 3.52		5/C 12.7 10.4	. 1	5/0 2.7 1.9	¥	nc 7.2 5.8				P LOW - 25	L. 16	W D	17 : 9 12	7 1:	W 5/	c	W OC			

ECUADOR 012 SAMPLED AS:

5 63FN-325 -012 ECUADOR-SHSS

SAMPLE NOS. 63P2167 - 217

DATE 11/30/83

U. S. DEPARTMENT OF AGRICULTURE SOIL CONSERVATION SERVICE NATIONAL SOIL SURVEY LABORATORY LINCOLN, NEBRASKA

GENERAL METHODS: ENGINEERING PRACTIONS JARK CALCULATED FROM USDA PRACTION SIZES

		CBAL					COMI	. 70	TANCI	LUNI	4/5		LCU	LALK	, FEL) n U S		PHACI	LON	312	.5		-30		100	-			
SY INTE	HZN NO.	(IA	0	1101	-	3 2	H C		T A 1 U E 4	3/4		4	S S 10	1 N	G 200	20	I P.	V E	USD 1.	.5	.25	.10	.05	DIAM 60	ETERS 50	TILE-	> <-PC	G UNI PI PHT T> CU	- COR- Y VTUR CC
8.xc167 332164 832169 812170 332171 932172 312173	13 25 35 45 55 63 75	20 - 33 - 40 - 51 -	40 51 54	A1 03 2AB 3BT: 30TE 46TE 48T3	11 12 13			100 100 99 99 99 100 100		100 100 97 97 93 100 100		100 130 97 97 98 100 100		99 96 97 90 100 99	73 56 63 74 74 45	38 26 32 45 52 70 10	19 12 15 26 37 59 71	3 13 27 52	37 97	99 100 97 97 98 100 99	97 96 95 96 97 99 98	81 66 72 82 80 89 92	43 50 63 66	0.08 0.07 0.04 0.03 0.01	0.06	2 -	4 3 1	19. 21. 19. 34. 71. 17. 5.	1 1.8 8 1.3 7 0.9 4 0.5 3 0.3
SAHLE		<	150 :	250	75 -2 -	5 75 -20 10L&	0 I 20 -5	-2 L	(2 >	75 -2 <	75 -20 -PCT	20 -5	CT I	<2 11->	39	SOIL	501	RVEY	SOL ENGI HOIS	NEER T SA	ING		110	<2 H SURV 5	A PRA	SACINE-	G/CC SATUR SATUR ATED 48	AT 1/	T105 3 8AB 2 <2 8H
83:163 83:163 83:164 83:170 35:2171 83:172 83:173	1 2 3 4 5 6 7	3			3 3 2	3 3 2			100 100 47 97 93 100 100	- 3	3 3 2			97		0.9 0.7: 0.di 0.9: 1.4 1.2	H (4 (4 (4 (4 (4 (4 (4 (4 (4 (4 (4 (4 (4	1.02 0.03 0.09 0.06 1.52 1.49	1.3 1.1 1.2 1.3 1.7 1.6	7 1 7 1 6 1 6 1 6 1	.57 .43 .52 .56 .dd .75	0.91 0.76 0.32 0.86 1.40 1.21	0 0	.98 .01 .86 .92 .45 .31	1.02 0.03 0.87 0.94 1.51 1.49	1.34 1.17 1.25 1.34 1.76 1.63	1.57 1.49 1.51 1.55 1.87 1.75	1.91 2.40 2.15 1.94 0.60 1.19	2.40 2.23 2.01 0.89 1.19
SALILII		>2 2	50 UP	u L 250 -75	75 -2 -	75 -20 -PC1	1) I 20 -5	-2 1HOI	(44)	 	.05-	L:) A	HES	-10 H/5	(A A)	ii ·	< 2	13 P 3 G- 510 57	4 ACT	C 10 N- 15 5A R H2O	L A 1 1/3 3AR 69	- (4HOL -1/5	E SOI	to (IN	ILITY <2 AM- /IN)-> OVEN -DHT 73	SOIL	LE <2 Hn
83.167 81:161 81:169 81:170 93:171 83:173	1 2 3 4 5 6 7	1 1		=======================================			=======================================	=======================================	100 100 99 19 19 100 100	14 17 15 12 17 9 6	14 13 15 17 21 13	1 1 4 1 4 2 4 2 4 2 4	11 12 25 40 10 7	43 43 46 37 47 50	10 11			1.38	3.1 5.3 0.9 0.3 0.2 0.2	2 1 10 5 0	.52	0.557 0.636 0.588 0.169 0.096		2.5 1.7 1.6 1.5 1.2 2.7 2.3	3.9 2.1 1.9 2.2 2.5 7.2 8.4	2.5 1.3 1.6 1.5 1.2 2.7 2.3	3.9 2.1 2.0 2.2 2.6 7.2 8.4	0.24 0.25 0.29 0.32 0.15 0.15	0.25 0.29 0.33 0.15 0.15
SAMPLE		>2 7	H 0 5 2 -2 OF 2	G H L F 10 2 -2 .	S 0 0 05 .	F R 5-	A C L LT .002	7 I	<2 M	N DS-	R A	C T VF AND+	SILI	H -	CL	(-1 E. (5 ET) (5 ET) (7 ET) (2 1	ATTA YUE	.05		14.1	ULAT LT .002 201-)	CF .0	1- 1: 1H (TES- ST.	CON-		IN. 1/3 3 158AR (VIIL	AR TO
832167 832163 832169 832170 832171 832172 832173	1 2 3 4 5 6 7	3 3 3	3 3 3		59 49 39 44 42	50	8 3 14 36 110 188	TR 1	TR TR TR TR	2 3 2 1 1 1 3	17 31 25 17 24 21 18	23 24 24 22 20 10 15	25 18 20 21 20 21 20 21	34	8 3 4 15 37 110 188		V V S. C	FSL FSL FSL	39.5 56.6 48.6 35.2 32.7 20.6 13.4	1 1 5 4 7	3.5 0.1 1.0 1.6 0.2 7.7	7.0 3.3 3.4 13.0 27.1 52.3 65.3		6.1 5.4 5.3 6.3 6.2					
SAMPLE 832167 832168	0-	LOW - 10 - 52	LON 10		7	AY 1.0	5AN 39. 56.	5	oc 4.33 1.65		5/C 5.6 17.2		5/0 5.6 4.3		oc 4.3 2.3			U	P LO 4 - 10 - 52	1	ON D.) 5	5/C .6		5/C	# nc 4.3 3.1	· ·		

ECHADOR 013

SAMPLED AS:

S 13PH-325 -013 ECUADOR-SHSS SAMPLE NOS. 03P2101 - 2185

DATH 11/30/63

U. S. DEPARTMENT OF AGRICULTURE SOIL CONSERVATION SERVICE NATIONAL SOIL SURVEY LABORATORY LINCOLN, MEBRASKA

GRARRAL ARTHODS: ENGINEERING PRACTIONS WERE CALCULATED FROM USDA PHACTION SIZES

	GRASHAL SELECTIONS: SEGRECATIONS ASKS CALCULATED FROM OSDA PRACTION 5/255																											
SVIDLE		(IN DEP		HOR- 17.05	P E	I	3/2	T & 1 H E	G E 3/4 5	3/8	P I	S S 10 U M	1 # 40 3 B	6 200 H->	5 5 20	D I E 5 Can	λ γ ε 2 35->	# CTO # SEC 1.	MUL.	LESS LESS .25	CO B	/E A B . 05	TDART DIAN 00 4P	IONS (< ETERS 50	(88) AT 10	ATTER- BERG LL PI <-PCT> 22 23	DNI- PMTY CU	CON-
83 118 1 8 3 118 2 8 3 118 3 8 3 118 4 8 3 2 18 5	18 28 33 45 55	12- 22-	30 42	. 12 23 CB 2 23 CB 2	100	114	100			103					92		70 a5	98 98 100 100	100	97 97 98 100	94 99	68 78 90 97	0.03	0.022		•	20.8 39.0 5.2 3.9	
SA WLE		>2 2	II 0 50 2 UP +	L 2 50 7 75 •	II T 5 75 75 -2 -20 7 VIIOL 19 30	: 0 I -5 : 20	-5 -5 7	<2 >	-<7 75 -2 <	5 34 -75 -10 PCT	90A 20 -1 08 C	.0013 -2 .75	3) 14 2 :1->		SOLL	0 20 K I	G B WITO LE RV F Y VEN	T SOI FSGI	P E L (NPP) (1 S	R	U N S(1/3	110	T 4 ~ < 2 M SUR 4 15	O L U M PRAC M YS	S K KOLT: SSBIDN: TZION:	G/CC) BING A SATUH -AFED 48	(VO RAT T 1/3 WHOLE	8AR <2
8 J 2 19 1 8 J 2 18 2 8 J 2 18 3 8 J 2 18 4 8 J 2 18 5	1 2 3 4 5				3 3 2 2 2 2			98		2			97 98 98 100 100		1.2 1.3 1.1	2 6 4	1.28 1.47 1.42	1.6	1 73 50	1.76 1.45 1.71	1.2	? 5 4	1.24 1.38 1.21	0.94 1.27 1.46 1.42 1.37	1.61 1.73 1.66	1.51 1.75 1.84 1.71 1.65	1.17 0.95 1.32	1.19 0.96 1.32
SVa La	ΗZ	>2 2	4 H -	0 L f 50 7 75 -		5 0 20 -5	I L 5 -2 VHO	(MM) <2 Le so	2- .05 DIL	7 1 .05-	/3 Lr .00	a A Po 2 D	R 2885 ? >	148 (01-	PI4 CLA	γ Έ	<2 C	44 E 2 C- 4114	PRAC I- IC	TION- 15 34k H20			¥HOL <-1/5 15	E SOLI	< 10 (I#/ 15	LITY 2 dd [IN] -> OVEN -DNY 73	SOIF MHOT	2 <2 88
37 (181 3.2192 832193 832184 832185	1 2 3 4 5	i 	 		1 1 1			49 49 49 100 100	14 13 4	18 25 12 7	14	15 12 5	41 34 37 52 55	11			4.20 2.05 0.45 0.20 0.22	1	3) 30 21	1.J3 0.59 0.51	0.17 9.07 0.10	2 1 9	3.1 0.8 0.7 2.0 2.5	4.6 1.6 7.6 7.6	3.2 0.8 0.7 2.0 7.5	1.6 2.6 7.6	0.24 0.18 0.08 0.12 0.12	0.18 0.09 0.12
SAMPLE		>2 7	H 0 5 2 -2 07 >	L E 0 2- -2 .0 2HH+S	5 0 I 05-	L LT 2.002 ILT>	 2 VC	-<2 H SA C	ส P . ม กร - ม . P CT	7 A F OP S	C T VP AND+	I C SILT C SILT) W - TS P	CL 4Y	(DET IN PIEL	2R# 1 0 1 C2 1	IYPD BT PSDA	2-	0 : PCT	: ILT 13- 002	CLAS LT .003	r (c . c	CA- L2 1 D14 -	REJ- ST. (DHMS <2	CON- DUCT. MHHOS MHHOS	(CDMULT 'ALT 'AG/ 15 'KG 15 ')	O .HI AU CV BAR A BHL;	P H2O R TO IRDRY SUIL)
832191 832182 832193 832104 932135	1 2 3 4 5	3 2 3	3 2 3	3 3 3	3 64 5 63 1 66 4 66 2 78	15 56 233	TR 1	TR TR Z	1 1 7 5	12	23 22 18 14 13	25 23 21 22 22 33	41 45 43	11 15 57 233 558		:	SICL SICL SIL	10. 20. 10.	.7 : .3 :	54.5 56.2 13.5 13.7	13. 36.5 70.0	1	5.9 6.2 5.6 6.3					
SAMPLE 832191 832192	ō.	LOV - 30 - 55	30	30	CLAY 10.0 13.1	30	. 5	oc 3.35 1.20	В	s/C 3.0 2.3		5/0 3.0 2.3	; W						iw 1	LO# D	1P 8 .	3/0	с и 3.	s/c 0	¥ 00 3.4			